

# TENSOR 2-PRODUCT FOR $\mathfrak{sl}_2$ : NEGATIVE HALF

MATTHEW MCMILLAN

ABSTRACT. We show that the abelian tensor 2-product construction for 2-representations of  $\mathfrak{sl}_2^+$  that was given by the author in a recent paper also gives a 2-representation of the full algebra  $\mathfrak{sl}_2$  when the inputs are 2-representations of the full algebra  $\mathfrak{sl}_2$ . As in the previous paper, the 2-product is given for a simple 2-representation  $\mathcal{L}(1)$  and an abelian 2-representation  $\mathcal{V}$  taken from the 2-category of algebras.

## CONTENTS

1. Introduction	1
1.1. Background and motivation	1
1.2. Result	2
1.3. Outline summary	3
1.4. Acknowledgements	3
2. Background: full action	3
3. More bimodules	4
4. Adjunction $(\tilde{E}, \tilde{F})$	10
5. Maps $\tilde{\rho}_\lambda$ are isomorphisms	11
5.1. Some tensor products in $C\text{-mod}$	11
5.2. Maps $\tilde{\rho}_\lambda$ : computation	20
5.3. Maps $\tilde{\rho}_\lambda$ : isomorphisms	38
References	48

## 1. INTRODUCTION

1.1. **Background and motivation.** This paper is the second part in a series by the author, starting with [McM], about an abelian tensor 2-product for 2-representations of Lie algebras. The 2-product is designed with a view to the program by Crane and Frenkel [CF] to develop a higher representation theory in order to upgrade known  $3d$  topological invariants to  $4d$  invariants, such as the TQFT of Witten-Reshetikhin-Turaev [Wit, RT]. The notion of 2-representation was developed in [CR, Lau, Roua, KLa, KLb]. A general definition of tensor 2-product of 2-representations of Kac-Moody algebras in the setting of  $\mathcal{A}_\infty$ -algebras is expected in [Roub]. A related tensor 2-product for the case of  $\mathfrak{gl}(1|1)^+$ , which does not involve the homotopical complications present for  $\mathfrak{sl}_2^+$ , is applied to recover part of Heegaard-Floer theory in [MR]. The reader will find further discussion in the introduction to [McM].

In [McM], the present author gave an explicit abelian model for a tensor 2-product  $\mathcal{L}(1) \otimes \mathcal{V}$  for  $\mathcal{L}(1)$  a certain simple 2-representation of  $\mathfrak{sl}_2$ , and  $\mathcal{V}$  a given abelian 2-representation of  $\mathfrak{sl}_2^+$  taken from the 2-category of algebras. It is natural to ask whether the product category has the structure of 2-representation of the full algebra  $\mathfrak{sl}_2$  if one assumes that  $\mathcal{V}$  also has this structure. This paper gives a proof that this is indeed true.

**1.2. Result.** Let  $\mathcal{U}$  be the strict 2-category associated with the enveloping algebra of  $\mathfrak{sl}_2$ , as given in Rouquier [Roua] or Vera [Ver, §3.2]. See [Roua, §5.1.1] for the definition of (integrable) 2-representation of  $\mathcal{U}$ . Since we only work with 2-representations of  $\mathcal{U}$  and not  $\mathcal{U}$  itself, we do not need a precise definition of  $\mathcal{U}$ . We work with 2-representations in the 2-category of algebras, bimodules, and bimodule maps.

As in [McM, §1.2], let  $A$  be a  $k$ -algebra for a field  $k$ , let  $E$  be an  $(A, A)$ -bimodule, and let  $x \in \text{End}(E)$ ,  $\tau \in \text{End}(E^2)$  be bimodule endomorphisms satisfying the nil affine Hecke relations:

$$\begin{aligned} \tau^2 &= 0, \\ \tau E \circ E \tau \circ \tau E &= E \tau \circ \tau E \circ E \tau, \\ \tau \circ E x &= x E \circ \tau + 1, \quad E x \circ \tau = \tau \circ x E + 1. \end{aligned}$$

This data determines a 2-representation  $\mathcal{V}$  of  $\mathcal{U}^+$ , the monoidal category associated to the positive half of the enveloping algebra of  $\mathfrak{sl}_2$ .

Now assume that  $\mathcal{V}$  has a weight decomposition  $A = \prod_{\lambda \in \mathbb{Z}} A_\lambda$  (cf. [McM, §4.3.1]). The data of  $(E, x, \tau)$  will extend to determine a 2-representation of the full 2-category  $\mathcal{U}$  when the functor  $E \otimes_A -$  admits a right adjoint functor  $F$  such that certain maps  $\rho_\lambda$  are isomorphisms in each weight  $\lambda \in \mathbb{Z}$ . The maps  $\rho_\lambda$  are determined by the data of  $(E, x, \tau)$ . See §2 below for the definitions.

**Theorem 1.1** (Main result). *Suppose  $(E, x, \tau)$  gives the data of a 2-representation of  $\mathcal{U}^+$  for the  $k$ -algebra  $A$  having a weight decomposition. Define the  $(A, A)$ -bimodule  $F = \text{Hom}_A({}_A E, A)$  right-dual to  $E$ . Suppose  $E$  has the following properties:*

- ${}_A E$  is finitely generated and projective,
- $E^n$  is free as a  $P_n$ -module,
- $E$  and  $F$  are locally nilpotent (i.e. the 2-representation is integrable),
- The maps  $\rho_\lambda$  defined in the canonical way (see §2) are isomorphisms for each  $\lambda \in \mathbb{Z}$ .

Let  $C$  be the  $k$ -algebra,  $\tilde{E}$  the  $(C, C)$ -bimodule, and  $\tilde{x}$  and  $\tilde{\tau}$  the endomorphisms defined in [McM] in terms of  $A$  and the data  $(E, x, \tau)$ . Note that  $\tilde{E}$  is locally nilpotent. Let  $\tilde{F} = \text{Hom}_C({}_C \tilde{E}, C)$ . Then:

- The canonical unit  $\tilde{\eta}$  and counit  $\tilde{\varepsilon}$  arising from the duality pairing give an adjunction  $(\tilde{E}, \tilde{F})$ ,
- ${}_A \tilde{F}$  is finitely generated and projective, and  $\tilde{F}$  is locally nilpotent,
- The new maps  $\tilde{\rho}_\lambda$  defined in the canonical way using  $\tilde{x}, \tilde{\tau}, \tilde{\varepsilon}, \tilde{\eta}$  are isomorphisms, so:

- The collection  $(\tilde{E}, \tilde{F}, \tilde{x}, \tilde{\tau}, \tilde{\varepsilon}, \tilde{\eta})$  gives the data of an integrable 2-representation of  $\mathcal{U}$  for  $C$ .

We emphasize that for  $\mathcal{V}$  a 2-representation of  $\mathcal{U}^+$  given by the data  $(E, x, \tau)$  for  $A$ , the fact that  $\mathcal{V}$  has an action of the full 2-category  $\mathcal{U}$  is equivalent to a property of that data: namely that the canonical commutator maps  $\rho_\lambda$  are isomorphisms. When this holds, then (according to the theorem) the maps  $\tilde{\rho}_\lambda$  of the product are also isomorphisms. So the new data  $(\tilde{E}, \tilde{x}, \tilde{\tau})$  for  $C$  inherits the property of extending to an action of the full  $\mathcal{U}$ .

**1.3. Outline summary.** The paper is organized as follows:

- In §2 we discuss the adjunction and the maps  $\rho_\lambda$  that are needed to define the extension of a 2-representation of  $\mathcal{U}^+$  to a 2-representation of the full 2-category  $\mathcal{U}$ . See [McM, §2] for additional background theory and notational conventions.
- In §3 we define and study some bimodules, giving concrete algebraic models for them in the manner of [McM, §3.2].
- In §4 we consider the right adjoint to  $\tilde{E}$ , namely  $\tilde{F} = \text{Hom}_C({}_C\tilde{E}, C)$ , and we show how to describe it concretely using the ‘ $D^b(B)$ ’ side of the Rickard equivalence constructed in [McM, §3.3.2].
- In §5.1 we compute explicitly the tensor products needed to compute explicit formulas for  $\tilde{\rho}_\lambda$ , namely the products  $\tilde{E} \otimes_C \tilde{E}$  and  $\tilde{E} \otimes_C \tilde{F}$  and  $\tilde{F} \otimes_C \tilde{E}$ . In §5.2 we compute explicit formulas for  $\tilde{\rho}_\lambda$ . In §5.3 we show directly that  $\tilde{\rho}_\lambda$  is an isomorphism for each  $\lambda \in \mathbb{Z}$ .

**1.4. Acknowledgements.** I thank Raphaël Rouquier for advice and encouragement during this project.

This work was supported by the NSF through grant DMS-1702305.

## 2. BACKGROUND: FULL ACTION

We refer the reader to [McM, §2] for many details and conventions about background structures and notation. We give here only the additional theory needed to describe the action of the negative half of  $\mathcal{U}$  and its compatibility with the positive half through the ‘commutator’ isomorphisms  $\rho_\lambda$ .

Assume we are given an algebra  $A$  and the data  $(E, x, \tau)$ , such that  ${}_A E$  is f.g. projective, and  $E^n$  is free as a  $P_n$ -module, and such that  $(E, x, \tau)$  is the data of a 2-representation of  $\mathfrak{sl}_2^+$  for the algebra  $A$ . Assume further that the 2-representation of  $\mathfrak{sl}_2^+$  afforded by  $A$  and  $(E, x, \tau)$  has a weight decomposition. For  $\lambda \in \mathbb{Z}$ , we use the notations  $A_\lambda$  and  ${}_\lambda E_\mu = A_\lambda E A_\mu$ , so that

$$A = \prod A_\lambda$$

and

$$E = \bigoplus_{\lambda, \mu} {}_\lambda E_\mu$$

give the weight decompositions.

Now we add the hypothesis that this data gives a 2-representation of the full algebra  $\mathfrak{sl}_2$ . We use the definition as in [Roua, p. 46] with  $\mathcal{V}_\lambda := A_\lambda\text{-mod}$ . (Except, note, our Hecke algebra uses the relations of that paper after substituting  $-\tau$  for  $\tau$ .) This hypothesis can be spelled out as follows.

Let  $F = \text{Hom}_A({}_A E, A)$ . Sometimes the notation  ${}^\vee E$  is used instead of  $F$ . The right adjoint  $\text{Hom}_A({}_A E, -)$  to the functor  ${}_A E \otimes_A -$  is canonically isomorphic to the functor  ${}_A F \otimes_A -$  because  ${}_A E$  is f.g. projective. Write  $\varepsilon : EF \rightarrow A$  and  $\eta : A \rightarrow FE$  for the unit and counit of the adjunction  $(E \otimes_A -, F \otimes_A -)$ .

We define  $\sigma : EF \rightarrow FE$  by:

$$\sigma = (FE\varepsilon) \circ (F\tau F) \circ (\eta EF) : EF \rightarrow FE.$$

For  $\lambda \in \mathbb{Z}^{\geq 0}$  we define:

$$(2.1) \quad \rho_\lambda = \sigma \oplus \bigoplus_{i=0}^{\lambda-1} \varepsilon \circ x^i F : EF_\lambda \rightarrow FE_\lambda \oplus A_\lambda^{\oplus \lambda},$$

and for  $\lambda \in \mathbb{Z}^{\leq 0}$ :

$$(2.2) \quad \rho_\lambda = \left( \sigma, \sum_{i=0}^{-\lambda-1} Fx^i \circ \eta \right) : EF_\lambda \oplus A_\lambda^{\oplus -\lambda} \rightarrow FE_\lambda.$$

We assume that  $\rho_\lambda$  is an  $(A, A)$ -bimodule isomorphism for each  $\lambda$ .

If a 2-representation is integrable, i.e. if  $E$  and  $F$  are locally nilpotent, then [Roua, Thm. 5.16] we know that there is another unit and counit giving a second adjunction  $(F, E)$ . When  $E$  and  $F$  are given by bimodules, the existence of the second adjunction implies that  $E_A$  and  ${}_A F$  also are f.g. projective. (Observe that for any left adjoint  $F'$ , the functor  $\text{Hom}_A({}_A F', -) \cong E \otimes_A -$  is exact, so  ${}_A F'$  is projective, and this functor also commutes with direct sums, so  ${}_A F'$  must be finitely generated. But then  $E_A = \text{Hom}({}_A F', A)$  is also f.g. projective.)

### 3. MORE BIMODULES

**Definition 3.1.** Let  $L_n$  denote  $\text{Hom}_{D^b(B)}(E'^n X_1, X_2)$ .

Note that  $L_1 = G_1$ . We will only need  $L_1$  and  $L_2$  in what follows. Observe that  $L_2$  has a right  $G_1^{\text{op}}$ -module structure. We now study  $L_2$  and provide it with the structure of  $(G_1^{\text{op}}, G_1^{\text{op}})$ -bimodule.

**Definition 3.2.** Let  $R \in \text{per } B$  be given by

$$\left( \begin{array}{c} E^2[y] \xrightarrow{(\pi E, \pi E \circ \tau)} E_y E \oplus E_y E \\ 0 \rightarrow E[y] \oplus E[y] \end{array} \right),$$

where the  $E_y$  action is by the canonical map  $E_y \otimes (E[y] \oplus E[y]) \rightarrow E_y E \oplus E_y E$ .

The matrix algebra structure of the nil-affine Hecke algebra shows that the following is an isomorphism of left  $A[y]$ -modules:

$$E^2[y] \xrightarrow[\tau y_1; \tau]{\sim} \tau y_1 E^2[y] \oplus \tau y_1 E^2[y].$$

**Lemma 3.3.** *There is an isomorphism  $R \xrightarrow{\sim} X'_2 \oplus X'_2$  in  $B$ -cplx given by the above isomorphism on the degree 0 term of the upper row, and the identity on all other terms. So  $R$  is strictly perfect.*

**Lemma 3.4.** *There is a quasi-isomorphism  $R \xrightarrow{q.i.} E'X_2$  determined by  $Id_{E^2[y]}$  on the degree 0 term of the upper row and  $\begin{pmatrix} 1 & 0 \\ 1 & -y_1 \end{pmatrix}$  on the degree 1 term of the lower row.*

*Remark 3.5.* This is not the same as the morphism in the proof of Prop. 3.8 in [McM, p. 13]. The existence only of a quasi-isomorphism was needed in Prop. 3.8, so I suggest changing that morphism to this one, since the right  $G_1^{\text{op}}$ -action on this one is easier to represent.

**Lemma 3.6.** *The complex  $R$  has a right action of the algebra  $G_1^{\text{op}}$ , where  $(\theta, \varphi) \in G_1^{\text{op}}$  acts by  $E\varphi \in \text{End}(E^2[y])$  on the degree 0 term  $E^2[y]$  in the top row, and by*

$$\begin{pmatrix} \varphi & 0 \\ \varphi_1 & \theta \end{pmatrix}$$

on the degree 1 term  $E[y]^{\oplus 2}$  in the bottom row, and by

$$\begin{pmatrix} E_y\varphi & 0 \\ E_y\varphi_1 & E_y\theta \end{pmatrix}$$

on the degree 1 term  $E_yE^{\oplus 2}$  in the top row. Through the quasi-isomorphism of the previous lemma, this action induces the canonical action of  $G_1^{\text{op}}$  on  $E'X_2$ .

*Proof.* First we check that the right action of  $(\theta, \varphi)$  gives a morphism of complexes of left  $B$ -modules. The action is clearly  $A^{\text{op}}[y]$ -linear in the top and bottom rows, and it is clearly linear over the off-diagonal generators in  $E_y$ . The action commutes with the differential on the bottom row. We check the top row:

$$\begin{aligned} \begin{pmatrix} E_y\varphi & 0 \\ E_y\varphi_1 & E_y\theta \end{pmatrix} \cdot \begin{pmatrix} \pi E \\ \pi E \circ \tau \end{pmatrix} &= \begin{pmatrix} E_y\varphi \circ \pi E \\ E_y\varphi_1 \circ \pi E + E_y\theta \circ \pi E \circ \tau \end{pmatrix} \\ &= \begin{pmatrix} \pi E \circ E\varphi \\ \pi E \circ E\varphi_1 + \pi E \circ \tau \circ E\theta \end{pmatrix} \\ &= \begin{pmatrix} \pi E \\ \pi E \circ \tau \end{pmatrix} \circ E\varphi. \end{aligned}$$

Next we check that the action commutes with multiplication in the algebra. In the algebra we have  $(\theta, \varphi) \cdot (\theta', \varphi') = (\theta\theta', \varphi' \circ \varphi)$ , while the action of the product in the degree 1 term of the bottom row is given by:

$$\begin{pmatrix} \varphi' & 0 \\ \varphi'_1 & \theta' \end{pmatrix} \cdot \begin{pmatrix} \varphi & 0 \\ \varphi_1 & \theta \end{pmatrix} = \begin{pmatrix} \varphi' \circ \varphi & 0 \\ \varphi'_1 \circ \varphi + (-\theta') \circ \varphi_1 & \theta\theta' \end{pmatrix}.$$

Note that

$$\varphi'_1 \circ \varphi + (-\theta') \circ \varphi_1 = \theta \cdot \varphi'_1 + \varphi'_1 \circ y_1\varphi_1 + \varphi_1 \cdot \theta'$$

and

$$\begin{aligned}\varphi' \circ \varphi - \_.\theta\theta' &= y_1((\_.\theta') \circ \varphi_1 + \varphi'_1 \circ (\_.\theta) + \varphi'_1 \circ y_1\varphi_1) \\ &= y_1(\varphi'_1 \circ \varphi + (\_.\theta') \circ \varphi_1),\end{aligned}$$

so the composition of the actions is indeed the action of the product for that term. The other terms are trivial to check.

Lastly we check that this action is compatible with the canonical action on  $E'X_2$  through the quasi-isomorphism of Lemma 3.4. Start with the degree 1 term in the bottom row:

$$\begin{aligned}\begin{pmatrix} 1 & 0 \\ 1 & -y_1 \end{pmatrix} \cdot \begin{pmatrix} \varphi & 0 \\ \varphi_1 & \theta \end{pmatrix} &= \begin{pmatrix} \varphi & 0 \\ \varphi - y_1\varphi_1 & -y_1\theta \end{pmatrix}, \\ \begin{pmatrix} \varphi & 0 \\ 0 & \theta \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 \\ 1 & -y_1 \end{pmatrix} &= \begin{pmatrix} \varphi & 0 \\ \theta & -y_1\theta \end{pmatrix}.\end{aligned}$$

Note that  $\varphi - y_1\varphi_1 = \theta$ , so these agree. The other terms are trivial to check.  $\square$

Now we may compute a model for  $L_2$  using the strictly perfect  $R$  as a replacement for  $E'X_2$ .

**Definition 3.7.** Define the following  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -sub-bimodule of  $F[y]^{\oplus 2} \oplus \text{Hom}_A({}_A E^2, E)[y]$ :

$$\begin{aligned}\bar{L}_2 &= \left\langle (f_1, f_2, \rho) \in F[y]^{\oplus 2} \oplus \text{Hom}_A(E^2, E)[y] \right. \\ &\quad \left. \rho = Ef_1 \circ \tau + Ef_2 + y_1 \circ \rho' \right. \\ &\quad \left. \text{for some } \rho' \in \text{Hom}_A(E^2, E)[y] \right\rangle.\end{aligned}$$

**Proposition 3.8.** *There is an isomorphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $\bar{L}_2 \xrightarrow{\sim} \text{Hom}_{K^b(B)}(R, X_2)$  determined by:*

$$(f_1, f_2, \rho) \mapsto \left( \left( \begin{pmatrix} ee, \begin{pmatrix} 0 \\ 0 \end{pmatrix} \\ 0, \begin{pmatrix} e \\ e' \end{pmatrix} \end{pmatrix} \right) \mapsto \begin{pmatrix} (\rho(ee), 0) \\ (0, f_2(e) + f_1(e')) \end{pmatrix} \right).$$

*Proof.* The proof is a direct computation of  $\text{Hom}_{K^b(B)}(R, X_2)$ . It is easy to check that the morphism given as the image of  $(f_1, f_2, \rho)$  is a morphism of complexes of left  $B$ -modules. The condition  $\rho = Ef_1 + Ef_2 \circ \tau + y_1 \circ \rho'$  is equivalent to the statement that this morphism has zero differential.  $\square$

**Corollary 3.9.** *There is an isomorphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $\bar{L}_2 \xrightarrow{\sim} L_2$  given by composing the isomorphism above with the isomorphism*

$$\text{Hom}_{K^b(B)}(R, X_2) \xrightarrow{\sim} \text{Hom}_{D^b(B)}(R, X_2).$$

**Proposition 3.10.** *There is an isomorphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $F^2[y] \xrightarrow{\sim} \text{Hom}_{K^b(B)}(R, X_1)$  given by*

$$F^2[y] \ni ff \mapsto \left( \left( \begin{array}{c} (ee, \begin{pmatrix} ee \\ ee' \end{pmatrix}) \\ (0, \begin{pmatrix} e \\ e' \end{pmatrix}) \end{array} \right) \mapsto \begin{pmatrix} ff(ee) \\ 0 \end{pmatrix} \right).$$

*Proof.* The proof is a direct computation of  $\text{Hom}_{K^b(B)}(R, X_1)$ .  $\square$

It is useful to give a description of  $G_2$  that is compatible with this description of  $L_2$ , i.e. that uses the replacement  $R$  for  $E'X_2$ .

**Definition 3.11.** Define the following  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -sub-bimodule of  $E[y]^{\oplus 2} \oplus \text{Hom}_A(E, E^2)[y]$ :

$$\begin{aligned} \bar{G}'_2 = \left\langle (e', e, \xi) \in E[y]^{\oplus 2} \oplus \text{Hom}_A(E, E^2)[y] \right. \\ \left. \begin{aligned} \xi = - \otimes e + y_2 \tau(- \otimes (e - y_1 e')) + y_1 y_2 \xi' \\ \text{for some } \xi' \in \text{Hom}_A(E, E^2)[y] \end{aligned} \right\rangle. \end{aligned}$$

**Proposition 3.12.** *There is an isomorphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $\bar{G}'_2 \xrightarrow{\sim} \text{Hom}_{K^b(B)}(X_2, R)$  determined by:*

$$(e', e, \xi) \mapsto \left( \left( \begin{array}{c} (e, 0) \\ (0, 1) \end{array} \right) \mapsto \begin{pmatrix} (\xi(e), 0) \\ e \\ e - y_1 e' \end{pmatrix} \right).$$

*Proof.* The proof is a direct computation of  $\text{Hom}_{K^b(B)}(X_2, R)$ .  $\square$

The quasi-isomorphism  $R \xrightarrow{q.i.} E'X_2$  determines an isomorphism  $\bar{G}'_2 \xrightarrow{\sim} \bar{G}_2$ , which is by  $(e', e, \xi) \mapsto (e, e - y_1 e', \xi)$ , with inverse given by  $(e_1, e_2, \xi) \mapsto (y_1^{-1}(e_1 - e_2), e_1, \xi)$ . In the remainder of this paper we will use  $\bar{G}'_2$  instead of  $\bar{G}_2$  as a model for  $G_2$ .

**Definition 3.13.** Let  $U$  denote  $\text{Hom}_{K^b(B)}(R, R)$ . It is canonically isomorphic to  $\text{Hom}_{D^b(B)}(E'X_2, E'X_2)$ .

As usual, we will define a model for  $U$  and work with the model in what follows.

**Definition 3.14.** Define the following  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -sub-bimodule of  $FE[y]^{\oplus 4} \oplus \text{Hom}_A(E^2, E^2)[y]$ :

$$\begin{aligned} \bar{U} = \left\langle (\Phi_{11}, \Phi_{21}, \Phi_{12}, \Phi_{22}, \Lambda) \in FE[y]^{\oplus 4} \oplus \text{Hom}_A(AE^2, E^2)[y] \right. \\ \left. \begin{aligned} \Lambda = \tau y_1 (E\Phi_{11} + E\Phi_{12} \circ \tau) - y_2 \tau y_1 (E\Phi_{21} + E\Phi_{22} \circ \tau) + y_1 y_2 \Lambda^\circ \\ \text{for some } \Lambda^\circ \in \text{Hom}_A(AE^2, E^2)[y] \end{aligned} \right\rangle. \end{aligned}$$

**Proposition 3.15.** *There is an isomorphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $\bar{U} \rightarrow U$  determined by*

$$(\Phi_{11}, \Phi_{21}, \Phi_{12}, \Phi_{22}, \Lambda) \mapsto \left( \left( \begin{array}{c} (ee, \begin{pmatrix} 0 \\ 0 \end{pmatrix}) \\ (0, \begin{pmatrix} e \\ e' \end{pmatrix}) \end{array} \right) \mapsto \left( \begin{array}{c} (\Lambda(ee), \begin{pmatrix} 0 \\ 0 \end{pmatrix}) \\ (0, [\Phi] \cdot \begin{pmatrix} e \\ e' \end{pmatrix}) \end{array} \right) \right).$$

*Proof.* The proof is a direct computation of  $U = \text{Hom}_{K^b(B)}(R, R)$ . We must show that the condition on  $\Lambda$  is equivalent to the statement that the image of  $(\Phi_{11}, \Phi_{21}, \Phi_{12}, \Phi_{22}, \Lambda)$  has zero differential. One computes directly that the image has zero differential if and only if the following pair of equations is valid:

$$\begin{cases} \pi E \circ \Lambda = E_y \Phi_{11} \circ \pi E + E_y \Phi_{12} \circ \pi E \circ \tau \\ \pi E \circ \tau \circ \Lambda = E_y \Phi_{21} \circ \pi E + E_y \Phi_{22} \circ \pi E \circ \tau. \end{cases}$$

That pair is manifestly equivalent to the pair:

$$(3.1) \quad \begin{cases} \Lambda = E\Phi_{11} + E\Phi_{12} \circ \tau + y_2 \Lambda' \\ \tau \circ \Lambda = E\Phi_{21} + E\Phi_{22} \circ \tau + y_2 \Lambda'' \end{cases}$$

for some  $\Lambda', \Lambda'' \in \text{Hom}_A({}_A E^2, E^2)[y]$ .

*Claim 3.16.* Suppose  $(\Phi, \Lambda)$  is given such that (3.1) holds for some  $\Lambda', \Lambda''$ . Then there is a  $\Lambda^\circ \in \text{Hom}_A({}_A E^2, E^2)[y]$  such that

$$(3.2) \quad \Lambda = \tau y_1 (E\Phi_{11} + E\Phi_{12} \circ \tau) - y_2 \tau y_1 (E\Phi_{21} + E\Phi_{22} \circ \tau) + y_1 y_2 \Lambda^\circ.$$

*Proof.* Multiply the second equation of the pair by  $\tau$  and obtain:

$$-\tau y_2 \Lambda'' = \tau \circ E\Phi_{21} + \tau \circ E\Phi_{22} \circ \tau.$$

Multiply the first by  $\tau$  and the second by  $\tau y_1$  and identify the results to obtain:

$$\tau y_2 \Lambda' = y_1 y_2 \tau \circ \Lambda'' + \tau y_1 \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) - \tau \circ (E\Phi_{11} + E\Phi_{12} \circ \tau).$$

Then:

$$\begin{aligned} \Lambda' &= (y_1 \tau - \tau y_2) \circ \Lambda' \\ &= y_1 \tau \circ \Lambda' - y_1 y_2 \tau \circ \Lambda'' - \tau y_1 \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) \\ &\quad + \tau \circ (E\Phi_{11} + E\Phi_{12} \circ \tau) \\ &= y_1 (\tau \circ \Lambda' - y_2 \tau \circ \Lambda'') - \tau y_1 \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) \\ &\quad + \tau \circ (E\Phi_{11} + E\Phi_{12} \circ \tau). \end{aligned}$$

Let  $\Lambda^\circ = \tau \circ \Lambda' - y_2 \tau \circ \Lambda''$ . Then:

$$\begin{aligned} \Lambda &= E\Phi_{11} + E\Phi_{12} \circ \tau + y_1 y_2 \Lambda^\circ \\ &\quad - y_2 \tau y_1 \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) + y_2 \tau \circ (E\Phi_{11} + E\Phi_{12} \circ \tau) \\ &= \tau y_1 \circ (E\Phi_{11} + E\Phi_{12} \circ \tau) - y_2 \tau y_1 \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) + y_1 y_2 \Lambda^\circ, \end{aligned}$$

as desired.  $\square$

*Claim 3.17.* Now suppose  $(\Phi, \Lambda)$  and  $\Lambda^\circ$  are given such that (3.2) holds. Then there are  $\Lambda', \Lambda''$  such that (3.1) holds.



*Proof.* Let

$$\begin{aligned}\Lambda' &= \tau \circ (E\Phi_{11} + E\Phi_{12} \circ \tau) - \tau y_1 \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) + y_1 \Lambda^\circ, \\ \Lambda'' &= \tau \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) + y_1 \tau \circ \Lambda^\circ.\end{aligned}$$

Multiplying the first by  $y_2$ , adding  $E\Phi_{11} + E\Phi_{12} \circ \tau$ , and simplifying with (3.2), we find:

$$y_2 \Lambda' + E\Phi_{11} + E\Phi_{12} \circ \tau = \Lambda.$$

Multiplying the second by  $y_2$  and adding  $E\Phi_{21} + E\Phi_{22} \circ \tau$ , we find:

$$\begin{aligned}y_2 \Lambda'' + E\Phi_{21} + E\Phi_{22} \circ \tau &= y_2 \tau \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) + y_2 y_1 \tau \circ \Lambda^\circ \\ &\quad + E\Phi_{21} + E\Phi_{22} \circ \tau \\ &= \tau y_1 \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) + \tau y_1 y_2 \circ \Lambda^\circ,\end{aligned}$$

while

$$\begin{aligned}\tau \circ \Lambda &= -\tau y_2 \tau y_1 \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) + \tau y_2 y_1 \Lambda^\circ \\ &= \tau y_1 \circ (E\Phi_{21} + E\Phi_{22} \circ \tau) + y_1 y_2 \tau \Lambda^\circ\end{aligned}$$

using (3.2). So the pair of equations (3.1) is satisfied.  $\square$

The proposition follows from the two claims.  $\square$

We will need to know the  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodule structure of the components of  $\tilde{E}$  and  $\tilde{E}^2$  and  $\tilde{F}$ . This information is implicit in the calculations in [McM] for the first three bimodules in the next proposition. The fourth bimodule is easy to compute using the map provided.

**Proposition 3.18.** *We have isomorphisms of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules:*

- $y_1 \dots y_n E^n[y] \xrightarrow{\sim} E^n[y]$  given by  $y_1^{-1} \dots y_n^{-1}$ .
- $L_1 = G_1 \xrightarrow{\sim} A[y] \oplus FE[y]$  given by  $(\theta, \varphi) \mapsto (\theta, \varphi_1)$ , where

$$\varphi_1 = y_1^{-1}(\varphi - \theta)$$

*is interpreted in  $FE[y]$ . Note that the summand  $FE[y]$  is a left  $G_1^{\text{op}}$ -submodule of  $G_1$ .*

- $G_2 \xrightarrow{\sim} E[y] \oplus E[y] \oplus FE^2[y]$  given by  $(e', e, \xi) \mapsto (e', e, \xi')$ , where

$$\xi' = y_1^{-1} y_2^{-1} (\xi - \_ \otimes e - \tau y_1 (\_ \otimes (e - y_1 e')))$$

*is interpreted in  $FE^2[y]$ . Note that the summand  $FE^2[y]$  is a left  $G_1^{\text{op}}$ -submodule of  $G_2$ .*

- $L_2 \xrightarrow{\sim} F[y] \oplus F[y] \oplus F^2E[y]$  given by  $(f_1, f_2, \rho) \mapsto (f_1, f_2, \rho_1)$ , where

$$\rho_1 = y_1^{-1}(\rho - Ef_1 - Ef_2 \circ \tau)$$

*is interpreted in  $F^2E[y]$ . Note that the summand  $F^2E[y]$  is a left  $G_1^{\text{op}}$ -submodule of  $L_2$ .*

- $U \rightarrow FE[y]^{\oplus 4} \oplus F^2E^2[y]$  given by

$$(\Phi_{11}, \Phi_{21}, \Phi_{12}, \Phi_{22}, \Lambda) \mapsto (\Phi_{11}, \Phi_{21}, \Phi_{12}, \Phi_{22}, \Lambda^\circ),$$

where

$$(3.3) \quad \Lambda = \tau y_1 (E\Phi_{11} + E\Phi_{12} \circ \tau) - y_2 \tau y_1 (E\Phi_{21} + E\Phi_{22} \circ \tau) + y_1 y_2 \Lambda^\circ$$

determines  $\Lambda^\circ$ , which is interpreted in  $F^2 E^2[y]$ . Note that the summand  $F^2 E^2[y]$  is a left  $G_1^{\text{op}}$ -submodule of  $U$ .

In what follows we will frequently use the bimodule descriptions on the right side of these isomorphisms for the components of  $\tilde{E}$  and  $\tilde{F}$ . Sometimes, to avoid confusion, we will use the shorthand expressions ‘submodule form’ and ‘bimodule form’ to distinguish the two sides. The data of an element given in the two forms will differ only in the last component, where in the ‘submodule form’ the last component gives the morphism induced on the degree 0 part of the top row of the  $B$ -module complexes, and in the ‘bimodule form’ the last component gives just the remainder term ‘ $\varphi_1$ ’, ‘ $\xi'$ ’, ‘ $\rho_1$ ’, ‘ $\chi''$ ’, or ‘ $\Lambda^\circ$ ’. Note that here ‘submodule’ indicates ‘ $(A^{\text{op}}[y], A^{\text{op}}[y])$ -sub-bimodule’, and ‘bimodule’ indicates ‘structure as  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodule’.

#### 4. ADJUNCTION $(\tilde{E}, \tilde{F})$

**Definition 4.1.** Let  $\tilde{F}$  denote the  $(C, C)$ -bimodule  $\text{Hom}_C({}_C \tilde{E}, C)$ .

We have seen that, under the hypotheses in [McM],  ${}_C \tilde{E}$  is f.g. projective. It follows that the right adjoint functor  $\text{Hom}_C({}_C \tilde{E}, -)$  of  $\tilde{E} \otimes_C -$  is canonically isomorphic to  $\tilde{F} \otimes_C -$ . We have already defined  $\tilde{x}$  and  $\tilde{\tau}$ . So we may define  $\tilde{\varepsilon} : \tilde{E} \tilde{F} \rightarrow C$  and  $\tilde{\eta} : C \rightarrow \tilde{F} \tilde{E}$  and  $\tilde{\sigma}$  and  $\tilde{\rho}_\lambda$  using the formulas in §2, but with  $A$  replaced by  $C$  and  $E, F, x, \tau$  replaced by  $\tilde{E}, \tilde{F}, \tilde{x}, \tilde{\tau}$ . Note that by this definition,  $\tilde{\varepsilon}$  is the usual evaluation functor, and  $\tilde{\eta}$  sends  $1 \in C$  to  $\text{Id}_{\tilde{E}} \in \tilde{F} \tilde{E}$ . (Where  $\tilde{F} \tilde{E}$  is viewed through the canonical isomorphism  $\text{Hom}(\tilde{E}, C) \otimes_C \tilde{E} \xrightarrow{\sim} \text{Hom}(\tilde{E}, \tilde{E})$ .)

Now we construct an isomorphism of  $(C, C)$ -bimodules

$$\tilde{F} \xrightarrow{\sim} \text{Hom}_{K^b(B)}(X_2 \oplus R, X)$$

as follows:

$$\begin{aligned} \tilde{F} &= \text{Hom}_C({}_C \tilde{E}, C) \\ &\xrightarrow{\sim} \text{Hom}_{D^b(C)}(\tilde{E}, C) \\ &\xrightarrow{\sim} \text{Hom}_{D^b(C)}(\mathcal{E}, \mathcal{C}) \\ &\xrightarrow{\sim} \text{Hom}_{D^b(C)}(\mathcal{H}om_B(X, E'X), \mathcal{H}om_B(X, X)) \\ &\xrightarrow{\sim} \text{Hom}_{D^b(B)}(E'X, X) \\ &\xrightarrow{\sim} \text{Hom}_{D^b(B)}(E'X_1 \oplus R, X) \\ &\xrightarrow{\sim} \text{Hom}_{K^b(B)}(E'X_1 \oplus R, X_1 \oplus X_2). \end{aligned}$$

(The fourth arrow comes from the Rickard equivalence.) The right action of  $C$  on the bimodule in the last line is given by post-composition. The left action is naturally defined using pre-composition, functoriality of  $E'$ , the isomorphism

$$\text{Hom}_{K^b(B)}(E'X_1, R) \xrightarrow{\sim} \text{Hom}_{K^b(B)}(E'X_1, E'X_2)$$

(since  $E'X_1$  is strictly perfect), and the right action of  $G_1^{\text{op}}$  on  $R$  (which commutes with the action on  $E'X_2$ ).

With this description of  $\tilde{F}$ , and using the direct sum decompositions, we can compute  $\tilde{F}$  as a  $2 \times 2$  matrix of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules, and we find:

$$\tilde{F} \xrightarrow{\sim} \begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix}.$$

## 5. MAPS $\tilde{\rho}_\lambda$ ARE ISOMORPHISMS

**5.1. Some tensor products in  $C\text{-mod}$ .** In this section we compute three tensor products of bimodules over  $C$ , namely  $\tilde{E}\tilde{E}$ ,  $\tilde{F}\tilde{E}$ , and  $\tilde{E}\tilde{F}$ , and describe the products in each case as matrices of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules. (That is to say, as bimodules over the subalgebra  $\begin{pmatrix} A^{\text{op}}[y] & 0 \\ 0 & A^{\text{op}}[y] \end{pmatrix} \subset C$ .) These calculations are used in the next section to verify that  $\tilde{\rho}_\lambda$  are isomorphisms. Note that the product  $\tilde{E}\tilde{E} = \tilde{E}^2$  is already given [McM, Prop. 3.41] a description as a matrix of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules using the identification with  $\text{Hom}_{K^b(B)}(X, E'^2 X)$ , but for a computation of  $\tilde{\sigma}$  it is also necessary to realize this matrix description as arising from the matrix description of  $\tilde{E}$  as a  $(C, C)$ -bimodule.

The tensor products are computed according to the general formulation described in [McM, §2.4]. First we take the tensor product over the subalgebra  $\Delta := \begin{pmatrix} A^{\text{op}}[y] & 0 \\ 0 & G_1^{\text{op}} \end{pmatrix} \subset C$ . This product is given on coefficients by matrix multiplication and tensor product over  $A^{\text{op}}[y]$  or  $G_1^{\text{op}}$ . After this we must take the quotient by the image of the map “ $I_B + I_C$ ” (cf. §2.4) that is produced using the action of the off-diagonal coefficients of  $C$ . This quotient is taken separately on each coefficient of the product over  $\Delta$ .

The basic technique for computing a quotient by the image of (say)  $I_B$  is to identify one of its projections as an isomorphism. (In §2.4, there is a projection of  $I_B$  to  $M_1 \otimes_A N_1$  and another projection to  $M_2 \otimes_D N_2$ .) In this situation the quotient by the image of  $I_B$  reduces to the summand of the other projection, because every element of the first (in the quotient) has a unique representative in the second. The basic technique for computing the quotient by the image of a sum  $I_B + I_C$  is to show that the projections of the second map have a compatibility with those of the first. Many of the coefficients encountered below are computed in this way, but a few of them require more complicated reasoning.

Let us write, in general,  $I'_\beta$  for the projection of  $I_B$  to the first summand, and  $-I''_\beta$  for the projection to the second. Similarly write  $I'_\delta$  and  $-I''_\delta$  for the projections of  $I_D$ . Here ‘first’ and ‘second’ and ‘ $I_B$ ’ and ‘ $I_D$ ’ are understood as in §2.4. In a tensor product of  $(C, C)$ -bimodules, each of the four coefficients will have its own set of maps  $I'_\beta, I''_\beta, I'_\delta, I''_\delta$ .

**5.1.1.  $\tilde{E}\tilde{E}$ .** For the product  $\tilde{E}\tilde{E}$ , we already know the structure of the coefficients of the matrix presentation. We will need to compute the action of  $\tilde{\tau}$  on certain elements of  $\tilde{E}\tilde{E}$  in order to compute  $\tilde{\sigma}$ , and for this it will be enough to compute the map from the tensor product over  $\Delta$  to the product over  $C$ . Write  $\Gamma$  for this map. Let the subscript ‘ $G$ ’ between concatenated

modules indicate the tensor product over  $G_1^{\text{op}}$ . (A lacking subscript indicates the product over  $A^{\text{op}}[y]$ .) So we have:

$$\begin{aligned}
 \tilde{E} \otimes_{\Delta} \tilde{E} &\cong \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \otimes_{\Delta} \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \\
 (5.1) \quad &\cong \begin{pmatrix} EE[y] \oplus E^2[y]_G G_1 & EE^2[y] \oplus E^2[y]_G G_2 \\ G_1 E[y] \oplus (G_2)_G G_1 & G_1 E^2[y] \oplus (G_2)_G G_2 \end{pmatrix} \\
 &\xrightarrow{\Gamma} \begin{pmatrix} E^2[y] & E^3[y] \\ G_2 & G_3 \end{pmatrix} \cong [\tilde{E}^2],
 \end{aligned}$$

and we wish to understand the map  $\Gamma$  on each coefficient. This map is the projection to the quotient by the image of  $I_B + I_D$ . In subsequent sections we must compute this quotient to find the structure of the image of  $\Gamma$ , but here, since we know the image to be  $\text{Hom}_{K^b(B)}(X_1 \oplus X_2, E'^2(X_1 \oplus X_2))$ , we are able to compute the map in terms of composition and functoriality of  $E'$ .

- For  $\Gamma_{11}$ , we have:
  - $\Gamma_{11} |_{EE[y]}$  is given by  $\text{Id}_{EE}$ ,
  - $\Gamma_{11} |_{(E^2[y])_G G_1}$  is given as the inverse of  $EE[y] \xrightarrow{\sim} (E^2[y])_G G_1$ ,  $ee \mapsto ee \otimes 1_{G_1}$ .
- For  $\Gamma_{21}$ , we have:
  - $\Gamma_{21} |_{G_1 E[y]}$  is given by

$$(\theta, \varphi_1) \otimes e \mapsto (\theta e, \theta y_1 e, \varphi_1(-) \otimes e) \in G_2,$$

- $\Gamma_{21} |_{(G_2)_G G_1}$  is given as the inverse of  $G_2 \xrightarrow{\sim} (G_2)_G G_1$ ,  $g_2 \mapsto g_2 \otimes 1_{G_1}$ .
- For  $\Gamma_{12}$ , we have:
  - $\Gamma_{12} |_{EE^2[y]}$  is given by  $\text{Id}_{E^3[y]}$ ,
  - $\Gamma_{12} |_{(E^2[y])_G G_2}$  is given by

$$ee \otimes (e', e, \xi') \mapsto y_1^{-1} y_2^{-1} y_3^{-1} (E\xi)(y_1 y_2 ee) \in E^3[y].$$

- For  $\Gamma_{22}$ , we have:
  - $\Gamma_{22} |_{G_1 E^2[y]}$  is given by

$$(\theta, \varphi_1) \otimes ee \mapsto (\theta y_1 y_2 ee, 0, 0, \varphi_1(-) \otimes ee)$$

(c.f. Diagram  $D_{1|2}(2, 1, 1)$  in [McM, §4.2]),

- $\Gamma_{22} |_{(G_2)_G G_2}$  is given by

$$(e_1, e_2, \xi) \otimes (\bar{e}_1, \bar{e}_2, \bar{\xi}) \mapsto (\bar{\xi}(e_1), e_2 \otimes \bar{e}_1, e_2 \otimes \bar{e}_2, E\bar{\xi} \circ \xi)$$

(c.f. Diagram  $D_{1|2}(2, 2, 1)$ ). We will need to compute this map for elements expressed in the form

$$(e', e, \xi') \in \text{Hom}_{K^b(B)}(X_2, R)$$

using  $e_1 = e$ ,  $e_2 = e - y_1 e'$ , and  $\xi = - \otimes e_1 + y_2 \tau(- \otimes e_2) + y_1 y_2 \xi'$ , and analogously for  $\bar{e}'$ ,  $\bar{e}$ , and  $\bar{\xi}'$ . We first compute  $E\bar{\xi} \circ \xi$ :

$$\begin{aligned}
 E\bar{\xi} \circ \xi &= (- \otimes \bar{e} + y_2 \tau(- \otimes (\bar{e} - y_1 \bar{e}')) + y_1 y_2 E\bar{\xi}') \\
 &\circ (- \otimes e + y_2 \tau(- \otimes (e - y_1 e')) + y_1 y_2 \xi') \\
 &= - \otimes (e\bar{e} + y_2 \tau(e\bar{e} - y_1 e\bar{e}')) + y_1 y_2 \bar{\xi}'(e) \\
 &+ y_3 \circ \tau E(- \otimes (e\bar{e} - y_2 e'\bar{e})) \\
 &+ y_2 y_3 \circ E\tau \circ \tau E(- \otimes (e - y_1 e')(\bar{e} - y_1 \bar{e}')) \\
 &+ y_1 y_2 y_3 (E\bar{\xi}' \circ \tau(- \otimes (e - y_1 e')) + E\tau(\xi' \otimes (\bar{e} - y_1 \bar{e}')) + \xi' \otimes \bar{e}').
 \end{aligned}$$

Using [McM, Prop. 3.21], we can read off the data of our desired formula:

$$\begin{aligned}
 (e', e, \xi') \otimes (\bar{e}', \bar{e}, \bar{\xi}') &\mapsto \\
 (e\bar{e} + y_2 \tau(e\bar{e} - y_1 e\bar{e}')) + y_1 y_2 \bar{\xi}'(e), &e\bar{e} - y_2 e'\bar{e}, (e - y_1 e')(\bar{e} - y_1 \bar{e}'), \\
 E\bar{\xi}' \circ \tau(- \otimes (e - y_1 e')) + E\tau(\xi' \otimes (\bar{e} - y_1 \bar{e}')) &+ \xi' \otimes \bar{e}'.
 \end{aligned}$$

Here the last component of the image has the form  $\chi''$ , so the image is  $(ee_1, ee_2, ee_3, \chi'')$  rather than  $(ee_1, ee_2, ee_3, \chi)$ .

5.1.2.  $\tilde{F}\tilde{E}$ . For the product  $\tilde{F}\tilde{E}$ , we can find the matrix presentation of its coefficients the same technique as for  $\tilde{F}$  and  $\tilde{E}^2$ . We have:

$$\begin{aligned}
 \tilde{F}\tilde{E} &= \text{Hom}_C({}_C\tilde{E}, C) \otimes_C \tilde{E} \\
 &\xrightarrow{\sim} \text{Hom}_C({}_C\tilde{E}, \tilde{E}) \\
 &\xrightarrow{\sim} \text{Hom}_{D^b(C)}(\tilde{E}, \tilde{E}) \\
 &\xrightarrow{\sim} \text{Hom}_{D^b(C)}(\mathcal{E}, \mathcal{E}) \\
 &\xrightarrow{\sim} \text{Hom}_{D^b(C)}(\mathcal{H}om_B(X, E'X), \mathcal{H}om_B(X, E'X)) \\
 &\xrightarrow{\sim} \text{Hom}_{D^b(B)}(E'X, E'X) \\
 &\xrightarrow{\sim} \text{Hom}_{K^b(B)}(E'X_1 \oplus R, E'X_1 \oplus R).
 \end{aligned}$$

So the matrix presentation is:

$$[\tilde{F}\tilde{E}] = \begin{pmatrix} G_1 & G_2 \\ L_2 & U \end{pmatrix}$$

where

$$\begin{aligned}
 U &:= \text{Hom}_{K^b(B)}(R, R) \\
 &\cong \text{Hom}_{D^b(B)}(E'X_2, E'X_2).
 \end{aligned}$$

Just as for  $\tilde{E}^2$ , we describe the morphisms from the coefficients of the product over  $\Delta$  to those of the product over  $C$ , that is, the coefficients of  $\Gamma$ :

$$\begin{aligned}
 \tilde{F} \otimes_{\Delta} \tilde{E} &\cong \begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix} \otimes_{\Delta} \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \\
 (5.2) \quad &\cong \begin{pmatrix} FE[y] \oplus (L_1)_G G_1 & FE^2[y] \oplus (L_1)_G G_2 \\ F^2E[y] \oplus (L_2)_G G_1 & F^2E^2[y] \oplus (L_2)_G G_2 \end{pmatrix} \\
 &\xrightarrow{\Gamma} \begin{pmatrix} G_1 & G_2 \\ L_2 & U \end{pmatrix}.
 \end{aligned}$$

- We have  $\Gamma_{11} : FE[y] \oplus G_1 \rightarrow G_1$  given by  $(\iota, \text{Id}_{G_1})$ :
  - $I'_{\beta} : FE[y]_G G_1 \xrightarrow{\sim} FE[y]$  given as the inverse of  $(fe \mapsto fe \otimes 1_{G_1})$ ,
  - $I''_{\beta} : FE[y]_G G_1 \xrightarrow{\iota \otimes G_1} (G_1)_G G_1 \cong G_1$ ,
  - $I'_{\delta} : (G_1)_G FE[y] \xrightarrow{\sim} FE[y]$  given as the inverse of  $(fe \mapsto 1_{G_1} \otimes fe)$ ,
  - $I''_{\delta} : (G_1)_G FE[y] \xrightarrow{G_1 \otimes \iota} (G_1)_G G_1 \cong G_1$ .

Here the map  $\iota : FE[y] \hookrightarrow G_1$  is the inclusion as the second summand indicated in Prop. 3.18. Using either  $I''_{\beta} \circ I'^{-1}_{\beta}$ , or  $I''_{\delta} \circ I'^{-1}_{\delta}$ , one can associate a unique representative in  $(G_1)_G G_1$  to each element of  $FE[y]$ . We also have  $(G_1)_G G_1 \xrightarrow{\sim} G_1$  as the inverse of  $(g \mapsto 1_{G_1} \otimes g)$ , from which we see that  $I''_{\beta} \circ I'^{-1}_{\beta} = I''_{\delta} \circ I'^{-1}_{\delta}$ , and the two associate the same representatives. The result follows.

- We have  $\Gamma_{21} : F^2E[y] \oplus L_2 \rightarrow L_2$  given by  $(\iota', \text{Id}_{L_2})$ :
  - $I'_{\beta} : F^2E[y]_G G_1 \xrightarrow{\sim} F^2E[y]$  given as the inverse of  $(ffe \mapsto ffe \otimes 1_{G_1})$ ,
  - $I''_{\beta} : F^2E[y]_G G_1 \xrightarrow{\iota' \otimes G_1} (L_2)_G G_1 \cong L_2$ ,
  - $I'_{\delta} : (L_2)_G FE[y] \rightarrow F^2E[y]$  given by  $(f', f, \rho') \otimes \bar{f}\bar{e} \mapsto (\bar{f} \circ \rho) \otimes \bar{e} = f \circ (Ef + Ef' \circ \tau + y_1 \rho') \otimes \bar{e}$ ,
  - $I''_{\delta} : (L_2)_G FE[y] \xrightarrow{L_2 \otimes \iota} (L_2)_G G_1 \cong L_2$ .

Here the map  $\iota' : F^2E[y] \hookrightarrow L_2$  is the inclusion as the second summand indicated in Prop. 3.18. Consider the first two maps. We have that  $I''_{\beta} \circ I'^{-1}_{\beta} = \iota'$  as maps  $F^2E[y] \rightarrow L_2$ . Consider the last two maps. One may check that  $\iota' \circ I'_{\delta} = I''_{\delta}$ . It follows that  $\text{Im}(I'_{\delta} - I''_{\delta}) \subset \text{Im}(I'_{\beta} - I''_{\beta})$ , so every element of  $F^2E[y]$  is associated to a unique element of  $L_2$  by applying the map  $\iota'$ .

- We have  $\Gamma_{12} : FE^2[y] \oplus G_2 \rightarrow G_2$  given by  $(\iota'', \text{Id}_{G_2})$ :
  - $I'_{\beta} : FE[y]_G G_2 \rightarrow FE^2[y]$  given by  $\bar{f}\bar{e} \otimes (e', e, \xi') \mapsto \bar{f} \otimes y_1^{-1} y_2^{-1} \xi(y_1 \bar{e}) = \bar{f} \otimes (\tau(\bar{e} \otimes e) - y_2 \tau(\bar{e} \otimes e') + \xi'(y_1 \bar{e}))$ ,
  - $I''_{\beta} : FE[y]_G G_2 \xrightarrow{\iota \otimes G_2} (G_1)_G G_2 \cong G_2$ ,
  - $I'_{\delta} : (G_1)_G FE^2[y] \xrightarrow{\sim} FE^2[y]$  given as the inverse of  $(fee \mapsto fee \otimes 1_{G_1})$ ,
  - $I''_{\delta} : (G_1)_G FE^2[y] \xrightarrow{G_1 \otimes \iota''} (G_1)_G G_2 \cong G_2$ .

Here the map  $\iota'' : FE^2[y] \hookrightarrow G_2$  is the inclusion as the second summand indicated in Prop. 3.18. Consider the last two maps. We have that  $I''_{\delta} \circ I'^{-1}_{\delta} = \iota''$  as maps  $FE^2[y] \rightarrow G_2$ . Now consider the first two maps. Observe that

$I''_\beta = \iota'' \circ I'_\beta$ . It follows that  $\text{Im}(I'_\beta - I''_\beta) \subset \text{Im}(I'_\delta - I''_\delta)$ , so every element of  $F^2E^2[y]$  is associated to a unique element of  $G_2$  by applying the map  $\iota''$ .

- We have  $\Gamma_{22} : F^2E^2[y] \oplus (L_2)_G G_2 \rightarrow U$  given by  $(\iota''', \text{Id}_U)$ :
  - $I'_\beta : F^2E[y]_G G_2 \rightarrow F^2E^2[y]$  given by  $\overline{f\bar{f}\bar{e}} \otimes (e', e, \xi') \mapsto \overline{f\bar{f}} \otimes y_1^{-1} y_2^{-1} \xi(y_1 \bar{e})$ ,
  - $I''_\beta : F^2E[y]_G G_2 \xrightarrow{\iota'' \otimes G_2} (L_2)_G G_2 \cong U$ ,
  - $I'_\delta : (L_2)_G F^2E^2[y] \rightarrow F^2E^2[y]$  given by  $(f', f, \rho') \otimes \bar{f}\bar{e}\bar{e} \mapsto (\bar{f} \circ \rho) \otimes \bar{e}\bar{e}$ ,
  - $I''_\delta : (L_2)_G F^2E^2[y] \xrightarrow{L_2 \otimes \iota''} (L_2)_G G_2 \cong U$ .

Define  $\iota''' : F^2E^2[y] \rightarrow U$  to be the inclusion as the second summand indicated in Prop. 3.18. Consider the first two maps. Observe that

$$I'_\beta(\overline{f\bar{f}\bar{e}} \otimes (e, y_1 e, \xi' = 0)) = \overline{f\bar{f}} \otimes (\bar{e} \otimes e) \in F^2E^2[y].$$

It follows that  $I'_\beta$  is surjective. Now we show that  $\iota''' \circ I'_\beta = I''_\beta$  and that  $\iota''' \circ I'_\delta = I''_\delta$ :

$$\begin{aligned} \iota'''(\overline{f\bar{f}} \otimes y_1^{-1} y_2^{-1} \xi(y_1 \bar{e})) &= (0, 0, 0, 0, \Lambda^\circ = \overline{f\bar{f}} \otimes y_1^{-1} y_2^{-1} \xi(y_1 \bar{e})), \\ I''_\beta(\overline{f\bar{f}\bar{e}} \otimes (e', e, \xi')) &= (0, 0, \overline{f\bar{f}\bar{e}}) \otimes_G (e', e, \xi') \\ &\mapsto (0, 0, 0, 0, \overline{f\bar{f}} \otimes y_1^{-1} y_2^{-1} \xi(y_1 \bar{e})); \end{aligned}$$

and

$$\begin{aligned} \iota'''((\bar{f} \circ \rho) \otimes \bar{e}\bar{e}) &= (0, 0, 0, 0, (\bar{f} \circ \rho) \otimes \bar{e}\bar{e}), \\ I''_\delta((f', f, \rho') \otimes \bar{f}\bar{e}\bar{e}) &= (f', f, \rho') \otimes (0, 0, \bar{f}\bar{e}\bar{e}) \\ &\mapsto (0, 0, 0, 0, (\bar{f} \circ \rho) \otimes \bar{e}\bar{e}). \end{aligned}$$

It follows that every element of  $F^2E^2[y]$  in the quotient is associated to a unique representative in  $U$  by applying  $\iota'''$ .

*Remark 5.1.* The map  $\iota'''$  describes the inclusion of the morphisms of  $\text{Hom}_{K^b(B)}(R, R)$  that factor through  $X_1$ . Furthermore, the maps  $I'_\beta$  and  $I'_\delta$  are in fact isomorphisms because  $R$  is a sum of summands of  $X_2$ .

5.1.3.  $\tilde{E}\tilde{F}$ . For the product  $\tilde{E}\tilde{F}$ , we do not already have a matrix presentation of its coefficients arising from the Rickard equivalence. Nonetheless, we can compute these coefficients using mostly the same techniques as those used for the other products above.

$$\begin{aligned} \tilde{E} \otimes_\Delta \tilde{F} &\cong \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \otimes_\Delta \begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix} \\ (5.3) \quad &\cong \begin{pmatrix} EF[y] \oplus E^2[y]_G F^2[y] & E[y]G_1 \oplus E^2[y]_G L_2 \\ G_1 F[y] \oplus (G_2)_G F^2[y] & G_1 G_1 \oplus (G_2)_G L_2 \end{pmatrix} \\ &\xrightarrow{\Gamma} \begin{pmatrix} EF[y] & E[y]G_1 \\ G_1 F[y] & G_1 G_1 \oplus EF[y] \end{pmatrix}. \end{aligned}$$

We wish to define  $\Gamma$ , show that it is surjective, and show that its kernel is the image of  $I_B + I_D$ . As before, we can proceed by studying the components separately.

- We have  $\Gamma_{11} : EF[y] \oplus E^2[y]_G F^2[y] \rightarrow EF[y]$  given by  $(\text{Id}_{EF[y]}, \omega)$ :

–  $I'_\beta : EE[y]_GF^2[y] \rightarrow EF[y]$  given by

$$\begin{aligned} e_1 \otimes e_2 \otimes f_2 f_1 &\mapsto e_1 \otimes (f_1 \circ E f_2 \circ (e_2, y_1 e_2, 0)) \\ &= e_1 \otimes f_1 \circ E f_2 \circ (- \otimes y_1 e_2) \\ &= e_1 \otimes f_1(-.f_2(y_1 e_2)) \\ &= e_1 \otimes f_2(y_1 e_2).f_1, \end{aligned}$$

–  $I''_\beta : EE[y]_GF^2[y] \xrightarrow{\text{Id}} E^2[y]_GF^2[y]$ ,

–  $I'_\delta : E^2[y]_GFF[y] \rightarrow EF[y]$  given by

$$\begin{aligned} e_1 e_2 \otimes f_2 \otimes f_1 &\mapsto y_1^{-1}((0, f_2, 0) \circ y_1 y_2 (e_1 e_2)) \otimes f_1 \\ &= E(f_2 \circ y_1)(e_1 e_2) \otimes f_1 \\ &= e_1.f_2(y_1 e_2) \otimes f_1 \\ &= e_1 \otimes f_2(y_1 e_2).f_1, \end{aligned}$$

–  $I''_\delta : E^2[y]_GFF[y] \xrightarrow{\text{Id}} E^2[y]_GF^2[y]$ .

Define  $\bar{\omega} : E^2 F^2[y] \rightarrow EF[y]$  by  $e_1 e_2 \otimes f_2 f_1 \mapsto e_1.f_2(e_2) \otimes f_1$ . Observe that  $e_1 \otimes y_1 \varphi_1(e_2) \otimes f_2 f_1$  and  $e_1 e_2 \otimes f_2 \circ y_1 \varphi_1 \otimes f_1$  are both sent to  $e_1.f_2(y_1 \varphi_1(e_2)) \otimes f_1$ . So  $\bar{\omega}$  is also middle-linear over generators in the second summand of  $A[y] \oplus y_1 FE[y] \cong G_1$ , and it descends to a morphism  $\omega : E^2[y]_GF^2[y] \rightarrow EF[y]$ . It is clear that  $I'_\beta = \omega$  and  $I'_\delta = \omega$  after identifying  $EE[y] \cong E^2[y]$  and  $FF[y] \cong F^2[y]$ . It follows that the kernel of  $\Gamma_{11}$  is the image of  $I'_\beta - I''_\beta$ , which is also the image of  $I'_\delta - I''_\delta$  and thus the image of the sum  $I'_\beta - I''_\beta + I'_\delta - I''_\delta = I_B + I_D$ .

*Remark 5.2.* The map  $\omega$  corresponds on the models to the map

$$\text{Hom}_{K^b(B)}(X_2, R) \otimes_{G_1^{\text{op}}} \text{Hom}_{K^b(B)}(R, X_2) \rightarrow \text{Hom}_{K^b(B)}(X_2, X_2)$$

given by composition. Note that  $G_1^{\text{op}} \rightarrow \text{End}_{K^b(B)}(R)$  comes from functoriality of  $E'$  and the quasi-isomorphism  $R \xrightarrow{q.i.} E' X_2$ .

• We have  $\Gamma_{21} : G_1 F[y] \oplus (G_2)_GF^2[y] \rightarrow G_1 F[y]$  given by  $(\text{Id}_{G_1 F[y]}, \omega')$ :

–  $I'_\beta : G_1 E[y]_GF^2[y] \rightarrow G_1 F[y]$  given by

$$\begin{aligned} g_1 e \otimes f_2 f_1 &\mapsto g_1 \otimes (f_1 \circ E f_2 \circ (e, y_1 e, 0)) \\ &= g_1 \otimes f_2(y_1 e).f_1, \end{aligned}$$

–  $I''_\beta : G_1 E[y]_GF^2[y] \rightarrow (G_2)_GF^2[y]$  given by

$$\begin{aligned} (\theta, \varphi_1) \otimes e \otimes f_2 f_1 &\mapsto ((e, y_1 e, 0) \circ (\theta, \varphi_1)) \otimes f_2 f_1 \\ &= (\theta e, \theta y_1 e, \varphi_1 \otimes e) \otimes f_2 f_1, \end{aligned}$$

–  $I'_\delta : (G_2)_GFF[y] \rightarrow G_1 F[y]$  given by

$$\begin{aligned} (e', e, \xi') \otimes f_2 f_1 &\mapsto ((0, f_2, 0) \circ (e', e, \xi')) \otimes f_1 \\ &= \left( E f_2 \circ (- \otimes e + y_2 \tau(- \otimes (e - y_1 e')) + y_1 y_2 \xi') \right) \otimes f_1 \\ &= (f_2(e), E f_2 \circ \tau(- \otimes e - y_1 e')) + E(f_2 \circ y_1) \circ \xi' \otimes f_1, \end{aligned}$$



$$- I''_\delta : (G_2)_G FF[y] \xrightarrow{\text{Id}} (G_2)_G F^2[y].$$

Let  $\omega' : (G_2)_G F^2[y] \rightarrow G_1 F[y]$  be the map  $I'_\delta$  after identifying  $FF[y]$  with  $F^2[y]$ . We show that  $\omega' \circ I''_\beta = I'_\beta$ :

$$\begin{aligned} \omega'((\theta e, \theta y_1 e, \varphi_1 \otimes e) \otimes f_2 f_1) &= (f_2(\theta y_1 e), E(f_2 \circ y_1) \circ (\varphi_1 \otimes e)) \otimes f_1 \\ &= (\theta f_2(y_1 e), \varphi_1 \cdot f_2(y_1 e)) \otimes f_1 \\ &= (\theta, \varphi_1) \cdot f_2(y_1 e) \otimes f_1 \\ &= I'_\beta((\theta, \varphi_1) \otimes e \otimes f_2 f_1). \end{aligned}$$

This shows that  $I'_\beta - I''_\beta = (\omega' - \text{Id})I''_\beta$ , whence  $\text{Im}(I'_\beta - I''_\beta) \subset \text{Im}(I'_\delta - I''_\delta)$ . It follows that the kernel of  $\Gamma_{21}$  is  $I_B + I_D$ .

- We have  $\Gamma_{12} : E[y]G_1 \oplus E^2[y]_G L_2 \rightarrow E[y]G_1$  given by  $(\text{Id}_{E[y]G_1}, \omega'')$ :
  - $I'_\beta : EE[y]_G L_2 \rightarrow E[y]G_1$  given by

$$\begin{aligned} e_1 e_2 \otimes (f', f, \rho') &\mapsto e_1 \otimes ((f', f, \rho') \circ (e_2, y_1 e_2, 0)) \\ &= e_1 \otimes (f(y_1 e_2) + f'(e_2), E f' \circ \tau(- \otimes e_2) + \rho'(- \otimes y_1 e_2)), \end{aligned}$$

$$- I''_\beta : EE[y]_G L_2 \xrightarrow{\text{Id}} E^2[y]_G L_2,$$

$$- I'_\delta : E^2[y]_G F[y]G_1 \rightarrow E[y]G_1 \text{ given (considering } I'_\delta \text{ of } \Gamma_{11}) \text{ by}$$

$$e_1 e_2 \otimes f_2 \otimes g_1 \mapsto e_1 \otimes f_2(y_1 e_2) \cdot g_1,$$

$$- I''_\delta : E^2[y]_G F[y]G_1 \rightarrow E^2[y]_G L_2 \text{ given by}$$

$$\begin{aligned} e_1 e_2 \otimes f \otimes (\theta, \varphi_1) &\mapsto e_1 e_2 \otimes ((\theta, \varphi_1) \circ (0, f, 0)) \\ &= e_1 e_2 \otimes (0, f \cdot \theta, \varphi_1 \circ E f). \end{aligned}$$

Let  $\omega'' : E^2[y]_G L_2 \rightarrow E[y]G_1$  be the map  $I'_\beta$  after identifying  $EE[y]$  with  $E^2[y]$ . We show that  $\omega'' \circ I''_\delta = I'_\delta$ :

$$\begin{aligned} \omega''(e_1 e_2 \otimes (0, f \cdot \theta, \varphi_1 \circ E f)) &= e_1 \otimes (f(y_1 e_2) \cdot \theta, (\varphi_1 \circ E f)(- \otimes y_1 e_2)) \\ &= e_1 \otimes (f(y_1 e_2) \cdot \theta, \varphi_1(- \cdot f(y_1 e_2))) \\ &= e_1 \otimes (f(y_1 e_2) \cdot \theta, f(y_1 e_2) \cdot \varphi_1) \\ &= e_1 \otimes f(y_1 e_2) \cdot (\theta, \varphi_1), \end{aligned}$$

while

$$\begin{aligned} I'_\delta(e_1 e_2 \otimes f \otimes (\theta, \varphi_1)) &= e_1 \otimes f(y_1 e_2) \cdot (\theta, \varphi_1) \\ &= (\omega'' \circ I''_\delta)(e_1 e_2 \otimes f \otimes (\theta, \varphi_1)). \end{aligned}$$

This shows that  $I'_\delta - I''_\delta = (\omega'' - \text{Id})I''_\delta$ , whence  $\text{Im}(I'_\delta - I''_\delta) \subset \text{Im}(I'_\beta - I''_\beta)$ . It follows that the kernel of  $\Gamma_{12}$  is  $I_B + I_D$ .

- We have  $\Gamma_{22} : G_1 G_1 \oplus (G_2)_G L_2 \rightarrow G_1 G_1 \oplus EF[y]$  given by  $\begin{pmatrix} \text{Id}_{G_1 G_1} & \omega''' \\ 0 & \kappa \end{pmatrix}$ :
  - $I'_\beta : G_1 E[y]_G L_2 \rightarrow G_1 G_1$  given (considering  $I'_\beta$  of  $\Gamma_{12}$ ) by

$$\begin{aligned} g_1 \otimes e \otimes (f', f, \rho') &\mapsto \\ g_1 \otimes (f'(e) + f(y_1 e), E f' \circ \tau(- \otimes e) + \rho'(- \otimes y_1 e)), & \end{aligned}$$

–  $I''_\beta : G_1 E[y]_G L_2 \rightarrow (G_2)_G L_2$  given by

$$(\theta, \varphi_1) \otimes e \otimes \ell_2 \mapsto (\theta e, \theta y_1 e, \varphi_1 \otimes e) \otimes \ell_2,$$

–  $I'_\delta : (G_2)_G F[y] G_1 \rightarrow G_1 G_1$  given (considering  $I'_\delta$  of  $\Gamma_{21}$ ) by

$$(e', e, \xi') \otimes f \otimes g_1 \mapsto (f(e), Ef \circ \tau(- \otimes e - y_1 e')) + E(f \circ y_1) \circ \xi' \otimes g_1,$$

–  $I''_\delta : (G_2)_G F[y] G_1 \rightarrow (G_2)_G L_2$  given by

$$g_2 \otimes f \otimes (\theta, \varphi_1) \mapsto g_2 \otimes (0, f.\theta, f \otimes \varphi_1).$$

We describe a map  $\omega''' : (G_2)_G L_2 \rightarrow G_1 G_1$ , and we show that  $\omega''' \circ I''_\beta = I'_\beta$  and  $\omega''' \circ I''_\delta = I'_\delta$ . Then we describe a decomposition of  $(G_2)_G L_2$  into  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -sub-bimodules,  $(G_2)_G L_2 \cong H \oplus EF[y]$ , where  $H$  is the image of  $I''_\beta + I''_\delta$ , and we let  $\kappa$  be the projection onto  $EF[y]$ , which lies in the kernel of  $\omega'''$ . It follows that the kernel of  $\Gamma_{22}$  is the image of  $I_B + I_D$ .

We first define a morphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $\bar{\omega}''' : G_2 \otimes_{A[y]} L_2 \rightarrow G_1 G_1$ . Considering that  $A[y] \hookrightarrow G_1^{\text{op}}$ , we show that  $\bar{\omega}'''$  descends to a morphism  $\omega''' : G_2 \otimes_{G_1^{\text{op}}} L_2 \rightarrow G_1 G_1$  by showing that it is middle-linear over generators of  $G_1^{\text{op}}$  in  $FE[y]$ . Let  $(e', e, \xi') \otimes (f', f, \rho') \in G_2 \otimes_{A[y]} L_2$  be an arbitrary simple tensor. For notational brevity we will write  $e_2$  for  $e - y_1 e'$ . Then define:

$$\begin{aligned} \bar{\omega}''' : (e', e, \xi') \otimes (f', f, \rho') &\mapsto \left( \varepsilon(e' \otimes f') + \varepsilon(e \otimes f), FE(\varepsilon \circ y_1 F)(\xi' \otimes f) \right. \\ &\quad \left. + FE\varepsilon(\xi' \otimes f') + \sigma(e \otimes f) - \sigma(y_1 e' \otimes f) \right) \otimes (1, 0) \\ &\quad + (1, 0) \otimes \left( 0, \varepsilon FE(e \otimes \rho') + \sigma(e' \otimes f') \right) \\ &\quad + \sigma FE(e \otimes \rho') - \sigma FE(y_1 e' \otimes \rho') + FE\sigma(\xi' \otimes f') + FE(\varepsilon \circ y_1 F)FE(\xi' \otimes \rho'). \end{aligned}$$

The last four terms, such as  $\sigma FE(e \otimes \rho')$ , give elements in  $FEFE[y]$ . These are interpreted in  $G_1 G_1$  using the decomposition

$$(5.4) \quad \begin{aligned} G_1 \otimes_{A[y]} G_1 &\xrightarrow{\sim} A[y]^{\text{op}} \oplus FE[y] \oplus FE[y] \oplus FEFE[y]. \\ (\theta, \varphi_1) \otimes (\theta', \varphi'_1) &\mapsto \theta \cdot \theta' \oplus \theta \cdot \varphi'_1 \oplus \varphi_1 \cdot \theta \oplus (\varphi_1 \otimes \varphi'_1). \end{aligned}$$

Each of the terms in this formula for  $\bar{\omega}'''$  is given by the application of a morphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules. We can also give a decomposition of  $G_2 \otimes_{A[y]} L_2$  into  $(A[y]^{\text{op}}, A[y]^{\text{op}})$ -bimodules:

$$\begin{aligned} G_2 \otimes_{A[y]} L_2 &\xrightarrow{\sim} EF[y]^{\oplus 4} \oplus FE^2 F[y]^{\oplus 2} \oplus EF^2 E[y]^{\oplus 2} \oplus FE^2 F^2 E[y] \\ (e', e, \xi') \otimes (f', f, \rho') &\mapsto (e' \otimes f', e' \otimes f, e \otimes f', e \otimes f) \\ &\quad \oplus (e' \otimes \rho', e \otimes \rho') \oplus (\xi' \otimes f', \xi' \otimes f) \oplus (\xi' \otimes \rho'). \end{aligned}$$

**Definition 5.3.** Using the ordered decompositions above, the map  $\bar{\omega}''' : G_2 \otimes_{A[y]} L_2 \rightarrow G_1 G_1$  is defined by the following matrix:

$$\begin{pmatrix} \varepsilon & 0 & 0 & \varepsilon & 0 & 0 & 0 & 0 & 0 \\ \sigma & 0 & 0 & 0 & 0 & \varepsilon FE & 0 & 0 & 0 \\ 0 & -\sigma \circ y_1 F & 0 & \sigma & 0 & 0 & FE\varepsilon & FE(\varepsilon \circ y_1 F) & 0 \\ 0 & 0 & 0 & 0 & -(\sigma \circ y_1 F)FE & \sigma FE & FE\sigma & 0 & FE(\varepsilon \circ y_1 F)FE \end{pmatrix}.$$

**Lemma 5.4.** *The map  $\bar{\omega}'''$  is middle-linear over the action of generators of  $G_1^{\text{op}}$  in  $FE[y]$ .*

*Proof.* We first compute the actions  $(e', e, \xi').\varphi_1$  and  $\varphi_1.(f', f, \rho')$  for  $\varphi_1 \in FE[y] \subset G_1^{\text{op}}$ ,  $(e', e, \xi') \in G_2$ , and  $(f', f, \rho') \in L_2$ . (Note that in the notation  $(\theta, \varphi) \in G_1^{\text{op}}$ , there is a relation  $\varphi = -\theta + y_1\varphi_1$ .) We have:

$$\begin{aligned} (e', e, \xi').\varphi_1 &= (\varphi_1(e), y_1\varphi_1(e), E\varphi_1 \circ \tau(-\otimes(e - y_1e')) + E(\varphi_1 y_1) \circ \xi') \\ \varphi_1.(f', f, \rho') &= (0, f \circ y_1\varphi_1 + f' \circ \varphi_1, Ef' \circ \tau \circ E\varphi_1 + \rho' \circ E(y_1\varphi_1)). \end{aligned}$$

One may compute the images under  $\bar{\omega}'''$  of

$$(\varphi_1(e), y_1\varphi_1(e), E\varphi_1 \circ \tau(-\otimes(e - y_1e')) + E(\varphi_1 y_1) \circ \xi') \otimes (f', f, \rho')$$

and

$$(e', e, \xi') \otimes (0, f \circ y_1\varphi_1 + f' \circ \varphi_1, Ef' \circ \tau \circ E\varphi_1 + \rho' \circ E(y_1\varphi_1))$$

using the formulas above to check that they agree.  $\square$

**Corollary 5.5.** *It follows from the lemma that  $\bar{\omega}'''$  determines a morphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $\omega''' : (G_2)_G L_2 \rightarrow G_1 G_1$ .*

We show next that  $I'_\beta = \omega''' \circ I''_\beta$  and  $I'_\delta = \omega''' \circ I''_\delta$ . The formula for  $\omega'''$  is determined by these conditions and may be derived from them. Evaluating the right side of the first equation:

$$\begin{aligned} &\omega''' \circ I''_\beta((\theta, \varphi_1) \otimes e \otimes (f', f, \rho')) \\ &= \omega'''((\theta e, \theta y_1 e, \varphi_1 \otimes e) \otimes (f', f, \rho')) \\ &= (f'(\theta e) + f(\theta y_1 e), \varphi_1.f(y_1 e) + \varphi_1.f'(e)) \otimes (1, 0) \\ &\quad + (1, 0) \otimes (0, \rho'(-\otimes \theta y_1 e) + Ef' \circ \tau(-\otimes \theta e)) \\ &\quad + (0, \varphi_1) \otimes (0, Ef' \circ \tau(-\otimes e) + \rho'(-\otimes y_1 e)), \\ &= (\theta.(f'(e) + f(y_1 e)), \varphi_1.(f'(e) + f(y_1 e))) \otimes (1, 0) \\ &\quad + (\theta, \varphi_1) \otimes (0, Ef' \circ \tau(-\otimes e) + \rho'(-\otimes y_1 e)) \\ &= (\theta, \varphi_1) \otimes (f'(e) + f(y_1 e), Ef' \circ \tau(-\otimes e) + \rho'(-\otimes y_1 e)), \\ &= I'_\beta((\theta, \varphi_1) \otimes e \otimes (f', f, \rho')). \end{aligned}$$

Now evaluating the right side of the second equation:

$$\begin{aligned}
 & \omega''' \circ I''_\delta((e', e, \xi') \otimes f \otimes (\theta, \varphi_1)) \\
 &= \omega'''((e', e, \xi') \otimes (0, f.\theta, f \otimes \varphi_1)) \\
 &= (f(e).\theta, E(f.\theta \circ y_1) \circ \xi' + E(f.\theta) \circ \tau(- \otimes (e - y_1 e'))) \otimes (1, 0) \\
 &\quad + (1, 0) \otimes (0, f(e).\varphi_1) + (0, Ef \circ \tau(- \otimes (e - y_1 e'))) \otimes (0, \varphi_1) \\
 &\quad + (0, E(fy_1) \circ \xi') \otimes (0, \varphi_1) \\
 &= (f(e), E(fy_1) \circ \xi' + E(f) \circ \tau(- \otimes (e - y_1 e'))) \otimes (\theta, 0) \\
 &\quad + (f(e), Ef \circ \tau(- \otimes (e - y_1 e')) + E(fy_1) \circ \xi') \otimes (0, \varphi_1) \\
 &= (f(e), Ef \circ \tau(- \otimes (e - y_1 e')) + E(fy_1) \circ \xi') \otimes (\theta, \varphi_1) \\
 &= I''_\delta((e', e, \xi') \otimes f \otimes (\theta, \varphi_1)).
 \end{aligned}$$

Consider the copy of  $EF[y]$  in  $(G_2)_{A[y]}L_2$  generated by terms of the form  $(0, e, 0) \otimes (f', 0, 0)$ , and let  $\bar{H}$  be the direct complement of this copy of  $EF[y]$  in  $(G_2)_{A[y]}L_2$ . The product  $(G_2)_GL_2$  is the quotient of the product  $(G_2)_{A[y]}L_2$  by the image of  $\gamma' - \gamma''$ , where:

–  $\gamma' : (G_2 \otimes FE[y]) \otimes L_2 \rightarrow (G_2)_{A[y]}L_2$  given by

$$\begin{aligned}
 & (e', e, \xi') \otimes \varphi_1 \otimes \ell_2 \mapsto \\
 & (\varphi_1(e), y_1\varphi_1(e), E\varphi_1 \circ \tau(- \otimes (e - y_1 e')) + E(\varphi_1 y_1) \circ \xi') \otimes \ell_2,
 \end{aligned}$$

–  $\gamma'' : G_2 \otimes (FE[y] \otimes L_2) \rightarrow (G_2)_{A[y]}L_2$  given by

$$\begin{aligned}
 & g_2 \otimes \varphi_1 \otimes (f', f, \rho') \mapsto \\
 & (0, f' \circ \varphi_1 + f \circ y_1\varphi_1, Ef' \circ \tau \circ E\varphi_1 + \rho' \circ E(y_1\varphi_1)).
 \end{aligned}$$

The images of  $\gamma'$  and  $\gamma''$  lie in  $\bar{H}$ , so the  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodule structure of  $(G_2)_GL_2$  is the sum  $H \oplus EF[y]$  where  $H$  is the quotient of  $\bar{H}$  by the image of  $\gamma' - \gamma''$ .

Now the image of  $I''_\beta$  includes every term of the form  $(e, y_1 e, \varphi_1 \otimes e) \otimes \ell_2$ , and the image of  $I''_\delta$  includes every term of the form  $g_2 \otimes (0, f, f \otimes \varphi_1)$ . By adding appropriate linear combinations of terms of the first form, one obtains any element  $(e, y_1 e, \xi') \otimes \ell_2$ , and similarly from terms of the second form one obtains any  $g_2 \otimes (0, f, \rho')$ . It follows that  $\text{Im}(I''_\beta + I''_\delta) = H$ .

**5.2. Maps  $\tilde{\rho}_\lambda$ : computation.** In this section we derive formulas for the maps  $\tilde{\rho}_\lambda$  in terms of the  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodule decompositions of the four coefficients of the matrix expressions of  $\tilde{E}\tilde{F}$  and  $\tilde{F}\tilde{E}$  and  $C$ .

5.2.1. *Map  $\tilde{\sigma}$ .* We begin by computing the map  $\tilde{\sigma} : \tilde{E}\tilde{F} \rightarrow \tilde{F}\tilde{E}$ . This map is defined already using  $\tilde{\eta}$  and  $\tilde{\varepsilon}$  and  $\tilde{\tau}$ , but we will need formulas for this map on each component of the matrix presentation of  $\tilde{E}\tilde{F}$ .

We use the following technique to derive the formulas. We start with an arbitrary element (or simple tensor) in a component of the matrix  $[\tilde{E}\tilde{F}]$ , as well as the element  $\tilde{\eta}(1) \in [\tilde{F}\tilde{E}]$ . Using the above computation of products, we represent the first element in the product of matrices  $[\tilde{E}] \cdot [\tilde{F}]$ , and the second

in  $[\tilde{F}] \cdot [\tilde{E}]$ . Now we have a term in the larger product  $[\tilde{F}] \cdot [\tilde{E}] \cdot [\tilde{E}] \cdot [\tilde{F}]$ . The inner components may be represented in  $[\tilde{E}^2]$ , and their image under  $\tilde{\tau}$  computed there and then expressed again in  $[\tilde{E}] \cdot [\tilde{E}]$ . The resulting element can be expressed in  $[\tilde{F}\tilde{E}] \cdot [\tilde{E}\tilde{F}]$ , and  $\tilde{\varepsilon}$  applied to the factor in  $[\tilde{E}\tilde{F}]$ , which lands in  $C$  and acts on the right on the remaining term in  $[\tilde{F}\tilde{E}]$ . We note that an element in  $[\tilde{E}\tilde{F}]$ , for example, will be expressed in  $[\tilde{E}] \cdot [\tilde{F}]$  by finding a representative in the product over  $\Delta$ . The final result will be independent of this choice.

- We have  $[\tilde{\sigma}]_{11} : [\tilde{E}\tilde{F}]_{11} \rightarrow [\tilde{F}\tilde{E}]_{11}$  given by  $\begin{pmatrix} \varepsilon \\ \sigma \end{pmatrix}$  using the decompositions:

$$\begin{aligned} & - [\tilde{E}\tilde{F}]_{11} \xrightarrow{\sim} EF[y], \\ & - [\tilde{F}\tilde{E}]_{11} \xrightarrow{\sim} (G_1)_G G_1 \xrightarrow{\sim} G_1 \xrightarrow{\sim} A[y] \oplus FE[y]. \end{aligned}$$

We take  $[\tilde{\eta}(1)]_{11} = (1, 0) \otimes (1, 0) \in (G_1)_G G_1 \cong [\tilde{F}\tilde{E}]_{11}$ , and an arbitrary  $e \otimes f \in EF[y] \cong [\tilde{E}\tilde{F}]_{11}$ . The product of these in  $[\tilde{F}\tilde{E}] \cdot [\tilde{E}\tilde{F}]$  can be represented in  $[\tilde{F}] \cdot [\tilde{E}] \cdot [\tilde{E}] \cdot [\tilde{F}]$  by

$$\begin{aligned} & \begin{pmatrix} 0 & (1,0) \\ 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (1,0) & 0 \end{pmatrix} \cdot \begin{pmatrix} e & 0 \\ 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix} \in \\ & \begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix} \cdot \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \cdot \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \cdot \begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix}. \end{aligned}$$

The middle factors produce  $(1, 0) \otimes e \in G_1 \otimes_{A[y]} E[y]$ . Passing through  $\Gamma_{21}$  of Eq. 5.1, this represents  $(e, y_1 e, 0) \in G_2 \cong [\tilde{E}^2]_{21}$ . Applying  $\tilde{\tau}$  yields  $(0, e, 0) \in G_2$ , which may be represented by

$$\begin{pmatrix} 0 & 0 \\ 0 & (0, e, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (1, 0) & 0 \end{pmatrix} \in [\tilde{E}] \cdot [\tilde{E}].$$

Then

$$\begin{pmatrix} 0 & 0 \\ (1, 0) & 0 \end{pmatrix} \cdot \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix} \xrightarrow{\tilde{\varepsilon}} \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} \in C$$

and

$$\begin{pmatrix} 0 & (1,0) \\ 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (0, e, 0) \end{pmatrix} \mapsto \begin{pmatrix} 0 & (0, e, 0) \\ 0 & 0 \end{pmatrix} \in [\tilde{F}\tilde{E}].$$

Finally, letting  $f \in C$  act on the right, we have

$$\begin{pmatrix} 0 & (0, e, 0) \\ 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} = \begin{pmatrix} (f(e), Ef \circ \tau(- \otimes e)) & 0 \\ 0 & 0 \end{pmatrix} \in \begin{pmatrix} G_1 & G_2 \\ L_2 & U \end{pmatrix}.$$

The coefficient in the upper right may be interpreted as  $(\varepsilon(e \otimes f), \sigma(e \otimes f))$ .

- We have  $[\tilde{\sigma}]_{21} : [\tilde{E}\tilde{F}]_{21} \rightarrow [\tilde{F}\tilde{E}]_{21}$  given by

$$\begin{pmatrix} 1 & 0 \\ 0 & F\varepsilon \\ 0 & F\sigma \end{pmatrix}$$

using the decompositions:

$$\begin{aligned} & - [\tilde{E}\tilde{F}]_{21} \xrightarrow{\sim} F[y] \oplus FEF[y], \\ & - [\tilde{F}\tilde{E}]_{21} \xrightarrow{\sim} F[y] \oplus F[y] \oplus F^2E[y]. \end{aligned}$$

Let us write

$$\eta(1) = \sum_a f_a \otimes e_a \in FE \subset FE[y].$$

The map

$$\text{Id}_R \in \text{Hom}_{K^b(B)}(R, R) \cong U$$

corresponds to the element  $[\tilde{\eta}(1)]_{22} \in (L_2)_G G_2$ , and it may be given in  $(L_2)_G G_2$  by the formula

$$\sum_a (f_a, 0, 0) \otimes (e_a, 0, 0) + \sum_b (0, f_b, 0) \otimes (0, e_b, 0).$$

This may be checked by composing the maps, i.e. considering the image under  $(L_2)_G G_2 \rightarrow U$ :

$$\begin{aligned} \sum_a (f_a, 0, 0) \otimes (e_a, 0, 0) &\mapsto (0, 0, 0, \text{Id}_{E[y]}, 0), \\ \sum_b (0, f_b, 0) \otimes (0, e_b, 0) &\mapsto (\text{Id}_{E[y]}, 0, 0, 0, 0), \end{aligned}$$

so the image is  $(\text{Id}_{E[y]}, 0, 0, \text{Id}_{E[y]}, 0)$ . Here the final terms  $\Lambda^\circ = 0$  are computed by composing the maps on the  $E^2[y]$  term in the degree 0 part of the upper row of  $R$ . For the first:

$$\begin{aligned} \Lambda &= \sum_a -y_2 \tau(- \otimes y_1 e_a) \circ E f_a \circ \tau \\ &= \sum_{a,d} -y_2 \tau y_1 (\tau(-)_{(1d)} \otimes f_a(\tau(-)_{(2d)}) \cdot e_a) \\ &= -y_2 \tau y_1 \tau = -y_2 \tau \end{aligned}$$

(in the second line we use a notation  $\tau(ee) = \sum_d \tau(ee)_{(1d)} \otimes \tau(ee)_{(2d)}$  for some choices of  $\tau(ee)_{(jd)}$ , and in the third line we use that  $\sum_a f_a(e') \cdot e_a = e'$ ). For the second:

$$\begin{aligned} \Lambda &= \sum_b (- \otimes e_b + y_2 \tau(- \otimes e_b)) \circ E f_b \\ &= \sum_b \tau y_1 (- \otimes e_b) \circ E f_b = \tau y_1. \end{aligned}$$

It follows from the formula for  $\Lambda^\circ$  (Eq. 3.3) that  $\Lambda^\circ = 0$  in both cases.

Then we take  $(\theta, \varphi_1) \otimes f \in G_1 F[y] \cong [\tilde{E}\tilde{F}]_{21}$ . Expressing everything in  $[\tilde{F}] \cdot [\tilde{E}] \cdot [\tilde{E}] \cdot [\tilde{F}]$ , we have:

$$\begin{aligned} &\sum_a \begin{pmatrix} 0 & 0 \\ 0 & (f_a, 0, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (e_a, 0, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (\theta, \varphi_1) & 0 \end{pmatrix} \cdot \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix} + \\ &\sum_b \begin{pmatrix} 0 & 0 \\ 0 & (0, f_b, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (0, e_b, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (\theta, \varphi_1) & 0 \end{pmatrix} \cdot \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix} \in \\ &\begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix} \cdot \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \cdot \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \cdot \begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix}. \end{aligned}$$

Now we must interpret  $\begin{pmatrix} 0 & 0 \\ 0 & (e_a, 0, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (\theta, \varphi_1) & 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 & 0 \\ 0 & (0, e_b, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (\theta, \varphi_1) & 0 \end{pmatrix}$  in  $[\tilde{E}^2]$  and apply  $\tilde{\tau}$ . We compute:

$$\begin{aligned} \Gamma_{21} : (e_a, 0, 0) \otimes (\theta, \varphi_1) &\mapsto (e_a, 0, 0) \cdot (\theta, \varphi_1) \\ &= (e_a \cdot \theta, 0, -E\varphi_1 \circ \tau(-\otimes y_1 e_a)) \in G_2 = [\tilde{E}^2]_{21} \\ &\xrightarrow{\tilde{\tau}} (0, e_a \cdot \theta, -\tau \circ E\varphi_1 \circ \tau(-\otimes y_1 e_a)), \\ \Gamma_{21} : (0, e_b, 0) \otimes (\theta, \varphi_1) &\mapsto (0, e_b, 0) \cdot (\theta, \varphi_1) \\ &= (\varphi_1(e_b), \varphi(e_b), E\varphi_1 \circ \tau(-\otimes e_b)) \in G_2 \\ &\xrightarrow{\tilde{\tau}} (0, \varphi_1(e_b), \tau \circ E\varphi_1 \circ \tau(-\otimes e_b)). \end{aligned}$$

We can represent these in  $[\tilde{E}] \cdot [\tilde{E}]$  using the natural map  $G_2 \xrightarrow{\sim} (G_2)_G G_1$ ,  $g_2 \mapsto g_2 \otimes (1, 0)$ . So, after applying  $\tilde{\tau}$  to the middle terms, we have:

$$\begin{aligned} &\sum_a \begin{pmatrix} 0 & 0 \\ 0 & (f_a, 0, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (0, e_a \cdot \theta, -\tau \circ E\varphi_1 \circ \tau y_1(-\otimes e_a)) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (1, 0) & 0 \end{pmatrix} \cdot \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix} + \\ &\sum_b \begin{pmatrix} 0 & 0 \\ 0 & (0, f_b, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (0, \varphi_1(e_b), \tau \circ E\varphi_1 \circ \tau(-\otimes e_b)) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (1, 0) & 0 \end{pmatrix} \cdot \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

Then  $\tilde{\varepsilon} : \begin{pmatrix} 0 & 0 \\ (1, 0) & 0 \end{pmatrix} \cdot \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix} \mapsto f \in F[y] \cong [C]_{21}$ , so by applying  $\tilde{F}\tilde{E}\tilde{\varepsilon}$  and viewing the first two factors in  $(L_2)_G G_2$  in  $[\tilde{F}\tilde{E}]_{22}$  we obtain:

$$\begin{pmatrix} 0 & 0 \\ 0 & \sum_a (f_a, 0, 0) \otimes (0, e_a \cdot \theta, -\tau \circ E\varphi_1 \circ \tau(-\otimes y_1 e_a)) \\ 0 & + \sum_b (0, f_b, 0) \otimes (0, \varphi_1(e_b), \tau \circ E\varphi_1 \circ \tau(-\otimes e_b)) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} \in [\tilde{F}\tilde{E}] \cdot C.$$

Then, finally, we express this element in  $[\tilde{F}\tilde{E}]$  by composing the first factors in  $L_2$  with the second factors in  $G_2$ , and also evaluating the action, which means post-composing with  $E'(f)$ :

$$\begin{aligned} &\sum_a (0, e_a \cdot \theta, -\tau \circ E\varphi_1 \circ \tau(-\otimes y_1 e_a)) \circ (f_a, 0, 0) \\ &= (0, 0, \theta, 0, -\tau \circ E\varphi_1 \circ \tau) \end{aligned}$$

where we have used

$$\begin{aligned} \Lambda &= \tau \cdot \theta - y_1 y_2 \tau \circ E\varphi_1 \circ \tau \text{ (pre-composing with } Ef_a \circ \tau \text{ and summing)} \\ &= \tau y_1 (E\theta \circ \tau) + y_1 y_2 \Lambda^\circ \text{ (formula 3.3)} \\ &= \tau \cdot \theta + y_1 y_2 (-\tau \circ E\varphi_1 \circ \tau) \text{ (equating with the first)}. \end{aligned}$$

Similarly:

$$\begin{aligned} & \sum_b (0, \varphi_1(e_b), \tau \circ E\varphi_1 \circ \tau(- \otimes e_b)) \circ (0, f_b, 0) \\ &= (\varphi_1, 0, 0, 0, \tau \circ E\varphi_1 \circ \tau) \end{aligned}$$

where we have used

$$\begin{aligned} \Lambda &= \tau y_1 \circ E\varphi_1 + y_1 y_2 \tau \circ E\varphi_1 \circ \tau \text{ (pre-composing with } Ef_b \text{ and summing)} \\ &= \tau y_1(E\varphi_1) + y_1 y_2 \Lambda^\circ \text{ (formula 3.3)} \\ &= \tau y_1 \circ E\varphi_1 + y_1 y_2 (\tau \circ E\varphi_1 \circ \tau) \text{ (equating with the first)}. \end{aligned}$$

Adding the two, we have

$$(\varphi_1, 0, \theta, 0, 0) \in FE[y]^{\oplus 4} \oplus F^2 E^2[y] \xrightarrow{\sim} U,$$

i.e.  $\Lambda^\circ = 0$  and  $\Lambda = \tau y_1(E\varphi_1 + E\theta \circ \tau) = \tau \circ E\varphi$ . Now post-composing with  $E'(f) = (0, f, 0) \in L_2$ , we obtain

$$(0, f, 0) \circ (\varphi_1, 0, \theta, 0, 0) = (\theta.f, f \circ \varphi_1, Ef \circ \tau \circ E\varphi_1)$$

where we have used

$$Ef \circ (\tau \circ E\theta + \tau y_1 \circ E\varphi_1) = E(\theta.f \circ \tau) + E(f \circ \varphi_1) + y_1 \circ Ef \circ \tau \circ E\varphi_1.$$

So the final expression is

$$\begin{pmatrix} 0 & 0 \\ (\theta.f, f \circ \varphi_1, Ef \circ \tau \circ E\varphi_1) & 0 \end{pmatrix} \in \begin{pmatrix} G_1 & G_2 \\ L_2 & U \end{pmatrix} = [\tilde{F}\tilde{E}].$$

The result follows because  $f \circ \varphi_1 = F\varepsilon(\varphi_1 \otimes f)$  and  $Ef \circ \tau \circ E\varphi_1 = F\sigma(\varphi_1 \otimes f)$ .

- We have  $[\tilde{\sigma}]_{12} : [\tilde{E}\tilde{F}]_{12} \rightarrow [\tilde{F}\tilde{E}]_{12}$  given by

$$\begin{pmatrix} 0 & \varepsilon E \\ 1 & y_1 \circ \varepsilon E \\ 0 & \sigma E \end{pmatrix}$$

using the decompositions:

- $E[y]G_1 \xrightarrow{\sim} E[y] \oplus EFE[y]$ ,
- $G_2 \xrightarrow{\sim} E[y] \oplus E[y] \oplus FE^2[y]$ .

We take  $[\tilde{\eta}(1)]_{11} = (1, 0) \otimes (1, 0) \in G_1 G_1 \cong [\tilde{F}\tilde{E}]_{11}$ , and an arbitrary  $e \otimes (\theta, \varphi_1) \in E[y]G_1 \cong [\tilde{E}\tilde{F}]_{12}$ . The product of these in  $[\tilde{F}\tilde{E}] \cdot [\tilde{E}\tilde{F}]$  can be represented in  $[\tilde{F}] \cdot [\tilde{E}] \cdot [\tilde{E}] \cdot [\tilde{F}]$  by

$$\begin{pmatrix} 0 & (1, 0) \\ 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (1, 0) & 0 \end{pmatrix} \cdot \begin{pmatrix} e & 0 \\ 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & (\theta, \varphi_1) \\ 0 & 0 \end{pmatrix},$$

and application of  $\tilde{F}\tilde{\tau}\tilde{F}$  gives

$$\begin{pmatrix} 0 & (1, 0) \\ 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (0, e, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (1, 0) & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & (\theta, \varphi_1) \\ 0 & 0 \end{pmatrix}.$$

This is sent by  $\tilde{F}\tilde{E}\tilde{\varepsilon}$  to

$$\begin{pmatrix} 0 & (0, e, 0) \\ 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (\theta, \varphi_1) \end{pmatrix} \in [\tilde{F}\tilde{E}] \cdot [C]$$



which gives  $(\varphi_1(e), \varphi(e), E\varphi_1 \circ \tau(- \otimes e)) \in G_2 \cong [\tilde{F}\tilde{E}]_{12}$ . The result follows from the fact that  $\varphi = -\theta + y_1\varphi_1$  and  $E\varphi_1 \circ \tau(- \otimes e) = \sigma E(e \otimes \varphi_1)$ .

- We have  $[\tilde{\sigma}]_{22} : [\tilde{E}\tilde{F}]_{22} \rightarrow [\tilde{F}\tilde{E}]_{22}$  given by

$$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & F\varepsilon E & 0 \\ \eta & y_1 & 0 & 0 & \sigma \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & F\sigma E & 0 \end{pmatrix}$$

using the ordered decompositions:

–  $G_1G_1 \oplus EF[y] \xrightarrow{\sim} A[y] \oplus FE[y] \oplus FE[y] \oplus FEFE[y] \oplus EF[y]$  determined from the ordered decomposition of Eq. 5.4,

–  $U \xrightarrow{\sim} FE[y] \oplus FE[y] \oplus FE[y] \oplus FE[y] \oplus F^2E^2[y]$  given in Prop. 3.18.

We compute  $[\tilde{\sigma}]_{22}$  first on  $G_1G_1$ , and afterwards on  $EF[y]$ . Let  $(\theta, \varphi_1) \otimes (\theta', \varphi'_1) \in G_1G_1$  be an arbitrary simple tensor. We can use the same presentation for  $[\tilde{\eta}(1)]_{22}$  as in the calculations for  $[\tilde{\sigma}]_{21}$ , and then the presentation for everything in  $[\tilde{F}] \cdot [\tilde{E}] \cdot [\tilde{E}] \cdot [\tilde{F}]$  is:

$$\begin{aligned} & \sum_a \begin{pmatrix} 0 & 0 \\ 0 & (f_a, 0, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (e_a, 0, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (\theta, \varphi_1) & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & (\theta', \varphi'_1) \\ 0 & 0 \end{pmatrix} + \\ & \sum_b \begin{pmatrix} 0 & 0 \\ 0 & (0, f_b, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (0, e_b, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ (\theta, \varphi_1) & 0 \end{pmatrix} \cdot \begin{pmatrix} 0 & (\theta', \varphi'_1) \\ 0 & 0 \end{pmatrix} \in \\ & \begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix} \cdot \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \cdot \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \cdot \begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix}. \end{aligned}$$

Using again the calculations for  $[\tilde{\sigma}]_{21}$ , we see that applying  $\tilde{F}(\tilde{E}\tilde{\varepsilon} \circ \tilde{\tau}\tilde{F})$  yields:

$$\begin{pmatrix} 0 & 0 \\ 0 & (\varphi_1, 0, \theta, 0, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (\theta', \varphi'_1) \end{pmatrix} \in [\tilde{F}\tilde{E}] \cdot C.$$

This time we compute the action of  $(\theta', \varphi'_1)$  on the right on  $U$  using the action of  $G_1^{\text{op}}$  on the right on  $R$ . This is given on the degree 0 term of the top row by composing with  $E\varphi'$ , and on the degree 1 part of the bottom row by multiplying by the matrix  $\begin{pmatrix} \varphi' & 0 \\ \varphi'_1 & \theta' \end{pmatrix}$ . So we have

$$\begin{pmatrix} \varphi' & 0 \\ \varphi'_1 & \theta' \end{pmatrix} \cdot \begin{pmatrix} \varphi_1 & \theta \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} \varphi' \circ \varphi_1 & \varphi' \circ -\theta \\ \varphi'_1 \circ \varphi_1 & \varphi'_1 \circ -\theta \end{pmatrix},$$

and

$$\begin{aligned}
 \Lambda &= E\varphi' \circ \tau \circ E\varphi \\
 &= E\theta' \circ \tau \circ E\theta + E\theta' \circ \tau \circ E(y_1\varphi_1) \\
 &\quad + E(y_1\varphi'_1) \circ \tau \circ E\theta + E(y_1\varphi'_1) \circ \tau \circ E(y_1\varphi_1), \\
 \Lambda &= \tau y_1(E(\varphi' \circ \varphi_1) + E(\varphi' \circ \theta) \circ \tau) \\
 &\quad - y_2\tau y_1(E(\varphi'_1 \circ \varphi_1) + E(\varphi'_1 \circ \theta) \circ \tau) + y_1 y_2 \Lambda^\circ \\
 &= \tau y_1 \circ E\varphi' \circ (E\varphi_1 + E\theta \circ \tau) \\
 &\quad - y_2\tau \circ E(y_1\varphi'_1) \circ (E\varphi_1 + E\theta \circ \tau) + y_1 y_2 \Lambda^\circ \\
 &= \tau y_1 \circ E\theta' \circ (E\varphi_1 + E\theta \circ \tau) \\
 &\quad + E(y_1\varphi'_1) \circ (E\varphi_1 + E\theta \circ \tau) + y_1 y_2 \Lambda^\circ.
 \end{aligned}$$

So (inspecting the difference)  $\Lambda^\circ = E\varphi'_1 \circ \tau \circ E\varphi_1$ , and the final expression in the  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodule form of  $U$  (Prop. 3.18) is:

$$\begin{pmatrix} 0 & 0 \\ 0 & (\varphi' \circ \varphi_1, \varphi'_1 \circ \varphi_1, \theta \cdot \varphi', \theta \cdot \varphi'_1, E\varphi'_1 \circ \tau \circ E\varphi_1) \end{pmatrix} \in [\tilde{F}\tilde{E}].$$

Using the fact that  $E\varphi'_1 \circ \tau \circ E\varphi_1 = F\sigma E(\varphi_1 \otimes \varphi'_1)$ , and  $\varphi'_1 \circ \varphi_1 = F\varepsilon E(\varphi_1 \otimes \varphi'_1) \in FE[y]$ , one recovers the first four columns of the matrix  $[\tilde{\sigma}]_{22}$ .

For the fifth column, we start with an arbitrary simple tensor  $e \otimes f' \in EF[y] \subset [\tilde{E}\tilde{F}]_{22}$ . This element can be represented by (i.e. has a preimage under  $\kappa$  given by)

$$(0, e, 0) \otimes (f', 0, 0) \in (G_2)_G L_2,$$

so we study now the element

$$\begin{aligned}
 &\sum_a \begin{pmatrix} 0 & 0 \\ 0 & (f_a, 0, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (e_a, 0, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (0, e, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (f', 0, 0) \end{pmatrix} + \\
 &\sum_b \begin{pmatrix} 0 & 0 \\ 0 & (0, f_b, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (0, e_b, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (0, e, 0) \end{pmatrix} \cdot \begin{pmatrix} 0 & 0 \\ 0 & (f', 0, 0) \end{pmatrix} \in \\
 &\begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix} \cdot \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \cdot \begin{pmatrix} E[y] & E^2[y] \\ G_1 & G_2 \end{pmatrix} \cdot \begin{pmatrix} F[y] & L_1 \\ F^2[y] & L_2 \end{pmatrix}.
 \end{aligned}$$

Apply  $\Gamma_{22}$  of  $[\tilde{E}\tilde{E}]$  to the elements  $(e_a, 0, 0) \otimes (0, e, 0)$  and  $(0, e_b, 0) \otimes (0, e, 0)$ . Observe first that under the isomorphism of Prop. 3.18, we have

$$\begin{aligned}
 (e_a, 0, 0) &\mapsto (0, -y_1 e_a, -\tau(-\otimes y_1 e_a)) \\
 (0, e_b, 0) &\mapsto (e_b, e_b, \tau y_1(-\otimes e_b)) \\
 (0, e, 0) &\mapsto (e, e, \tau y_1(-\otimes e)).
 \end{aligned}$$

The formula for  $\Gamma_{22}$  may be applied to the elements on the right to find representatives in  $G_2$ , where  $[\tilde{\tau}]_{22}$  has been defined [McM, Def. 4.4]. We

obtain:

$$\begin{aligned}
 & (0, -y_1 e_a, 0) \otimes (e, e, 0) \\
 & \xrightarrow{\Gamma_{22}} (0, -y_2(e_a \otimes e), -y_2(e_a \otimes e), 0) \\
 & \xrightarrow{[\tilde{\tau}]_{22}} (e_a \otimes e, e_a \otimes e, -\tau y_2(e_a \otimes e), 0), \\
 & (e_b, e_b, 0) \otimes (e, e, 0) \\
 & \xrightarrow{\Gamma_{22}} (\tau y_1(e_b \otimes e), e_b \otimes e, e_b \otimes e, 0) \\
 & \xrightarrow{[\tilde{\tau}]_{22}} (\tau(e_b \otimes e), \tau(e_b \otimes e), \tau(e_b \otimes e), 0).
 \end{aligned}$$

Here the components  $(ee_1, ee_2, ee_3, \chi'') \in G_2$  are interpreted using [McM, Prop. 3.16], except we record the data of  $\chi''$  in place of the data of  $\chi$  according to [McM, Prop. 3.21]:

$$\chi = - \otimes ee_1 + y_3 \circ \tau E(- \otimes ee_2) + y_2 y_3 \circ E \tau \circ \tau E(- \otimes ee_3) + y_1 y_2 y_3 \chi''.$$

The next step is to express  $(e_a e, e_a e, -\tau y_2(e_a e), 0)$  and  $(\tau(e_b e), \tau(e_b e), \tau(e_b e), 0)$  back in  $(G_2)_G G_2$  (i.e. find a preimage under  $\Gamma_{22} |_{(G_2)_G G_2}$ ) in order to view them in  $[\tilde{E}] \cdot [\tilde{E}]$ . We will need the notation  $\tau(ee) = \sum_d \tau(ee)_{(1d)} \otimes \tau(ee)_{(2d)}$  introduced to compute  $[\tilde{\sigma}_{21}]$  above.

*Claim.* We have:

$$\begin{aligned}
 \Gamma_{22} : & \sum_d (0, \tau(e_a e)_{(1d)}, 0) \otimes (\tau(e_a e)_{(2d)}, y_1 \tau(e_a e)_{(2d)}, 0) \\
 & - \sum_d (0, \tau y_1(e_a e)_{(1d)}, 0) \otimes (0, \tau y_1(e_a e)_{(2d)}, 0) \\
 & \mapsto (e_a e, e_a e, -\tau y_2(e_a e), 0), \\
 \Gamma_{22} : & \sum_d (0, \tau(e_b e)_{(1d)}, 0) \otimes (0, \tau(e_b e)_{(2d)}, 0) \\
 & \mapsto (\tau(e_b e), \tau(e_b e), \tau(e_b e), 0)
 \end{aligned}$$

where  $\Gamma_{22}$  is the one computed for  $[\tilde{E}] \cdot [\tilde{E}]$  using elements described in the form  $(e', e, \xi') \in G_2$ .

*Proof.* A direct calculation using the formula for  $\Gamma_{22} |_{(G_2)_G G_2}$ .  $\square$

We have now an element of  $[\tilde{F}] \cdot [\tilde{E}] \cdot [\tilde{E}] \cdot [\tilde{F}]$ , and we need to apply  $[\tilde{F}] \cdot [\tilde{E}] \cdot \tilde{\varepsilon}$  and then realize the result in  $[\tilde{F}\tilde{E}]$ . Observe that

$$(0, e, 0) \otimes (f', 0, 0) \xrightarrow{\tilde{\varepsilon}} 0.$$

It follows that

$$\begin{aligned}
 & (0, \tau y_1(e_a e)_{(2d)}, 0) \otimes (f', 0, 0) \mapsto 0, \\
 & (0, \tau(e_b e)_{(2d)}, 0) \otimes (f', 0, 0) \mapsto 0.
 \end{aligned}$$

Therefore only one of the three terms will remain, namely the term with factor  $(\tau(e_a e)_{(2d)}, y_1 \tau(e_a e)_{(2d)}, 0)$ . We have

$$\begin{aligned} & (\tau(e_a e)_{(2d)}, y_1 \tau(e_a e)_{(2d)}, 0) \otimes (f', 0, 0) \\ & \xrightarrow{\tilde{\varepsilon}} \left( f'(\tau(e_a e)_{(2d)}), E f' \circ \tau \circ (- \otimes y_1 \tau(e_a e)_{(2d)}) \right) \\ & =: (\theta, \varphi)_{a,d} \in G_1. \end{aligned}$$

Expanding the second component, we have

$$\begin{aligned} & E f' \circ \tau y_1 (- \otimes \tau(e_a e)_{(2d)}) \\ & = - \otimes f'(\tau(e_a e)_{(2d)}) + y_1 (E f' \circ \tau(- \otimes \tau(e_a e)_{(2d)})), \end{aligned}$$

so

$$(\varphi_1)_{a,d} = E f' \circ \tau(- \otimes \tau(e_a e)_{(2d)}).$$

Now we compose the first two factors (i.e. apply  $\Gamma_{22} |_{(L_2)_G G_2}$  from  $[\tilde{F}] \cdot [\tilde{E}]$ ) to obtain an element of  $[\tilde{F}\tilde{E}]_{22}$ , i.e. of  $U \cong \text{End}_{K^b(B)}(R)$ , and we use the right action of  $G_1$  on  $R$  to compute the action of  $(\theta, \varphi)_{a,d}$  on this element. As we have described, this right action is given by composing the element of  $U$  with a certain morphism of  $\text{End}_{K^b(B)}(R)$  determined by the morphism  $E'(\theta, \varphi)_{a,d} \in \text{End}_{K^b(B)}(E'X_2)$  given by functoriality of  $E'$ . The formulas are simpler if we perform the two compositions together. The resulting element of  $U$  determines an endomorphism of the degree 0 term  $E^2[y]$  of the top row of  $R$  as follows:

$$\begin{aligned} & E^2 f' \circ E \tau(- \otimes y_1 \tau(e_a e)_{(2d)}) \\ & \circ (\tau y_1 (- \otimes \tau(e_a e)_{(1d)})) \circ (E f_a \circ \tau) \\ & = E^2 f' \circ E \tau \circ \left( \tau y_1 (E f_a \circ \tau(-)) \otimes \tau(e_a e)_{(1d)} \otimes y_1 \tau(e_a e)_{(2d)} \right) \\ & = E^2 f' \circ E \tau \circ \tau E \circ y_2 y_1 (E f_a \circ \tau(-)) \otimes \tau(e_a e)_{(1d)} \otimes \tau(e_a e)_{(2d)} \end{aligned}$$

Summing over  $d$  and  $a$  we obtain:

$$\begin{aligned} & \underbrace{\sum_{a,d}}_{\rightsquigarrow} E^2 f' \circ E \tau \circ \tau E \circ y_2 y_1 (E \tau \circ \tau E (- \otimes e)) \\ & = E^2 f' \circ E \tau \circ \tau E \circ E \tau(- \otimes y_1 e) =: \Lambda. \end{aligned}$$

Now the matrix giving the morphism on the degree 1 term of the bottom row of  $R$  arising from the element

$$\Gamma_{22}((f_a, 0, 0) \otimes (0, \tau(e_a e)_{(1d)}, 0)) \in U$$

is

$$\begin{pmatrix} 0 & f_a(-) \cdot \tau(e_a e)_{(1d)} \\ 0 & 0 \end{pmatrix}.$$

Taking the product we have:

$$\begin{aligned} & \begin{pmatrix} Ef' \circ \tau(- \otimes y_1 \tau(e_a e)_{(2d)}) & 0 \\ Ef' \circ \tau(- \otimes \tau(e_a e)_{(2d)}) & f'(\tau(e_a e)_{(2d)}) \end{pmatrix} \cdot \begin{pmatrix} 0 & f_a(-) \cdot \tau(e_a e)_{(1d)} \\ 0 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & Ef' \circ \tau(f_a(-) \cdot \tau(e_a e)_{(1d)} \otimes y_1 \tau(e_a e)_{(2d)}) \\ 0 & Ef' \circ \tau(f_a(-) \cdot \tau(e_a e)_{(1d)} \otimes \tau(e_a e)_{(2d)}) \end{pmatrix}. \end{aligned}$$

Summing over  $a$  and  $d$  we find:

$$\underbrace{\sum_{a,d}}_{\rightarrow} \begin{pmatrix} 0 & Ef' \circ \tau \circ y_1 \tau(- \otimes e) \\ 0 & Ef' \circ \tau(\tau(- \otimes e)) \end{pmatrix} = \begin{pmatrix} 0 & Ef' \circ \tau(- \otimes e) \\ 0 & 0 \end{pmatrix}.$$

We need the remainder term  $\Lambda^\circ$  to find the image in the  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodule expression for  $U$ . Observe that since  $\Phi_{11} = \Phi_{21} = \Phi_{22} = 0$ , we have by Prop. 3.3:

$$\begin{aligned} \Lambda &= \tau y_1 \circ (E^2 f' \circ E \tau(- \otimes e) \circ \tau) + y_1 y_2 \Lambda^\circ \\ &= \tau y_1 \circ E^2 f' \circ E \tau \circ \tau E(- \otimes e) + y_1 y_2 \Lambda^\circ \\ &= y_2 E^2 f' \circ \tau E \circ E \tau \circ \tau E(- \otimes e) \\ &\quad + E^2 f' \circ E \tau \circ \tau E(- \otimes e) + y_1 y_2 \Lambda^\circ, \end{aligned}$$

while from the above:

$$\begin{aligned} \Lambda &= E^2 f' \circ E \tau \circ \tau E \circ E \tau(- \otimes y_1 e) \\ &= E^2 f' \circ E \tau \circ \tau E \circ y_2 E \tau(- \otimes e) \\ &\quad + E^2 f' \circ E \tau \circ \tau E(- \otimes e) \\ &= E^2 f' \circ E \tau \circ y_3 \tau E \circ E \tau(- \otimes e) \\ &\quad + E^2 f' \circ E \tau \circ E \tau(- \otimes e) \\ &\quad + E^2 f' \circ E \tau \circ \tau E(- \otimes e) \\ &= y_2 E^2 f' \circ E \tau \circ \tau E \circ E \tau(- \otimes e) \\ &\quad + E^2 f' \circ E \tau \circ \tau E(- \otimes e), \end{aligned}$$

so  $\Lambda^\circ = 0$  because  $E \tau \circ \tau E \circ E \tau = \tau E \circ E \tau \circ \tau E$ , and the final expression is

$$(0, 0, Ef' \circ \tau(- \otimes e), 0, 0) \in U \cong [\tilde{F} \tilde{E}]_{22}.$$

Observe that  $\sigma(e \otimes f') = Ef' \circ \tau(- \otimes e)$ . From this we have the fifth column of the matrix of  $[\tilde{\sigma}]_{22}$  as a map of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules, and the justification of all components of that matrix is now complete.

5.2.2. *Maps  $\tilde{\varepsilon} \circ \tilde{x}^i \tilde{F}$  and  $\tilde{F} \tilde{x}^i \circ \tilde{\eta}$ .* We continue by computing the maps  $\tilde{\varepsilon} \circ \tilde{x}^i \tilde{F}$  and  $\tilde{F} \tilde{x}^i \circ \tilde{\eta}$  on the various components of the matrices  $[\tilde{F} \tilde{E}]$  and  $[\tilde{E} \tilde{F}]$ .

- We have  $[\tilde{\varepsilon} \circ \tilde{x}^i \tilde{F}]_{11} : [\tilde{E} \tilde{F}]_{11} \rightarrow [C]_{11}$  given by  $\varepsilon \circ (x^i y_1) F$  using the decompositions:
  - $[\tilde{E} \tilde{F}]_{11} \xrightarrow{\sim} EF[y]$ ,
  - $[C]_{11} \xrightarrow{\sim} A[y]$ .

The endomorphism  $\tilde{x} \in \text{End}(\tilde{E})$  given in [McM, Def. 4.4] determines an endomorphism of  $[\tilde{E}\tilde{F}]_{11}$  given by  $xF$  on  $EF[y]$ . The morphism  $\tilde{\varepsilon}$  performs the composition of elements in  $\tilde{E}$  and  $\tilde{F}$  when these are interpreted in  $\text{Hom}_{K^b(B)}(X, E'X)$  and  $\text{Hom}_{K^b(B)}(E'X, X)$ . In particular,  $e \in E[y]$  represents the morphism given by  $1 \mapsto y_1e$  in degree 0 of the top row of  $X_1$  and  $E'X_1$ , and  $f \in F[y]$  represents the morphism given by  $e \mapsto f(e)$  in the same place.

- We have  $[\tilde{F}\tilde{x}^i \circ \tilde{\eta}]_{11} : [C]_{11} \rightarrow [\tilde{F}\tilde{E}]_{11}$  given by:

$$\left( \begin{array}{c} y^i \\ F(h_{i-1}(x, y)) \circ \eta \end{array} \right)$$

using the decompositions:

- $[C]_{11} \xrightarrow{\sim} A[y]$ ,
- $[\tilde{F}\tilde{E}]_{11} \xrightarrow{\sim} A[y] \oplus FE[y]$ .

Here  $h_i(A_1, \dots, A_n)$  is the complete homogeneous symmetric polynomial of degree  $i$  in the variables  $\{A_1, \dots, A_n\}$ . Observe first that  $[\tilde{\eta}]_{11}$  is given by  $1 \mapsto \text{Id}_{E[y]} \in G_1$ , and  $\text{Id}_{E[y]} = (1, 0)$  in the bimodule presentation of  $G_1$ . According to [McM, Def. 4.4],  $[\tilde{x}]_{11}$  acts on  $G_1$  (in its submodule presentation) by  $\tilde{x}^i \cdot (\theta, \varphi) = (y^i\theta, x^i \circ \varphi)$ . One computes:

$$\begin{aligned} x^i \circ \varphi &= x^i \cdot \theta + x^i y_1 \varphi_1 \\ &= y^i \theta + (x^i - y^i) \cdot \theta + y_1 x^i \varphi_1 \\ &= y^i \theta + y_1 (h_{i-1}(x, y) \cdot \theta + x^i \circ \varphi_1), \end{aligned}$$

so in the bimodule presentation we have  $\tilde{x}^i \cdot (\theta, \varphi_1) = (y^i\theta, h_{i-1}(x, y) \cdot \theta + x^i \circ \varphi_1)$ . In particular,  $\tilde{x}^i \cdot (1, 0)$  has the specified form.

Note the small case interpretations:

$$\begin{cases} h_{i-1}(x, y) = 0 & i = 0 \\ h_{i-1}(x, y) = 1 & i = 1 \\ h_{i-1}(x, y) = x + y & i = 2 \\ \dots & \dots \end{cases}$$

- We have  $[\tilde{\varepsilon} \circ \tilde{x}^i \tilde{F}]_{21} : [\tilde{E}\tilde{F}]_{21} \rightarrow [C]_{21}$  given by  $(x^i, F(\varepsilon \circ (x^i y_1) F))$  using the decompositions:

- $[\tilde{E}\tilde{F}]_{21} \xrightarrow{\sim} F[y] \oplus FEF[y]$ ,
- $[C]_{21} \xrightarrow{\sim} F[y]$ .

(Note that  $x.f = f \circ x$  for  $f \in F[y]$ .) The endomorphism  $[\tilde{x}]_{21}$  acts on  $G_1$  as described above:  $x^i \circ \varphi = x^i \cdot \theta + x^i y_1 \varphi_1$ . Then

$$\begin{aligned} G_1 \otimes_{A[y]} F[y] &\xrightarrow{[\tilde{\varepsilon}]_{21}} F[y], \\ (\tilde{x}^i \cdot (\theta, \varphi_1)) \otimes f &\mapsto f \circ x^i \cdot \theta + f \circ x^i y_1 \varphi_1, \end{aligned}$$

and the component data follows immediately.

- We have  $[\tilde{F}\tilde{x}^i \circ \tilde{\eta}]_{21} : [C]_{21} \rightarrow [\tilde{F}\tilde{E}]_{21}$  given by:

$$\begin{pmatrix} 0 \\ y^i \\ F(F(h_{i-1}(x, y)) \circ \eta) \end{pmatrix}$$

using the decompositions:

- $[C]_{21} \xrightarrow{\sim} F[y]$ ,
- $[\tilde{F}\tilde{E}]_{21} \xrightarrow{\sim} F[y] \oplus F[y] \oplus F^2E[y]$ .

Let

$$\begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} \in \begin{pmatrix} A^{\text{op}}[y] & E[y] \\ F[y] & G_1 \end{pmatrix} = [C],$$

and observe that

$$\begin{aligned} \tilde{\eta} \left( \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} \right) &= \tilde{\eta} \left( \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right) \\ &= \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} \cdot \tilde{\eta} \left( \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right) \\ &= \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} \cdot \begin{pmatrix} (1, 0) & 0 \\ 0 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & 0 \\ (0, f, 0) & 0 \end{pmatrix} \in \begin{pmatrix} G_1 & G_2 \\ L_2 & U \end{pmatrix} = [\tilde{F}\tilde{E}]. \end{aligned}$$

Here  $(0, f, 0)$  is given in the bimodule form of  $L_2$ . Elements  $f \in F[y]$  act on  $X_2$  by  $f \cdot e \mapsto f(e)$  for  $e$  in the degree 0 part of the top row of  $X_2$ . These elements act on  $E'X_2$  by  $Ef$  in the degree 0 part of the top row of  $E'X_2$ , and by  $(0, f) : E[y] \oplus E[y] \rightarrow A[y]$  in the degree 1 part of the bottom row of  $E'X_2$ .

Now we compute  $[\tilde{F}\tilde{x}^i]_{21}$  on these elements. Take

$$(0, f, 0) \otimes (1, 0) \in (L_2)_G G_1 \cong [\tilde{F}\tilde{E}]_{21}.$$

This arises from  $[\tilde{F}]_{22} \cdot [\tilde{E}]_{21}$ . We have already seen that

$$\tilde{x}^i \cdot (1, 0) = (y^i, h_{i-1}(x, y)),$$

and we then view this in  $L_2$  using the action of  $G_1$  on  $L_2$  on the right, which is given by composition. We obtain  $(0, y^i f, x_1^i \circ Ef)$  in submodule form, or

$$(0, y^i f, h_{i-1}(x_1, y) \circ Ef)$$

in bimodule form using:

$$\begin{aligned} x_1^i \circ Ef &= (y^i + y_1 h_{i-1}(x_1, y)) \circ Ef \\ &= E(y^i f) + y_1 \rho', \end{aligned}$$

so

$$\begin{aligned} \rho' &= h_{i-1}(x_1, y) \circ Ef \\ &= F(h_{i-1}(x, y) \circ \eta)(f). \end{aligned}$$

- We have  $[\tilde{\varepsilon} \circ \tilde{x}^i \tilde{F}]_{12} : [\tilde{E}\tilde{F}]_{12} \rightarrow [C]_{12}$  given by  $(x^i, (\varepsilon \circ (x^i y_1) F) E)$  using the decompositions:

- $[\tilde{E}\tilde{F}]_{12} \xrightarrow{\sim} E[y] \oplus EFE[y]$ ,
- $[C]_{12} \xrightarrow{\sim} E[y]$ .

The endomorphism  $[\tilde{x}]_{12}$  acts on  $E[y] \cong [\tilde{E}]_{11}$  as  $x$ , and thus on  $E[y]G_1 \cong [\tilde{E}\tilde{F}]_{12}$  as  $xG_1$ . Then

$$\begin{aligned} E[y] \otimes_{A[y]} G_1 &\xrightarrow{[\tilde{\varepsilon}]_{12}} E[y], \\ x^i(e) \otimes (\theta, \varphi_1) &\mapsto \varphi(x^i y_1 e) \\ &= y_1(x^i e \theta + \varphi_1(y_1 x^i e)), \end{aligned}$$

and the component data follows immediately. (Recall that  $e \in E[y]$  on either side corresponds to the map  $A[y] \rightarrow E[y]$ ,  $1 \mapsto y_1 e$ .)

- We have  $[\tilde{F}\tilde{x}^i \circ \tilde{\eta}]_{12} : [C]_{12} \rightarrow [\tilde{F}\tilde{E}]_{12}$  given by:

$$\begin{pmatrix} y^i \\ y^i y_1 \\ (F(h_{i-1}(x, y)) \circ \eta) E \end{pmatrix}$$

using the decompositions:

- $[C]_{12} \xrightarrow{\sim} E[y]$ ,
- $[\tilde{F}\tilde{E}]_{12} \xrightarrow{\sim} E[y] \oplus E[y] \oplus FE^2[y]$ .

By reasoning that is parallel to the  $[\tilde{F}\tilde{x}^i \circ \tilde{\eta}]_{21}$  case, we have:

$$\tilde{\eta} : \begin{pmatrix} 0 & e \\ 0 & 0 \end{pmatrix} \mapsto \begin{pmatrix} 0 & (e, y_1 e, 0) \\ 0 & 0 \end{pmatrix} \in \begin{pmatrix} G_1 & G_2 \\ L_2 & U \end{pmatrix} = [\tilde{F}\tilde{E}],$$

using again the bimodule form of  $G_2$ . We also have

$$(e, y_1 e, 0) \mapsto (1, 0) \otimes (e, y_1 e, 0) \in (G_1)_G G_2$$

under the canonical isomorphism  $G_2 \xrightarrow{\sim} (G_1)_G G_2$ , giving a presentation of  $[\tilde{\eta}]_{12}(e)$  in  $[\tilde{F}]_{12} \cdot [\tilde{E}]_{22}$ . We can calculate  $[\tilde{F}\tilde{x}^i]_{12}$  on an element  $(1, 0) \otimes g_2$  by calculating  $[\tilde{x}^i]_{12}$  on  $g_2$ .

The element  $(e, y_1 e, 0)$  (bimodule form) corresponds to the element  $(e, 0, - \otimes y_1 e)$  (submodule form) under the isomorphism  $y_1 e' \mapsto (e_1 - e_2)$ ,  $e \mapsto e_1$ . Then applying [McM, Def. 4.4] we compute:

$$\begin{aligned} &[\tilde{x}^i]_{22} \cdot (y_1 e, 0, - \otimes y_1 e) \\ &= \left( y^i y_1 e, 0, - \otimes y^i y_1 e + y_1 y_2 (h_{i-1}(x_2, y) (- \otimes e)) \right), \end{aligned}$$

using:

$$\begin{aligned} &x_2^i \circ (- \otimes y_1 e) \\ &= y^i \circ (- \otimes y_1 e) + h_{i-1}(x_2, y) y_2 \circ (- \otimes y_1 e) \\ &= - \otimes y^i y_1 e + y_1 y_2 h_{i-1}(x_2, y) (- \otimes e). \end{aligned}$$

Converting back to the  $(e', e)$  data and bimodule form, we have

$$[\tilde{x}^i]_{22} \cdot (e, y_1 e, 0) = (y^i e, y^i y_1 e, h_{i-1}(x_2, y) (- \otimes e)).$$

Note that

$$(F(h_{i-1}(x, y)) \circ \eta) E : e \mapsto h_{i-1}(x_2, y) (- \otimes e).$$



- We have  $[\tilde{\varepsilon} \circ \tilde{x}^i \tilde{F}]_{22} : [\tilde{E}\tilde{F}]_{22} \rightarrow [C]_{22}$  given by:

$$\begin{pmatrix} y^i & 0 & 0 & 0 & -\varepsilon \circ h_{i-1}(x, y)F \\ h_{i-1}(x, y) \circ \eta & x^i E & Fx^i & F(\varepsilon \circ x^i y_1 F)E & \begin{matrix} -FE\varepsilon \circ F(\tau \circ h_{i-1}(x_1, x_2))F \circ \eta EF \\ -FE\varepsilon \circ F(h_{i-2}(x_1, x_2, y))F \circ \eta EF \end{matrix} \end{pmatrix}$$

using the ordered decompositions (recall Eq. 5.4):

$$\begin{aligned} - [\tilde{E}\tilde{F}]_{22} &\xrightarrow{\sim} A[y] \oplus FE[y] \oplus FE[y] \oplus FEFE[y] \oplus EF[y], \\ - [C]_{22} &\xrightarrow{\sim} A[y] \oplus FE[y]. \end{aligned}$$

Consider the first four summands first, i.e.  $G_1 G_1$ . Take an arbitrary simple tensor  $(\theta, \varphi_1) \otimes (\theta', \varphi'_1)$ , which may be viewed as an element of  $[\tilde{E}]_{21} \cdot [\tilde{F}]_{12}$ . Then:

$$[\tilde{x}^i \tilde{F}]_{22} : (\theta, \varphi_1) \otimes (\theta', \varphi'_1) \mapsto (y^i \theta, h_{i-1}(x, y) \cdot \theta + x^i \circ \varphi_1) \otimes (\theta', \varphi'_1).$$

Now applying  $[\tilde{\varepsilon}]_{22} : G_1 G_1 \rightarrow G_1$ , which is given by composition:

$$\begin{aligned} [\tilde{\varepsilon} \circ \tilde{x}^i \tilde{F}]_{22} : (\theta, \varphi_1) \otimes (\theta', \varphi'_1) &\mapsto \\ &(y^i \theta \theta', (-\theta') \circ h_{i-1}(x, y) \cdot \theta + (-\theta') \circ x^i \circ \varphi_1 \\ &+ \varphi'_1 \circ (-y^i \theta) + \varphi'_1 \circ (x^i - y^i) \cdot \theta + \varphi'_1 \circ y_1 x^i \circ \varphi_1) \\ &= (y^i \theta \theta', h_{i-1}(x, y) \cdot \theta \theta' + \varphi'_1 \circ x^i \cdot \theta + x^i \cdot \theta' \circ \varphi_1 + \varphi'_1 \circ y_1 x^i \circ \varphi_1). \end{aligned}$$

The first four columns of the matrix of  $[\tilde{\varepsilon} \circ \tilde{x}^i \tilde{F}]_{22}$  can be read off this formula.

The last column gives the restriction of  $[\tilde{\varepsilon} \circ \tilde{x}^i \tilde{F}]_{22}$  to a map  $EF[y] \rightarrow A[y] \oplus FE[y]$ . This is computed on a simple tensor  $e \otimes f$  by interpreting it in  $(G_2)_G L_2$  as  $(0, e, 0) \otimes (f, 0, 0)$ , then applying  $[\tilde{x}]_{22}$  to the first factor, and finally composing factors after interpreting in

$$\text{Hom}_{K^b(B)}(X_2, R) \otimes \text{Hom}_{K^b(B)}(R, X_2)$$

to give an element of  $G_2 \cong \text{End}_{K^b(B)}(X_2)$ .

Rewriting  $(0, e, 0)$  in the submodule form for a morphism in  $\text{Hom}_{K^b(B)}(X_2, E'X_2)$ , obtain  $(e, e, \tau y_1(- \otimes e))$ . Then:

$$[\tilde{x}^i \tilde{F}]_{22} : (e, e, \tau y_1(- \otimes e)) \otimes (f, 0, 0) \mapsto (y^i e, x^i e, x_2^i \circ \tau y_1(- \otimes e)) \otimes (f, 0, 0).$$

Now compute the composite in  $G_1$ , starting with the submodule form:

$$\begin{aligned} &(f(y_1^{-1}(y^i e - x^i e)), Ef \circ \tau \circ x_2^i \circ \tau y_1(- \otimes e)) \\ &= \left( f(-(h_{i-1}(x, y))e), Ef \circ \tau \circ x_2^i \circ \tau y_1(- \otimes e) \right) \\ &= \left( -\varepsilon(h_{i-1}(x, y)e \otimes f), Ef \circ \tau \circ x_2^i \circ \tau y_1(- \otimes e) \right). \end{aligned}$$

It remains to convert the expression to bimodule form. In this calculation we will use the facts that:

$$x_2^i \circ \tau = \tau \circ x_1^i - h_{i-1}(x_1, x_2),$$

and

$$x_2^j = y^j + y_2(h_{i-1}(x_2, y)).$$

We have:

$$\begin{aligned}
 & Ef \circ \tau \circ x_2^i \circ \tau y_1(-\otimes e) \\
 &= -Ef \circ \tau \circ (h_{i-1}(x_1, x_2))y_1(-\otimes e) \\
 &= -Ef \circ \tau y_1 \circ (h_{i-1}(x_1, x_2)(-\otimes e)) \\
 &= -Ef \circ (h_{i-1}(x_1, x_2)(-\otimes e)) - y_1 \circ Ef \circ \tau \circ (h_{i-1}(x_1, x_2)(-\otimes e)) \\
 &= -Ef \circ \left( \sum_{j+k=i-1} x_1^j (y^k + y_2 h_{k-1}(x_2, y))(-\otimes e) \right) \\
 &\quad - y_1 \circ Ef \circ \tau \circ (h_{i-1}(x_1, x_2)(-\otimes e)) \\
 &= -Ef \circ (h_{i-1}(x_1, y))(-\otimes e) - y_1 \left( Ef \circ \tau \circ h_{i-1}(x_1, x_2)(-\otimes e) \right. \\
 &\quad \left. + Ef \circ h_{i-2}(x_1, x_2, y)(-\otimes e) \right).
 \end{aligned}$$

So the second component of the bimodule form is:

$$-Ef \circ \tau \circ h_{i-1}(x_1, x_2)(-\otimes e) - Ef \circ h_{i-2}(x_1, x_2, y)(-\otimes e).$$

This can be rewritten as the application to  $e \otimes f$  of the following morphism:

$$-FE\varepsilon \circ F(\tau \circ h_{i-1}(x_1, x_2) + h_{i-2}(x_1, x_2, y))F \circ \eta EF.$$

- We have  $[\tilde{F}\tilde{x}^i \circ \tilde{\eta}]_{22} : [C]_{22} \rightarrow [\tilde{F}\tilde{E}]_{22}$  given by:

$$\begin{pmatrix}
 Fy^i \circ \eta & y^i y_1 \\
 -F(h_{i-1}(x, y)) \circ \eta & y^i \\
 0 & 0 \\
 Fx^i \circ \eta & 0 \\
 F^2(h_{i-1}(x_1, x_2) \circ \tau - h_{i-2}(x_1, x_2, y)) \circ \eta^2 & F^2(h_{i-1}(x_2, y)) \circ F\eta E
 \end{pmatrix}$$

using the ordered decompositions:

$$- [C]_{22} \xrightarrow{\sim} A[y] \oplus FE[y],$$

$$- [\tilde{F}\tilde{E}]_{22} \xrightarrow{\sim} FE[y] \oplus FE[y] \oplus FE[y] \oplus FE[y] \oplus F^2E^2[y].$$

Observe first that  $[\tilde{\eta}]_{22} : G_1 \rightarrow U$  by  $(1, 0) = \text{Id}_{G_1} \mapsto \text{Id}_R \in U$ . Now we would like to express  $[\tilde{\eta}]_{22}((\theta, \varphi_1))$  in  $(L_2)_G G_2 \cong U$  in order to compute  $[\tilde{F}\tilde{x}^i]_{22}$  by applying  $[\tilde{x}^i]_{22}$  to the factor in  $G_2$ , after which we will compose the factors again to recover an element in  $U$ . Recall (from the calculation of  $[\tilde{E}\tilde{F}]_{21}$ ) that  $\text{Id}_R \in U$  is represented in  $(L_2)_G G_2$  by:

$$\sum_a (f_a, 0, 0) \otimes (e_a, 0, 0) + \sum_b (0, f_b, 0) \otimes (0, e_b, 0) \in (L_2)_G G_2,$$

where

$$\eta(1) = \sum_a f_a \otimes e_a \in FE.$$

We also have that

$$[\tilde{\eta}]_{22}((\theta, \varphi_1)) = \text{Id}_R \cdot (\theta, \varphi_1)$$

using the right action of  $G_1$  on  $G_2$ . So we compute the right action of  $(\theta, \varphi_1)$  on our representative of  $\text{Id}_R$ , apply  $[\tilde{x}^i]_{22}$ , and then compose.

Let us first restrict attention to the case  $(\theta, \varphi_1) = (1, 0)$ , which gives the first column of the matrix. The right action does nothing. We have

$$[\tilde{x}^i]_{22} : (e_a, 0, 0) \mapsto (x^i e_a, 0, h_{i-1}(x_1, x_2)(-\otimes e_a)).$$

This is computed as follows. The map on degree 0 in the top:

$$\begin{aligned} & x_2^i \circ y_2 \tau(-\otimes(-y_1 e_a)) \\ &= y_2 x_2^i \tau(-\otimes(-y_1 e_a)) \\ &= y_2 \tau x_1^i(-\otimes(-y_1 e_a)) + y_1 y_2 h_{i-1}(x_1, x_2)(-\otimes e_a) \\ &= y_2 \tau(-\otimes(-y_1 x^i e_a)) + y_1 y_2 h_{i-1}(x_1, x_2)(-\otimes e_a). \end{aligned}$$

The formula for the first coordinate is easy to compute from the definition, and then the remainder term can be read off the last expression. By composing the maps, we obtain in  $U$ :

$$\begin{aligned} & \sum_a (x^i e_a, 0, h_{i-1}(x_1, x_2)(-\otimes e_a)) \circ (f_a, 0, 0) \\ &= (0, 0, 0, x^i, \Lambda^\circ) = (0, 0, 0, x^i, h_{i-1}(x_1, x_2) \circ \tau), \end{aligned}$$

where the last component is computed using

$$\begin{aligned} \Lambda &= (y_2 \tau(-\otimes(-y_1 x^i e_a)) + y_1 y_2 h_{i-1}(x_1, x_2)(-\otimes e_a)) \circ E f_a \circ \tau \\ &= -y_2 \tau y_1 x_1^i \tau + y_1 y_2 h_{i-1}(x_1, x_2) \tau \end{aligned}$$

together with  $\Phi_{11} = \Phi_{12} = \Phi_{21} = 0$  and the rule

$$\begin{aligned} \Lambda &= 0 - y_2 \tau y_1 \circ E \Phi_{22} \circ \tau + y_1 y_2 \Lambda^\circ \\ &= -y_2 \tau y_1 \circ E x^i \circ \tau + y_1 y_2 \Lambda^\circ. \end{aligned}$$

For the other term we have:

$$\begin{aligned} [\tilde{x}^i]_{22} : (0, e_b, 0) &\mapsto \\ &(-h_{i-1}(x_1, y) e_b, y^i e_b, -h_{i-2}(x_1, x_2, y)(-\otimes e_b)). \end{aligned}$$

The first two components are easily computed from the definition, and then the last component is read off the map on degree 0 in the top row, which is:

$$\begin{aligned} & x_2^i \circ \tau y_1(-\otimes e_b) \\ &= \tau y_1(-\otimes x^i e_b) - y_1 h_{i-1}(x_1, x_2)(-\otimes e_b) \\ &= \tau y_1(-\otimes x^i e_b) - y_1 h_{i-1}(x_1, y)(-\otimes e_b) \\ &\quad - y_1 y_2 h_{i-2}(x_1, x_2, y)(-\otimes e_b) \\ &= -\otimes x^i e_b + y_2 \tau(-\otimes x^i e_b) - -\otimes(x^i - y^i) e_b \\ &\quad - y_1 y_2 h_{i-2}(x_1, x_2, y)(-\otimes e_b) \\ &= -\otimes y^i e_b + y_2 \tau(-\otimes x^i e_b) - y_1 y_2 h_{i-2}(x_1, x_2, y)(-\otimes e_b), \end{aligned}$$

using

$$y_2 h_{i-2}(x_1, x_2, y) = h_{i-1}(x_1, x_2) - h_{i-1}(x_1, y).$$

By composing the maps, we obtain in  $U$ :

$$\begin{aligned} & \sum_b (-h_{i-1}(x_1, y)e_b, y^i e_b, -h_{i-2}(x_1, x_2, y)(- \otimes e_b)) \circ (0, f_b, 0) \\ &= (y^i, -h_{i-1}(x_1, y), 0, 0, -h_{i-2}(x_1, x_2, y)). \end{aligned}$$

The first four components are clear. The fifth is derived using

$$\begin{aligned} \Lambda &= x_2^i \circ \tau y_1(- \otimes e_b) \circ E f_b \\ &= x_2^i \tau y_1 = \tau x_1^i y_1 - y_1 h_{i-1}(x_1, x_2) \end{aligned}$$

together with the rule

$$\begin{aligned} \Lambda &= \tau y_1 \circ E(y^i) - y_2 \tau y_1 \circ E(-h_{i-1}(x_1, y)) + y_1 y_2 \Lambda^\circ \\ &= \tau y_1 y^i + y_2 \tau(y_1 h_{i-1}(x_1, y)) + y_1 y_2 \Lambda^\circ \\ &= \tau y_1 y^i + y_2 \tau(x_1^i - y^i) + y_1 y_2 \Lambda^\circ \\ &= y^i + y_2 \tau x_1^i + y_1 y_2 \Lambda^\circ \\ &= \tau x_1^i y_1 - y_1 h_{i-1}(x_1, y) + y_1 y_2 \Lambda^\circ, \end{aligned}$$

so

$$\begin{aligned} y_2 \Lambda^\circ &= -h_{i-1}(x_1, x_2) + h_{i-1}(x_1, y), \\ \Lambda^\circ &= -h_{i-2}(x_1, x_2, y). \end{aligned}$$

Taking the sum of the two terms, we conclude that  $[\tilde{F}\tilde{x}^i]_{22} : A[y] \rightarrow U$  by

$$1 \mapsto (y^i, -h_{i-1}(x_1, y), 0, x^i, h_{i-1}(x_1, x_2) \circ \tau - h_{i-2}(x_1, x_2, y)).$$

By considering the coefficients in  $FE[y]$  and  $F^2E^2[y]$  instead of in  $\text{End}(E[y])$  and  $\text{End}(E^2[y])$ , we obtain the formulas in the first column.

We turn now to the second column, the map  $FE[y] \rightarrow U$ . This is found using the same method but with  $(\theta, \varphi_1) = (0, \varphi_1)$  for any  $\varphi_1 \in FE[y]$ . We have

$$\begin{aligned} (e_a, 0, 0) \cdot (0, \varphi_1) &= (0, 0, E\varphi_1 \circ \tau(- \otimes -y_1 e_a)) \\ (0, e_b, 0) \cdot (0, \varphi_1) &= (\varphi_1(e_b), y_1 \varphi_1(e_b), E\varphi_1 \circ \tau(- \otimes e_b)), \end{aligned}$$

where we have used the calculations

$$E(y_1 \varphi_1) \circ y_2 \tau(- \otimes -y_1 e_a) = y_1 y_2 E\varphi_1 \circ \tau(- \otimes -y_1 e_a)$$

and

$$E(y_1 \varphi_1) \circ \tau y_1(- \otimes e_b) = - \otimes y_1 \varphi_1(e_b) + y_1 y_2 E\varphi_1 \circ \tau(- \otimes e_b).$$

Next we compute

$$[\tilde{x}^i]_{22} : (0, 0, E\varphi_1 \circ \tau(- \otimes -y_1 e_a)) \mapsto (0, 0, E\varphi_1 \circ x_2^i \circ \tau(- \otimes -y_1 e_a)),$$

and finally in  $U$ :

$$\begin{aligned} & \sum_a (0, 0, E\varphi_1 \circ x_2^i \circ \tau(- \otimes -y_1 e_a)) \circ (f_a, 0, 0) \\ &= (0, 0, 0, 0, \Lambda^\circ) = (0, 0, 0, 0, -E\varphi_1 \circ x_2^i \tau) \end{aligned}$$

using the calculation

$$\begin{aligned} & x_2^i \circ y_1 y_2 E \varphi_1 \circ \tau(- \otimes - y_1 e_a) \circ E f_a \circ \tau \\ &= -x_2^i \circ y_1 y_2 E \varphi_1 \circ \tau y_1 \circ \tau \\ &= -y_1 y_2 E \varphi_1 \circ x_2^i \tau. \end{aligned}$$

Then we also compute

$$\begin{aligned} [\tilde{x}^i]_{22} : (\varphi_1(e_b), y_1 \varphi_1(e_b), E \varphi_1 \circ \tau(- \otimes e_b)) &\mapsto \\ (y^i \varphi_1(e_b), y^i y_1 \varphi_1(e_b), E \varphi_1 \circ x_2^i \circ \tau(- \otimes e_b) &+ h_{i-1}(x_2, y)(- \otimes \varphi_1(e_b))) \end{aligned}$$

using the calculation

$$\begin{aligned} & x_2^i \circ (- \otimes y_1 \varphi_1(e_b) + y_1 y_2 E \varphi_1 \circ \tau(- \otimes e_b)) \\ &= - \otimes y^i y_1 \varphi_1(e_b) + y_2 h_{i-1}(x_2, y)(- \otimes y_1 \varphi_1(e_b)) \\ &\quad + y_1 y_2 E \varphi_1 \circ x_2^i \circ \tau(- \otimes e_b) \\ &= - \otimes y^i y_1 \varphi_1(e_b) + y_1 y_2 (h_{i-1}(x_2, y)(- \otimes \varphi_1(e_b)) \\ &\quad + E \varphi_1 \circ x_2^i \circ \tau(- \otimes e_b)) \\ &= - \otimes y^i y_1 \varphi_1(e_b) + y_1 y_2 (-y_1 h_{i-2}(x_1, x_2, y) E \varphi_1(- \otimes e_b) \\ &\quad + E \varphi_1 \circ \tau \circ x_1^i(- \otimes e_b)), \end{aligned}$$

and finally in  $U$ :

$$\begin{aligned} & \sum_b (y^i \varphi_1(e_b), y^i y_1 \varphi_1(e_b), E \varphi_1 \circ x_2^i \circ \tau(- \otimes e_b) \\ &\quad + h_{i-1}(x_2, y) E \varphi_1(- \otimes e_b)) \circ (0, f_b, 0) \\ &= (y^i y_1 \varphi_1, y^i \varphi_1, 0, 0, \Lambda^\circ) = (y^i y_1 \varphi_1, y^i \varphi_1, 0, 0, h_{i-1}(x_2, y) E \varphi_1 + E \varphi_1 \circ x_2^i \tau) \end{aligned}$$

using the calculation

$$\begin{aligned} \Lambda &= x_2^i \circ (- \otimes y_1 \varphi_1(e_b) + y_1 y_2 E \varphi_1 \circ \tau(- \otimes e_b)) \circ E f_b \\ &= x_2^i y_1 E \varphi_1 + y_1 y_2 E \varphi_1 \circ x_2^i \tau \\ &= y^i y_1 E \varphi_1 + y_2 h_{i-1}(x_2, y) y_1 E \varphi_1 + y_1 y_2 E \varphi_1 \circ x_2^i \tau, \end{aligned}$$

together with the rule

$$\begin{aligned} \Lambda &= \tau y_1 (y^i y_1 E \varphi_1) - y_2 \tau y_1 (y^i E \varphi_1) + y_1 y_2 \Lambda^\circ \\ &= y^i y_1 E \varphi_1 + y_1 y_2 \Lambda^\circ. \end{aligned}$$

Taking the sum of the two terms, we conclude that  $[\tilde{F} \tilde{x}^i]_{22} : FE[y] \rightarrow U$  by

$$\varphi_1 \mapsto (y^i y_1 \varphi_1, y^i \varphi_1, 0, 0, h_{i-1}(x_2, y) E \varphi_1).$$

The last component is the same as  $(F^2(h_{i-1}(x_2, y)) \circ F \eta E)(\varphi_1)$ . This gives the formulas in the second column of the matrix of  $[\tilde{F} \tilde{x}^i \circ \tilde{\eta}]_{22}$ .

**5.3. Maps  $\tilde{\rho}_\lambda$ : isomorphisms.** Now we have formulas for all of the components of the maps  $\tilde{\sigma}$ ,  $\tilde{\varepsilon} \circ \tilde{x}^i \tilde{F}$ , and  $\tilde{F} \tilde{x}^i \circ \tilde{\eta}$  that are used to construct the maps  $\tilde{\rho}_\lambda$ . It remains to make use of the isomorphisms  $\rho_\lambda$  constructed from  $\sigma$ ,  $\varepsilon \circ x^i F$ , and  $F x^i \circ \eta$ , together with these formulas, to show that  $\tilde{\rho}_\lambda$  are isomorphisms, and we accomplish that in this section.

We remind the reader of our notational convention that  $E_\lambda = E e_\lambda$  for  $e_\lambda \in A_\lambda$  the idempotents of a weight decomposition. Recall that the bimodule  $E$  satisfies  $e_j E e_i = \delta_{i+2,j} \cdot e_{i+2} E e_i$ , and similarly for  $F$  but with  $i - 2$  instead of  $i + 2$ . Finally, recall [McM, Prop. 4.25] which gives the weight idempotents for the algebra  $C$  of the 2-product.

- We have for  $[\tilde{\rho}_\lambda]_{11}$ ,  $\lambda \geq 0$ :

$$[\tilde{\rho}_\lambda]_{11} : EF_{\lambda+1}[y] \rightarrow A_{\lambda+1}[y] \oplus FE_{\lambda+1}[y] \oplus A_{\lambda+1}[y]^{\oplus \lambda}$$

by

$$[\tilde{\rho}_\lambda]_{11} = \varepsilon \oplus \sigma \oplus \bigoplus_{i=0}^{\lambda-1} \varepsilon \circ (x^i y_1) F.$$

- We have for  $[\tilde{\rho}_\lambda]_{11}$ ,  $\lambda \leq 0$ :

$$[\tilde{\rho}_\lambda]_{11} : EF_{\lambda+1}[y] \oplus A_{\lambda+1}[y]^{\oplus -\lambda} \rightarrow A_{\lambda+1}[y] \oplus FE_{\lambda+1}[y]$$

by

$$[\tilde{\rho}_\lambda]_{11} = \left( \begin{pmatrix} \varepsilon \\ \sigma \end{pmatrix}, \sum_{i=0}^{-\lambda-1} \left( F(h_{i-1}(x, y)) \circ \eta \right) \right).$$

**Proposition 5.6.** *The morphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $[\tilde{\rho}_\lambda]_{11}$  is an isomorphism for all  $\lambda$ .*

*Proof.* When  $\lambda \geq 0$  and therefore  $\lambda + 1 \geq 0$ , the map

$$\sigma \oplus \bigoplus_{i=0}^{\lambda} \varepsilon \circ x^i F : EF_{\lambda+1}[y] \xrightarrow{\sim} FE_{\lambda+1}[y] \oplus A_{\lambda+1}[y]^{\oplus \lambda+1}$$

is just  $\rho_{\lambda+1} \otimes_k k[y]$ . It is an isomorphism because  $\rho_{\lambda+1}$  is an isomorphism (by the hypothesis that  $\mathcal{V}$  is a 2-representation).

*Claim 5.7.* When  $\lambda \geq 0$ , the map

$$\sigma \oplus \varepsilon \oplus \bigoplus_{i=0}^{\lambda-1} \varepsilon \circ (x^i y_1) F : EF_{\lambda+1}[y] \rightarrow FE_{\lambda+1}[y] \oplus A_{\lambda+1}[y]^{\oplus \lambda+1}$$

is also an isomorphism.

*Proof.* Let

$$M_{-y} \in \text{End}_{A_{\lambda+1}[y]} (A_{\lambda+1}[y]^{\oplus \lambda+1})$$

be the endomorphism with matrix coefficients

$$[M_{-y}] \in \text{Mat}_{(\lambda+1) \times (\lambda+1)} (A_{\lambda+1}[y]^{\text{op}})$$

given by 1 on the diagonal and  $-y$  on the subdiagonal, 0 elsewhere. This matrix is invertible, and  $M_{-y}$  is an isomorphism. Observe that

$$\varepsilon \circ (-x^{i-1} y F) = -y \cdot \varepsilon \circ (x^{i-1} F).$$

Using this we write the map in question as a composition of isomorphisms:

$$\sigma \oplus \varepsilon \oplus \bigoplus_{i=0}^{\lambda-1} \varepsilon \circ (x^i y_1 F) = \begin{pmatrix} \text{Id}_{FE_{\lambda+1}[y]} & 0 \\ 0 & M_{-y} \end{pmatrix} \circ \left( \sigma \oplus \bigoplus_{i=0}^{\lambda} \varepsilon \circ (x^i F) \right).$$

By reordering the first two summands in the codomain, we obtain the map  $[\tilde{\rho}_\lambda]_{11}$ .  $\square$

When  $\lambda = 0$ , the two formulas for  $[\tilde{\rho}_\lambda]_{11}$  agree. Let us assume  $\lambda < 0$ , so  $\lambda + 1 \leq 0$  and the map

$$(5.5) \quad \left( \sigma, \sum_{i=0}^{-(\lambda+1)-1} Fx^i \circ \eta \right) : EF_{\lambda+1} \oplus A_{\lambda+1}[y]^{\oplus -(\lambda+1)} \xrightarrow{\sim} FE_{\lambda+1}[y]$$

is  $\rho_{\lambda+1} \otimes_k k[y]$ , an isomorphism.

*Claim 5.8.* When  $\lambda < 0$ , the map

$$\left( \sigma, \sum_{i=1}^{-\lambda-1} F(h_{i-1}(x, y)) \circ \eta \right) : EF_{\lambda+1}[y] \oplus A_{\lambda+1}[y]^{\oplus -(\lambda+1)} \rightarrow FE_{\lambda+1}[y]$$

is also an isomorphism.

*Proof.* This time we define an isomorphism

$$M_h \in \text{End}_{A_{\lambda+1}[y]} (A_{\lambda+1}[y]^{\oplus -(\lambda+1)})$$

with components  $[M_h]_{ii} = 1$ ,  $[M_h]_{ij} = y^{j-i}$  for  $j > i$ , and  $[M_h]_{ij} = 0$  for  $j < i$ . This is an upper-triangular invertible matrix:

$$\begin{pmatrix} 1 & y & y^2 & \dots & y^{-(\lambda+1)-1} \\ 0 & 1 & y & \dots & y^{-(\lambda+1)-2} \\ 0 & 0 & 1 & \dots & y^{-(\lambda+1)-3} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix}.$$

Now observe that  $F(x^i y^j) \circ \eta = Fx^i \circ \eta \cdot y^j$ . Using this, we can write:

$$\begin{aligned} \sum_{i=1}^{-\lambda-1} F(h_{i-1}(x, y)) \circ \eta &= \sum_{i=0}^{-\lambda-2} \sum_{\substack{j+k \\ =i}} (Fx^j) \circ \eta \cdot y^k \\ &= \left( \sum_{i=0}^{-(\lambda+1)-1} Fx^i \circ \eta \right) \circ M_h, \end{aligned}$$

and it follows from this and the isomorphism above the claim that the map of the claim is an isomorphism.  $\square$

Now the map

$$\left( \begin{pmatrix} \varepsilon \\ \sigma \end{pmatrix}, \sum_{i=0}^{-\lambda-1} \begin{pmatrix} y^i \\ F(h_{i-1}(x, y)) \circ \eta \end{pmatrix} \right)$$

can be written out:

$$\begin{pmatrix} \varepsilon & \text{Id} & y & \dots & y^{-\lambda-1} \\ \sigma & 0 & \eta & \dots & F(h_{-\lambda-2}(x, y)) \circ \eta \end{pmatrix}.$$

Interchanging the first two summands of the domain, we obtain the form:

$$\begin{pmatrix} \text{Id} & (\varepsilon, y, y^2, \dots, y^{-\lambda-1}) \\ 0 & \left( \sigma, \sum_{i=1}^{-\lambda-1} F(h_{i-1}(x, y)) \circ \eta \right) \end{pmatrix},$$

which (by the claim) is manifestly an isomorphism.  $\square$

- We have for  $[\tilde{\rho}_\lambda]_{21}$ ,  $\lambda \geq 0$ :

$$[\tilde{\rho}_\lambda]_{21} : F_{\lambda+1}[y] \oplus FEF_{\lambda+1}[y] \rightarrow F_{\lambda+1}[y] \oplus F_{\lambda+1}[y] \oplus F^2E_{\lambda+1}[y] \oplus F_{\lambda+1}[y]^{\oplus \lambda}$$

by

$$[\tilde{\rho}_\lambda]_{21} = \begin{pmatrix} 1 & 0 \\ 0 & F\varepsilon \\ 0 & F\sigma \\ \bigoplus_{i=0}^{\lambda-1} x^i & \bigoplus_{i=0}^{\lambda-1} F(\varepsilon \circ (x^i y_1)F) \end{pmatrix}.$$

- We have for  $[\tilde{\rho}_\lambda]_{21}$ ,  $\lambda \leq 0$ :

$$[\tilde{\rho}_\lambda]_{21} : F_{\lambda+1}[y] \oplus FEF_{\lambda+1}[y] \oplus F_{\lambda+1}[y]^{\oplus -\lambda} \rightarrow F_{\lambda+1}[y] \oplus F_{\lambda+1}[y] \oplus F^2E_{\lambda+1}[y]$$

by

$$[\tilde{\rho}_\lambda]_{21} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & F\varepsilon & \sum_{i=0}^{-\lambda-1} y^i \\ 0 & F\sigma & \sum_{i=0}^{-\lambda-1} F(F(h_{i-1}(x, y)) \circ \eta) \end{pmatrix}.$$

**Proposition 5.9.** *The morphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $[\tilde{\rho}_\lambda]_{21}$  is an isomorphism for all  $\lambda$ .*

*Proof.* When  $\lambda \geq 0$ , we have that

$$\begin{aligned} F\varepsilon \oplus F\sigma \oplus \bigoplus_{i=0}^{\lambda-1} F(\varepsilon \circ (x^i y_1)F) & : FEF_{\lambda+1}[y] \\ & \rightarrow F_{\lambda+1}[y] \oplus F^2E_{\lambda+1}[y] \oplus F_{\lambda+1}[y]^{\oplus \lambda} \end{aligned}$$

is an isomorphism, using Claim 5.7 and the fact that horizontal composition of the identity functor on  $F$  with an isomorphism gives an isomorphism. Then  $[\tilde{\rho}_\lambda]_{21}$  may be compressed to a lower-triangular  $2 \times 2$  matrix with an isomorphism in position  $(2, 2)$ , so it is an isomorphism.

When  $\lambda = 0$ , the two formulas for  $[\tilde{\rho}_\lambda]_{21}$  agree. Assume now that  $\lambda < 0$ , so the map

$$\left( F\sigma, \sum_{i=1}^{-\lambda-1} F(F(h_{i-1}(x, y)) \circ \eta) \right) : FEF_{\lambda+1}[y] \oplus F_{\lambda+1}[y]^{\oplus -(\lambda+1)} \rightarrow F^2E_{\lambda+1}[y]$$



is an isomorphism using Claim 5.8. Now expand the notation of the map  $[\tilde{\rho}_\lambda]_{21}$  in the third row:

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & F\varepsilon & 1 & \sum_{i=1}^{-\lambda-1} y^i \\ 0 & F\sigma & 0 & \sum_{i=1}^{-\lambda-1} F(F(h_{i-1}(x, y)) \circ \eta) \end{pmatrix}.$$

After switching the second and third summands of the domain, we obtain an upper-triangular matrix with isomorphisms on the diagonal, so  $[\tilde{\rho}_\lambda]_{21}$  is an isomorphism.  $\square$

- We have for  $[\tilde{\rho}_\lambda]_{12}$ ,  $\lambda \geq 0$ :

$$[\tilde{\rho}_\lambda]_{12} : E_{\lambda-1}[y] \oplus EFE_{\lambda-1}[y] \rightarrow E_{\lambda-1}[y] \oplus E_{\lambda-1}[y] \oplus FE_{\lambda-1}^2[y] \oplus E_{\lambda-1}[y]^{\oplus \lambda}$$

by

$$[\tilde{\rho}_\lambda]_{12} = \begin{pmatrix} 0 & \varepsilon E \\ 1 & y_1 \circ \varepsilon E \\ 0 & \sigma E \\ \bigoplus_{i=0}^{\lambda-1} x^i & \bigoplus_{i=0}^{\lambda-1} (\varepsilon \circ (x^i y_1) F) E \end{pmatrix}.$$

- We have for  $[\tilde{\rho}_\lambda]_{12}$ ,  $\lambda \leq 0$ :

$$[\tilde{\rho}_\lambda]_{12} : E_{\lambda-1}[y] \oplus EFE_{\lambda-1}[y] \oplus E_{\lambda-1}[y]^{\oplus -\lambda} \rightarrow E_{\lambda-1}[y] \oplus E_{\lambda-1}[y] \oplus FE_{\lambda-1}^2[y]$$

by

$$[\tilde{\rho}_\lambda]_{12} = \begin{pmatrix} 0 & \varepsilon E & \sum_{i=0}^{-\lambda-1} y^i \\ 1 & y_1 \circ \varepsilon E & \sum_{i=0}^{-\lambda-1} y^i y_1 \\ 0 & \sigma E & \sum_{i=0}^{-\lambda-1} (F(h_{i-1}(x, y)) \circ \eta) E \end{pmatrix}.$$

**Proposition 5.10.** *The morphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $[\tilde{\rho}_\lambda]_{12}$  is an isomorphism for all  $\lambda$ .*

*Proof.* When  $\lambda \geq 0$ , we have that

$$\begin{aligned} \varepsilon E \oplus \sigma E \oplus \bigoplus_{i=0}^{\lambda-1} (\varepsilon \circ (x^i y_1) F) E : EFE_{\lambda-1}[y] \\ \rightarrow E_{\lambda-1}[y] \oplus FE_{\lambda-1}^2[y] \oplus E_{\lambda-1}[y]^{\oplus \lambda} \end{aligned}$$

is an isomorphism, using Claim 5.7 with  $E$  applied on the right. Note that  $E$  applied on the right here is the same as  ${}_{\lambda+1}E_{\lambda-1}$  applied on the right, and this raises the weight by 2, so we still invoke the isomorphism  $\rho_{\lambda+1}$  for weight  $\lambda + 1$ .

We perform some row operations on the matrix of  $[\tilde{\rho}_\lambda]_{12}$ . Subtract  $y_1$  times the first row from the second to eliminate the coefficient  $y_1 \circ \varepsilon E$ . Then exchange the first and second rows, then exchange the second and

third rows, then collapse the second and third into the notation of the fourth. We obtain:

$$\begin{pmatrix} 1 & & & 0 \\ 0 \oplus 0 \oplus \bigoplus_{i=0}^{\lambda-1} x^i & \sigma E \oplus \varepsilon E \oplus \bigoplus_{i=0}^{\lambda-1} (\varepsilon \circ (x^i y_1) F) E \end{pmatrix},$$

which is upper-triangular with isomorphisms on the diagonal, so the original matrix for  $[\tilde{\rho}_\lambda]_{12}$  is an isomorphism.

When  $\lambda = 0$ , the two formulas for  $[\tilde{\rho}_\lambda]_{12}$  agree. Assume now that  $\lambda < 0$ , so the map

$$\left( \sigma E, \sum_{i=1}^{-\lambda-1} (F(h_{i-1}(x, y)) \circ \eta) E \right) : EFE_{\lambda-1}[y] \oplus E_{\lambda-1}[y]^{\oplus -(\lambda+1)} \rightarrow FE_{\lambda-1}^2[y]$$

is an isomorphism using Claim 5.8. Now expand the notation of the map  $[\tilde{\rho}_\lambda]_{12}$  in the third row:

$$[\tilde{\rho}_\lambda]_{12} = \begin{pmatrix} 0 & \varepsilon E & 1 & \sum_{i=1}^{-\lambda-1} y^i \\ 1 & y_1 \circ \varepsilon E & y_1 & \sum_{i=1}^{-\lambda-1} y^i y_1 \\ 0 & \sigma E & 0 & \sum_{i=1}^{-\lambda-1} (F(h_{i-1}(x, y)) \circ \eta) E \end{pmatrix}.$$

Exchange the first and second rows, then the first and third columns, then collapse the third and fourth columns into the notation of the third, and obtain:

$$\begin{pmatrix} 1 & y_1 & \left( y_1 \circ \varepsilon E, \sum_{i=1}^{-\lambda-1} y^i y_1 \right) \\ 0 & 1 & \left( \varepsilon E, \sum_{i=1}^{-\lambda-1} y^i \right) \\ 0 & 0 & \left( \sigma E, \sum_{i=1}^{-\lambda-1} (F(h_{i-1}(x, y)) \circ \eta) E \right) \end{pmatrix}.$$

Since this is upper-triangular with isomorphisms on the diagonal, the original matrix  $[\tilde{\rho}_\lambda]_{12}$  is an isomorphism.  $\square$

- We have for  $[\tilde{\rho}_\lambda]_{22}$ ,  $\lambda \geq 0$ :

$$(5.6) \quad \begin{aligned} [\tilde{\rho}_\lambda]_{22} : A_{\lambda-1}[y] \oplus FE_{\lambda-1}[y]^{\oplus 2} \oplus FEFE_{\lambda-1}[y] \oplus EF_{\lambda-1}[y] \\ \rightarrow FE_{\lambda-1}[y]^{\oplus 4} \oplus F^2 E_{\lambda-1}^2[y] \oplus A_{\lambda-1}[y]^{\oplus \lambda} \oplus FE_{\lambda-1}[y]^{\oplus \lambda} \end{aligned}$$

by  $[\tilde{\rho}_\lambda]_{22} =$

$$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & F\varepsilon E & 0 \\ \eta & y_1 & 0 & 0 & \sigma \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & F\sigma E & 0 \\ \bigoplus_{i=0}^{\lambda-1} y^i & 0 & 0 & 0 & \bigoplus_{i=0}^{\lambda-1} -\varepsilon \circ (h_{i-1}(x, y))F \\ \bigoplus_{i=0}^{\lambda-1} h_{i-1}(x, y) \circ \eta & \bigoplus_{i=0}^{\lambda-1} x^i E & \bigoplus_{i=0}^{\lambda-1} Fx^i & \bigoplus_{i=0}^{\lambda-1} F(\varepsilon \circ x^i y_1 F)E & \Theta \end{pmatrix},$$

where

$$\Theta = \bigoplus_{i=0}^{\lambda-1} -FE\varepsilon \circ F(\tau \circ h_{i-1}(x_1, x_2) - h_{i-2}(x_1, x_2, y))F \circ \eta EF.$$

- We have for  $[\tilde{\rho}_\lambda]_{22}$ ,  $\lambda \leq 0$ :

$$(5.7) \quad [\tilde{\rho}_\lambda]_{22} : A_{\lambda-1}[y] \oplus FE_{\lambda-1}[y]^{\oplus 2} \oplus FEF E_{\lambda-1}[y] \oplus EF_{\lambda-1}[y] \\ \oplus A_{\lambda-1}[y]^{\oplus -\lambda} \oplus FE_{\lambda-1}[y]^{\oplus -\lambda} \rightarrow FE_{\lambda-1}[y]^{\oplus 4} \oplus F^2 E_{\lambda-1}^2[y]$$

by  $[\tilde{\rho}_\lambda]_{22} =$

$$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 & \sum_{i=0}^{-\lambda-1} Fy^i \circ \eta & \sum_{i=0}^{-\lambda-1} y^i y_1 \\ 0 & 0 & 0 & F\varepsilon E & 0 & \sum_{i=0}^{-\lambda-1} -F(h_{i-1}(x, y)) \circ \eta & \sum_{i=0}^{-\lambda-1} y^i \\ \eta & y_1 & 0 & 0 & \sigma & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & \sum_{i=0}^{-\lambda-1} Fx^i \circ \eta & 0 \\ 0 & 0 & 0 & F\sigma E & 0 & \Theta' & \sum_{i=0}^{-\lambda-1} F^2(h_{i-1}(x_2, y)) \circ F\eta E \end{pmatrix},$$

where

$$\Theta' = \sum_{i=0}^{-\lambda-1} \left( F^2(h_{i-1}(x_1, x_2) \circ \tau - h_{i-2}(x_1, x_2, y)) \circ \eta^2 \right).$$

**Proposition 5.11.** *The morphism of  $(A^{\text{op}}[y], A^{\text{op}}[y])$ -bimodules  $[\tilde{\rho}_\lambda]_{22}$  is an isomorphism for all  $\lambda$ .*

*Proof.* When  $\lambda > 0$  and therefore  $\lambda - 1 \geq 0$ , the map

$$\sigma \oplus \bigoplus_{i=0}^{\lambda-2} -\varepsilon \circ x^i F : EF_{\lambda-1}[y] \rightarrow FE_{\lambda-1}[y] \oplus A_{\lambda-1}[y]^{\oplus \lambda-1}$$

is an isomorphism. (The minus sign does not interfere.)

*Claim 5.12.* When  $\lambda > 0$ , the map

$$\sigma \oplus \bigoplus_{i=1}^{\lambda-1} -\varepsilon \circ (h_{i-1}(x, y))F : EF_{\lambda-1}[y] \rightarrow FE_{\lambda-1}[y] \oplus A_{\lambda-1}[y]^{\oplus \lambda-1}$$

is an isomorphism.

*Proof.* Define an isomorphism

$$M'_h \in \text{End}_{A_{\lambda-1}[y]} (A_{\lambda-1}[y]^{\oplus \lambda-1})$$

with components  $[M'_h]_{ii} = 1$ ,  $[M'_h]_{ij} = y^{i-j}$  for  $i > j$ , and  $[M'_h]_{ij} = 0$  for  $i < j$ . This is a lower-triangular invertible matrix:

$$\begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ y & 1 & 0 & \dots & 0 \\ y^2 & y & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ y^{\lambda-2} & y^{-\lambda-3} & y^{\lambda-4} & \dots & 1 \end{pmatrix}.$$

Now observe that  $\varepsilon \circ (x^i y^j)F = y^j \cdot \varepsilon \circ x^i F$ . Using this, we can write:

$$\begin{aligned} \bigoplus_{i=1}^{\lambda-1} -\varepsilon \circ (h_{i-1}(x, y))F &= \bigoplus_{i=0}^{\lambda-2} \sum_{\substack{j+k \\ =i}} y^k \cdot (-\varepsilon \circ x^j F) \\ &= M'_h \circ \left( \bigoplus_{i=0}^{\lambda-2} -\varepsilon \circ x^i F \right), \end{aligned}$$

and it follows from this and the isomorphism above the claim that the map of the claim is an isomorphism.  $\square$

Now assume  $\lambda > 0$  and reorder the summands of the domain and codomain to permute the rows and columns of the matrix of  $[\tilde{\rho}_\lambda]_{22}$ . Let the domain be given in the order:

$$FE_{\lambda-1}[y]^{\oplus 2} \oplus A_{\lambda-1}[y] \oplus EF_{\lambda-1}[y] \oplus FEF_{\lambda-1}[y],$$

where the first two identical summands appear in the same order as before, and the others are unique. Let the codomain be given in the order:

$$\begin{aligned} FE_{\lambda-1}[y]^{\oplus 2} \oplus A_{\lambda-1}[y] \oplus FE_{\lambda-1}[y] \oplus A_{\lambda-1}[y]^{\oplus \lambda-1} \\ \oplus F^2 E_{\lambda-1}^2[y] \oplus FE_{\lambda-1}[y] \oplus FE_{\lambda-1}[y]^{\oplus \lambda-1}, \end{aligned}$$

where the new summand number and corresponding old summand number are given more specifically by the following chart:

$$\begin{array}{cccccccccccccccc} \text{new:} & 1 & 2 & 3 & 4 & 5 & 6 & 7 & \dots & \lambda+3 & \lambda+4 & \lambda+5 & \lambda+6 & \dots & 2\lambda+5 \\ \text{old:} & 4 & 1 & 6 & 3 & 7 & 8 & 9 & \dots & \lambda+5 & 2 & 5 & \lambda+6 & \dots & 2\lambda+5. \end{array}$$

Writing the matrix of  $[\tilde{\rho}_\lambda]_{22}$ , for  $\lambda > 0$ , with columns and rows changed by the above permutations, we obtain:

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ y_1 & 0 & \eta & \sigma & 0 & 0 \\ 0 & 0 & \bigoplus_{i=1}^{\lambda-1} y^i & \bigoplus_{i=1}^{\lambda-1} -\varepsilon \circ (h_{i-1}(x, y))F & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & F\varepsilon E \\ 0 & 0 & 0 & 0 & 0 & F\sigma E \\ \bigoplus_{i=0}^{\lambda-1} x^i E & \bigoplus_{i=0}^{\lambda-1} Fx^i & \bigoplus_{i=0}^{\lambda-1} h_{i-1}(x, y) \circ \eta & \Theta & \bigoplus_{i=0}^{\lambda-1} F(\varepsilon \circ x^i y_1 F)E \end{pmatrix}.$$

After compressing the notation of rows 3 and 4 of this matrix, and also of rows 6-8, we obtain a lower-triangular matrix. The last two diagonal entries are

$$\begin{pmatrix} & \sigma \\ \bigoplus_{i=1}^{\lambda-1} -\varepsilon \circ (h_{i-1}(x, y))F & \end{pmatrix},$$

which is an isomorphism by the claim, and

$$\begin{pmatrix} F\varepsilon E \\ F\sigma E \\ \bigoplus_{i=0}^{\lambda-1} F(\varepsilon \circ x^i y_1 F)E \end{pmatrix} : FEF E_{\lambda-1}[y] \rightarrow FE_{\lambda-1}[y] \oplus F^2 E_{\lambda-1}^2[y] \oplus FE_{\lambda-1}[y]^{\oplus \lambda},$$

which is an isomorphism for  $\lambda > 0$ , and therefore  $\lambda + 1 \geq 0$ , using Claim 5.7 with  $F$  applied on the left,  $E$  on the right.

When  $\lambda = 0$  the matrix of  $[\tilde{\rho}_\lambda]_{22}$  is given by removing rows 3,  $5-(\lambda + 3)$ , and  $(\lambda + 6)-(2\lambda + 5)$ :

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ y_1 & 0 & \eta & \sigma & 0 \\ 0 & 0 & 0 & 0 & F\varepsilon E \\ 0 & 0 & 0 & 0 & F\sigma E \end{pmatrix}.$$

When  $\lambda = 0$  we also have isomorphisms:

$$(\eta, \sigma) : EF_{\lambda-1}[y] \oplus A[y]_{\lambda-1} \xrightarrow{\sim} FE_{\lambda-1}[y]$$

and

$$\begin{pmatrix} F\varepsilon E \\ F\sigma E \end{pmatrix} : F(FE_{\lambda+1})E[y] \xrightarrow{\sim} F_{\lambda+1}E[y] \oplus F(EF_{\lambda+1})E[y],$$

so we see that again the matrix can be written as a lower-triangular matrix with invertible diagonal entries.

Finally, assume  $\lambda < 0$ . We have an isomorphism:

$$\left( \sigma, \sum_{i=0}^{-\lambda} Fx^i \circ \eta \right) : EF_{\lambda-1}[y] \oplus A_{\lambda-1}[y]^{\oplus -(\lambda-1)} \xrightarrow{\sim} FE_{\lambda-1}[y],$$

which is the isomorphism  $\rho_{\lambda-1} \otimes_k k[y]$ . There is a final claim to check:

*Claim 5.13.* When  $\lambda < 0$ , the map

$$\left( \sigma, \eta, \sum_{i=0}^{-\lambda-1} -F(x^i y_1) \circ \eta \right) : EF_{\lambda-1}[y] \oplus A_{\lambda-1}[y]^{\oplus -(\lambda-1)} \rightarrow FE_{\lambda-1}[y]$$

is an isomorphism.

*Proof.* Define an isomorphism

$$M'_{-y} \in \text{End}_{A_{\lambda-1}[y]} (A_{\lambda-1}[y]^{\oplus \lambda-1})$$

with components  $[M_h]_{ij}$  given by 1 along the diagonal and  $-y$  along the subdiagonal. This is a lower-triangular invertible matrix. We write the map in question as a composition of isomorphisms:

$$\begin{aligned} & \left( \sigma, \eta, \sum_{i=0}^{-\lambda-1} -F(x^i y_1) \circ \eta \right) = \left( \sigma, \eta, \sum_{i=1}^{-\lambda} Fx^i \circ \eta \right) \\ & \circ \begin{pmatrix} \text{Id}_{EF_{\lambda-1}[y]} & 0 & 0 \\ 0 & \text{Id}_{A_{\lambda-1}[y]} & 0 \\ 0 & 0 & -\text{Id}_{A_{\lambda-1}[y]^{\oplus-\lambda}} \end{pmatrix} \circ \begin{pmatrix} \text{Id}_{EF_{\lambda-1}[y]} & 0 \\ 0 & M'_{-y} \end{pmatrix}. \end{aligned}$$

□

Now let  $W$  be the endomorphism of the codomain of  $[\tilde{\rho}_\lambda]_{22}$  given by the invertible matrix:

$$[W] = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -y_1 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

We show that  $[W] \cdot [\tilde{\rho}_\lambda]_{22}$  is equivalent to a lower-triangular matrix after giving a suitable permutation of the domain and codomain summands. Let the domain be given in the order:

$$\begin{aligned} & EF_{\lambda-1}[y] \oplus A_{\lambda-1}[y]^{-(\lambda-1)} \oplus FE_{\lambda-1}[y] \\ & \oplus FEF_{\lambda-1}[y] \oplus FE_{\lambda-1}[y]^{-(\lambda+1)} \oplus FE_{\lambda-1}[y]^{\oplus 2}, \end{aligned}$$

where the change of summand numbers is given by the following chart:

new:	1	2	3	4	...	$-\lambda + 2$	$-\lambda + 3$	$-\lambda + 4$
old:	5	1	6	7	...	$-\lambda + 5$	2	4
new:	$-\lambda + 5$	$-\lambda + 6$	...	$-2\lambda + 4$	$-2\lambda + 5$	$-2\lambda + 6$		
old:	$-\lambda + 7$	$-\lambda + 8$	...	$-2\lambda + 5$	$-\lambda + 6$	3.		

Let the codomain be given in the order:

$$FE_{\lambda-1}[y]^{\oplus 4} \oplus F^2 E_{\lambda-1}^2[y],$$

where the change of summand numbers is given by the following chart:

new:	1	2	3	4	5
old:	3	4	5	2	1.

The matrix of  $[W] \cdot [\tilde{\rho}_\lambda]_{22}$  for  $\lambda < 0$  agrees with that for  $[\tilde{\rho}_\lambda]_{22}$  except in the third row, where it is:

$$\left( \eta \ 0 \ 0 \ 0 \ \sigma \ \sum_{i=0}^{-\lambda-1} -F(x^i y_1) \circ \eta \ 0 \ 0 \right).$$

Writing now the matrix of  $[W] \cdot [\tilde{\rho}_\lambda]_{22}$  with columns and rows changed by the above permutations, and compressing some columns, we obtain:

$$\begin{pmatrix} \left( \sigma, \eta, \sum_{i=0}^{-\lambda-1} -F(x^i y_1) \circ \eta \right) & 0 & (0, 0) & 0 & 0 \\ \left( 0, 0, \sum_{i=0}^{-\lambda-1} Fx^i \circ \eta \right) & 1 & (0, 0) & 0 & 0 \\ (0, 0, \Theta') & 0 & \left( F\sigma E, \sum_{i=1}^{-\lambda-1} F^2(h_{i-1}(x_2, y)) \circ F\eta E \right) & 0 & 0 \\ \left( 0, 0, \sum_{i=0}^{-\lambda-1} -F(h_{i-1}(x, y)) \circ \eta \right) & 0 & \left( F\varepsilon E, \sum_{i=1}^{-\lambda-1} y^i \right) & 1 & 0 \\ \left( 0, 0, \sum_{i=0}^{-\lambda-1} Fy^i \circ \eta \right) & 0 & \left( 0, \sum_{i=1}^{-\lambda-1} y^i y_1 \right) & y_1 & 1 \end{pmatrix}.$$

The upper left map is an isomorphism by the Claim proved above. The middle diagonal map is an isomorphism because it is the isomorphism of Claim 5.8 with  $F$  applied on the left and  $E$  on the right. So the matrix is lower-triangular with isomorphisms along the diagonal.  $\square$

## REFERENCES

- [CF] Louis Crane and Igor Frenkel, *Four-dimensional topological quantum field theory, Hopf categories, and the canonical bases*, no. 10, 5136–5154.
- [CR] Joseph Chuang and Raphaël Rouquier, *Derived equivalences for symmetric groups and  $\mathfrak{sl}_2$ -categorification*, no. 1, 245–298.
- [KLa] Mikhail Khovanov and Aaron D. Lauda, *A diagrammatic approach to categorification of quantum groups I*, no. 14, 309–347.
- [KLb] ———, *A diagrammatic approach to categorification of quantum groups II*, no. 5, 2685–2700.
- [Lau] Aaron D. Lauda, *A categorification of quantum  $\mathfrak{sl}(2)$* , no. 6, 3327–3424.
- [McM] Matthew McMillan, *A tensor 2-product of 2-representations of  $\mathfrak{sl}(2)^+$* .
- [MR] Andrew Manion and Raphaël Rouquier, *Higher representations and cornered Heegaard Floer homology*.
- [Roua] Raphaël Rouquier, *2-Kac-Moody algebras*.
- [Roub] ———, *Hopf categories*.
- [RT] Nikolai Reshetikhin and Vladimir Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, no. 3, 547–597.
- [Ver] Laurent Vera, *Faithfulness of simple 2-representations of  $\mathfrak{sl}_2$* .
- [Wit] Edward Witten, *Quantum field theory and the Jones polynomial*, no. 3, 351–399.