

The general definition of Kähler angle

Yongpin Zhu¹

Abstract

In this paper, we give a general definition of Kähler angle. There are many results about Kähler angle one can try to generalize to the general case.

1 Introduction

It is well-known that the algebraic structure of a vector space can be generalized to the corresponding objects on the manifold by a pointwise way. The geometric version of the algebraic property of Wirtinger's inequality was given, such as, in [2]. In this paper, we give the general definition of the Kähler angle by the geometric version of Wirtinger's inequality.

Suppose that (N, J, \langle, \rangle) is a Hermite manifold, where \langle, \rangle is the Riemannian metric on N , J is a complex structure compatible with \langle, \rangle , i.e.

$$\langle JU, JV \rangle = \langle U, V \rangle.$$

The Kähler form ω on N is defined by

$$\omega(U, V) = \langle JU, V \rangle.$$

2 The general definition of Kähler angle

For the convenience of discussion, we first state an algebraic version of Wirtinger's inequality which takes from [3], the proof is almost the same as the geometric version.

Theorem 2.1. *Let (V, J, \langle, \rangle) be a euclidian vector space endowed with a compatible almost complex structure J , and the associated fundamental form ω . Let $W \subset V$ be an oriented subspace of dimension $2m$. The induced scalar product on W together with the chosen orientation define a natural volume form $\text{vol}_W \in \bigwedge^{2m} W^*$. Then*

$$\omega^m|_W \leq m! \text{Vol}_W,$$

and that equality holds if and only if $W \subset V$ is a complex subspace, i.e. $J(W) = W$, and the orientation is the one induced by the almost complex structure.

The geometric version of Wirtinger's inequality was given by a pointwise way.

¹ School of Mathematical Sciences, University of Science and Technology of China, Hefei 230026, P.R. of China

Theorem 2.2 (Wirtinger inequation). *Let N be a n -dimensional Hermite manifold, let $M \subset N$ is any $2m$ -dimensional orientable real submanifold. At any point $p \in M$, let dV_p denote the volume form of the induced metric on M . Then the restriction of the m -power $\omega^m = \omega \wedge \cdots \wedge \omega$ of the Kähler form of N to T_pM satisfies*

$$\frac{\omega^m}{m!} \leq dV_p,$$

and equality holds if and only if T_pM is a complex subspace of T_pN , with the canonical orientation.

Proof. If $m = 1$, then for any unit vectors $v, w \in T_pN$,

$$\omega(w, v)^2 = \langle Jv, w \rangle^2 \leq |Jv|^2 |w|^2 = |v|^2 |w|^2 = 1.$$

and that equality holds $\iff w = Jv$, i.e. $\text{span}\{w, v\}$ is the one-dimensional complex subspace of T_pN . For the general case, we now consider the form ω restricted to T_pM , if we choose the oriented orthonormal basis $\{e_1, \dots, e_{2m}\}$ of T_pM , then $\omega(e_i, e_j) = \langle Je_i, e_j \rangle = J_i^j$. By $J^2 = -id$ and the elementary linear algebra, we know that there exists the oriented orthonormal basis, which still denote by $\{e_1, \dots, e_{2m}\}$, such that J can be represented by

$$\begin{pmatrix} 0 & \lambda_1 & & & & \\ -\lambda_1 & 0 & & & & \\ & & \ddots & & & \\ & & & 0 & \lambda_m & \\ & & & -\lambda_m & 0 & \end{pmatrix},$$

Then we have $\omega(e_{2k-1}, e_{2k}) = \lambda_k, k = 1, 2, \dots, m$. Let $\{\theta^i\}$ be the dual basis of $\{e_i\}$, then

$$\omega = \sum_{k=1}^m \lambda_k \theta^{2k-1} \wedge \theta^{2k},$$

and $|\lambda_k| \leq 1$. Thus, we have

$$\omega^m = m! \lambda_1 \cdots \lambda_m \theta^1 \wedge \cdots \wedge \theta^{2m} = m! \lambda_1 \cdots \lambda_m dV_p,$$

so $|\omega^m| \leq m! dV_p$, and equality holds $\iff |\lambda_1 \cdots \lambda_m| = 1 \iff \omega(e_{2k-1}, e_{2k}) = \pm 1, \forall k = 1, \dots, m \iff e_{2k-1} = \pm J e_{2k}, 1 \leq k \leq m \iff T_pM$ is a complex subspace of T_pN . Equality holds without the absolute value sign exactly when the orientation agree with the one induced by J . \square

Note that the statement of the theorem is pointwise, so it still holds true for the almost-Hermite manifolds, in fact, it is a geometric version of the algebraic version of Wirtinger's inequality. As the top form over M , note that the dimension of the top form space is 1, we can introduce the function

$$\frac{\omega^m|_{T_pM}}{m! dV_p} \leq 1$$

on M to characterize the difference between the two top form. When $n = 2, m = 1$, the function is $\cos \alpha$ in [1], which was first introduced in [1]. And then

α was called the Kähler angle because of its geometric meaning in that case: $\cos \alpha = \omega(e_1, e_2) = \langle Je_1, e_2 \rangle$, i.e. the angle between the unit vector Je_1 and e_2 , where e_1, e_2 are local orthonormal frame fields on M . Therefore, we still denote the function by $\cos \alpha$ and call α the Kähler angle.

Definition 2.3 (Kähler angle). *Let N be a n -dimensional almost-Hermite manifold, let $M \subset N$ is any $2m$ -dimensional orientable real submanifold. Then we define Kähler angle α by*

$$\cos \alpha = \frac{\omega^m|_{TM}}{m!dV}, \quad (*)$$

where ω is the Kähler form of N , dV denote the volume form of the induced metric on M .

Next, we consider the case that $\cos \alpha \equiv 1$. First, we recall the well-known Newlander – Nirenberg theorem.

Theorem 2.4 (Newlander-Nirenberg). *Let (M, J) be an almost-Hermite manifold, if J is integrabel, i.e.the torsion tensor of J is 0, then J is a complex structure induced by a complex manifold structure of M .*

When $\cos \alpha \equiv 1$, by the Wirtinger inequality, for any $p \in M$, T_pM is a complex subspace of T_pN , i.e. $J_p(T_pM) \subset T_pM$, thus $(M, J|_M, g)$ is a almost-Hermite submanifold, where g is an induced metric of M . If N is a Hermite manifold, then $(M, J|_M, g)$ is a Hermite submanifold. In fact, by the definition of the torsion tensor of the almost complex structure, the torsion tensor of $J|_M$ is the limit of the torsion tensor of the complex structure J on M , so it is 0. Then it is known from the Newlander – Nirenberg theorem that it is a complex submanifold. We call the point of $\alpha = 0$ (i.e. $\cos \alpha = 1$) a complex point, and the point of $\alpha = \pi$ (i.e. $\cos \alpha = -1$) a anti-complex point.

By the definition, Kähler angles $\alpha \in [0, \pi]$ are determined by (*) and the function arccos, thus the function α is continuous on M and smooth except for the complex points and the anti-complex points. When $m = 1$, and we consider it locally, we can always allow the angle of the two vector fields to take a value outside $[0, \pi]$, and when we calculate the derivative, the two angle differ by at most one negative sign, so that $|\nabla \alpha|$ is consistent as a function on M . And when we consider the subsequent question, we always get the derivative of angle α by the derivative of the function $\cos \alpha$, so we can also directly consider the function (*). But at least, when $m = 1$, expressing the function (*) as $\cos \alpha$ can effectively simplify the expression by using the properties of $\cos \alpha$ (e.g., $|\cos(\alpha)| \leq 1$, $\cos \alpha^2 + \sin \alpha^2 = 1$, etc.).

In the general dimensionality, after a choice of the appropriate local unitary frames, the function(*) = $\cos \alpha_1 \cdots \cos \alpha_m$, where $\cos \alpha_k = \omega(e_{2k-1}, e_{2k}), k = 1, \cdots, m$. Thus the α_k can be treated similarly, note that α_k depend on the choice of the local unitary frames, but not for Kähler angle α , so we study the function (*) directly. For convenience, we call the function (*) is the Kähler function.

In this paper, we only focus on the definition of Kähler angle. But after giving the general definition of Kähler angle, the original results about Kähler angle can be generalized from N is a Kähler manifold, and $m = 1, n = 2$, to the general case: N is a almost-Hermite manifold, and any dimension $2m$ and any codimension $2(n - m)$.

References

- [1] S. S. Chern and J. Wolfson, Minimal surfaces by moving frames, Amer. J. Math. 105(1983), 59-83.
- [2] Lawson, H.B. Lectures on Minimal Submanifolds. 1. University of California, 1980.
- [3] D.Huybrechts, Complex geometry. An introduction [M]. J.H.w. Dietz nachf. g.m.b.h, 2005.