

Moduli stabilization, F-term uplifting and soft supersymmetry breaking terms

Hiroyuki Abe^{1,*}, Tetsutaro Higaki^{2,†}, Tatsuo Kobayashi^{3,‡} and Yuji Omura^{4,§}

¹*Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*

²*Department of Physics, Tohoku University, Sendai 980-8578, Japan*

³*Department of Physics, Kyoto University, Kyoto 606-8502, Japan*

³*Department of Physics, Kyoto University, Kyoto 606-8501, Japan*

Abstract

We study moduli stabilization with F-term uplifting. As a source of uplifting F-term, we consider spontaneous supersymmetry breaking models, e.g. the Polonyi model and the Intriligator-Seiberg-Shih model. We analyze potential minima by requiring almost vanishing vacuum energy and evaluate the size of modulus F-term. We also study soft SUSY breaking terms. In our scenario, the mirage mediation is dominant in gaugino masses. Scalar masses can be comparable with gaugino masses or much heavier, depending on couplings with spontaneous supersymmetry breaking sector.

*E-mail address: abe@yukawa.kyoto-u.ac.jp

†E-mail address: tetsu@tuhep.phys.tohoku.ac.jp

‡E-mail address: kobayash@gauge.scphys.kyoto-u.ac.jp

§E-mail address: omura@scphys.kyoto-u.ac.jp

1 Introduction

In superstring theory, moduli stabilization is one of important issues to study, because their vacuum expectation values (VEVs) determine several types of couplings in 4D effective field theory of massless modes and through moduli stabilization supersymmetry (SUSY) may break.

Non-perturbative superpotential, e.g. the gaugino condensation, is important to stabilize them. Even if non-perturbative superpotential is generated, there is a problem still. The good candidate for the potential minimum is a supersymmetric vacuum, but it leads to a negative vacuum energy if the gravitino mass $m_{3/2}$ is non-vanishing. It is not straightforward to realize the de Sitter vacuum with the almost vanishing vacuum energy, where moduli fields are stabilized. One may obtain such vacuum when we consider rather complicated superpotential and Kähler potential of moduli fields including several free parameters.

Recently, in Ref. [1] a new scenario was proposed to lead to a de Sitter (or Minkowski) vacuum, where all of moduli are stabilized in type IIB string models, and it is the so-called KKLT scenario. The KKLT scenario consists of three steps. At the first step, it is assumed that the dilaton and complex structure moduli are stabilized through flux compactification [2]. At the second step, non-perturbative superpotential terms, which depend on the Kähler moduli, are introduced. Thus, such superpotential stabilizes remaining Kähler moduli. However, that leads to a supersymmetric anti de Sitter (AdS) vacuum. At the third step, the AdS vacuum is uplifted by introducing anti-D3 branes, which break SUSY explicitly, at the tip of warp throat.

Phenomenological aspects like soft SUSY breaking terms have been studied [3]. The KKLT scenario predicts the unique pattern of SUSY breaking terms, where modulus mediation and anomaly mediation [4] are comparable for gaugino masses, scalar masses and A-terms. Recently, such type of SUSY breaking mediation is called as the mirage mediation, and this pattern of s-particle spectrum has significant phenomenological implications [5]-[9].

In this paper, we study on the third step of the KKLT scenario, i.e. uplifting. In the KKLT scenario, the uplifting potential is realized by explicit SUSY breaking. Here, we consider the possibility for uplifting by spontaneous SUSY breaking within supersymmetric theory, i.e. the F-term uplifting. Such possibility has been studied in Ref. [10]. Here we study F-term uplifting scenario by use of concrete SUSY breaking models. We consider the Polonyi model [11] and Intriligator-Seiberg-Shih (ISS) model [12] as the source of spontaneous SUSY breaking. (See also for recent works related with the ISS model Ref. [13].) Then, we require almost vanishing vacuum energy. For this purpose, we fine-tune parameters. A difference between our approach and the KKLT scenario is that in our approach the vacuum energy is assumed to vanish within supergravity theory, while the vacuum energy is canceled with uplifting energy from outside of supergravity in the KKLT scenario. That might lead to phenomenological aspects different from those in the

KKLT scenario. We study soft SUSY breaking terms and show the modulus and anomaly mediation are comparable in the gaugino masses. The size of scalar masses depends on how to couple with the spontaneous SUSY breaking sector.

This paper is organized as follows. In the next section, we review on the Polonyi model and the ISS model. We consider the ISS model within the framework of supergravity. In section 3, we combine the moduli stabilization with the Polonyi model and the ISS model to realize de Sitter (Minkowski) vacuum. We analyze potential minima and evaluate the magnitude of modulus F-term. In section 4, we study soft SUSY breaking terms. Section 5 is devoted to conclusion and discussion.

2 SUSY breaking models

In this section, we give a brief review on spontaneous SUSY breaking models, which shall be used as the source of uplifting in the next section. We consider the Polonyi model and ISS model. We use the unit with $M_{Pl} = 1$, where M_{Pl} denotes the reduced Planck scale.

2.1 Polonyi model

The Polonyi model is given by the following Kähler potential K and superpotential W :

$$K = |\Phi|^2, \quad W = c + \mu^2 \Phi,$$

where Φ is the Polonyi field and c and μ^2 are constants. The scalar potential

$$V = e^G (G^{I\bar{J}} G_I G_{\bar{J}} - 3), \quad G = K + \ln |W|^2,$$

is minimized by a real VEV, $\langle \Phi \rangle = \phi$, satisfying the stationary condition

$$V_\Phi \Big|_0 = e^G G_\Phi (G_{\Phi\Phi} + G_\Phi^2 - 2) \Big|_0 = 0,$$

where

$$\begin{aligned} (G_{\Phi\Phi} + G_\Phi^2 - 2) \Big|_0 &= \tilde{W}^{-1}(\phi^3 + \tilde{c}\phi^2 - 2\tilde{c}), \\ G_\Phi \Big|_0 &= \tilde{W}^{-1}(\phi^2 + \tilde{c}\phi + 1), \end{aligned}$$

$\tilde{W} = \mu^{-2}W|_0 = \phi + \tilde{c}$, $\tilde{c} = \mu^{-2}c$, and $f|_0$ denotes $f|_{\Phi=\langle\Phi\rangle}$ for a function $f = f(\Phi)$.

If the parameters c and μ^2 are within the range $-2 < \tilde{c} < 2$, we find $G_\Phi \Big|_0 > 0$ and the SUSY point is not a stationary point of the scalar potential. Instead, the potential has a SUSY breaking stationary point determined by

$$\phi^3 + \tilde{c}\phi^2 - 2\tilde{c} = 0.$$

Requiring that this SUSY breaking minimum has a vanishing vacuum energy

$$V\Big|_0 = e^G(G_\Phi^2 - 3)\Big|_0 = 0,$$

that is

$$(\phi^2 + \tilde{c}\phi + 1)^2 = 3(\phi + \tilde{c})^2,$$

the VEV ϕ and the parameter \tilde{c} are determined as

$$\phi = \sqrt{3} - 1, \quad \tilde{c} = 2 - \sqrt{3}.$$

This is the so-called Polonyi solution, which corresponds to a SUSY-breaking Minkowski minimum.

2.2 ISS model in supergravity

Here, we review briefly on the ISS model and study its supergravity extension. We consider the following Kähler and superpotential,

$$\begin{aligned} K &= \sum_{i,c} (|\phi_i^c|^2 + |\tilde{\phi}_c^i|^2) + \sum_{i,j} |\Phi_j^i|^2, \\ W &= h(\mu^2 \Phi_i^i - \phi_i^c \Phi_j^i \tilde{\phi}_c^j), \end{aligned}$$

where $i = 1, 2, \dots, N_f, c = 1, 2, \dots, N$ and $N = N_f - N_c$. This theory is dual to the $SU(N_c)$ theory with N_f flavors of “quarks” q^i and \bar{q}_i , which have the superpotential

$$W = h\mu^2 q\bar{q}. \tag{1}$$

In a global SUSY analysis of Ref. [12], this model has a SUSY breaking vacuum

$$\langle \phi \rangle = (\phi_0, 0), \quad \langle \tilde{\phi} \rangle = \begin{pmatrix} \tilde{\phi}_0 \\ 0 \end{pmatrix}, \quad \langle \Phi \rangle = \begin{pmatrix} 0 & 0 \\ 0 & \Phi_0 \end{pmatrix},$$

where $\tilde{\phi}_0\phi_0 = \mu^2 1_N$, and Φ_0 is a $(N_f - N) \times (N_f - N)$ matrix. The vacuum energy at this minimum is given by

$$V_0 = (N_f - N)|h\mu^2|^2.$$

Here we study the supergravity extension of this model. For simplicity, we consider the case with $N_f = 2$ and $N_c = 1$. Although the model with $N_c = 1$ has no dual theory, we can analyze the model with $N_c > 1$ in a similar way. We parameterize fields as

$$\phi_i = (\chi, \rho), \quad \tilde{\phi}^i = \begin{pmatrix} \tilde{\chi} \\ \tilde{\rho} \end{pmatrix}, \quad \Phi_j^i = \begin{pmatrix} Y & z \\ \tilde{z} & X \end{pmatrix}.$$

Then the system is described by

$$\begin{aligned} K &= |\chi|^2 + |\rho|^2 + |\tilde{\chi}|^2 + |\tilde{\rho}|^2 + |Y|^2 + |X|^2 + |z|^2 + |\tilde{z}|^2, \\ W &= c + h\mu^2(Y + X) - h(\chi\tilde{\chi}Y + \rho\tilde{\rho}X + \chi\tilde{\rho}z + \rho\tilde{\chi}\tilde{z}), \end{aligned}$$

where we add a constant superpotential term, c , for generality. The minimum obtained in the above global SUSY analysis corresponds to

$$\langle \chi \rangle = \langle \tilde{\chi} \rangle = \mu, \quad \langle X \rangle = \Phi_0,$$

and others are all vanishing. Note that these VEVs generate a SUSY mass of $O(h\mu)$ for most directions such as $\chi + \tilde{\chi}$ and Y , while X and the real part of $\chi - \tilde{\chi}$ receive a SUSY breaking mass of $O(h^2\mu/8\pi)$ at the 1-loop level. In addition, we have some Goldstone modes associated with the broken flavor symmetries as shown in Ref. [12]. The ISS model corresponds to generalized O’Raifeartaigh model.

On the other hand, supergravity scalar potential is given by

$$V = e^K (K^{I\bar{J}} (D_I W)(D_{\bar{J}} \bar{W}) - 3|W|^2),$$

where $D_I W = W_I + K_I W$ is given, e.g. as

$$\begin{aligned} D_\chi W &= -h(Y\tilde{\chi} + z\tilde{\rho}) + \bar{\chi}W, \\ D_\rho W &= -h(\tilde{z}\tilde{\chi} + \tilde{\rho}X) + \bar{\rho}W, \\ D_z W &= -h\chi\tilde{\rho} + \bar{z}W, \\ D_Y W &= -h\chi\tilde{\chi} + h\mu^2 + \bar{Y}W. \end{aligned}$$

Noticing that conditions $D_{\chi, \tilde{\chi}, \rho, \tilde{\rho}, z, \tilde{z}, Y} W = 0$ can be satisfied by

$$\rho, \tilde{\rho}, z, \tilde{z} = 0, \quad hY = O(W) = O(m_{3/2}), \quad \chi, \tilde{\chi} \simeq \mu,$$

we find that, similarly to the global SUSY case, most fields are stabilized (prior to X) by the SUSY condition $D_{\chi, \tilde{\chi}, \rho, \tilde{\rho}, z, \tilde{z}, Y} W = 0$ due to a large SUSY mass of $O(h\mu)$. Although some directions such as pseudo moduli and Goldstone modes should remain as light fields in the effective potential below, these effects are irrelevant to the following analysis of X which is essentially the source of SUSY breaking, and we will omit the terms involving these modes.

Then, the effective potential for the remaining X is described by

$$\begin{aligned} V &= V^{(0)} + V^{(1)}, \\ V^{(0)} &= e^K (K^{X\bar{X}} (D_X W)(D_{\bar{X}} \bar{W}) - 3|W|^2), \\ V^{(1)} &= m_X^2 |X|^2, \end{aligned}$$

where

$$K = |X|^2, \quad W = c + \mu^2 X.$$

In the superpotential, c and μ are redefined as $c + \langle (\mu^2 - \chi\tilde{\chi})hY \rangle \rightarrow c$, μ as $\sqrt{h}\mu \rightarrow \mu$, respectively. The mass m_X represents a SUSY breaking mass for X generated at the 1-loop level.

The form of the tree level scalar potential $V^{(0)}$ is the same as the Polonyi model analyzed in the previous section. (The Polonyi field Φ is now replaced by X .) However, we have a SUSY breaking mass for X in $V^{(1)}$. We expand the derivative of the scalar potential, $V_X = \partial_X V$, under the assumption that $c \sim \mu^2$, $\langle X \rangle = x \sim \mu^2$ and $\mu^2 \ll 1$ in the unit $M_{Pl} = 1$, and find

$$V_X = V_X^{(0)} + V_X^{(1)} = \mu^6 e^K W^{-1} \left(\mu^{-4} c \{ (\mu^{-1} m_X)^2 x - 2c \} + O(\mu^4) \right).$$

The stationary condition $V_X = 0$ results in

$$x \simeq 2 \left(\frac{\mu}{m_X} \right)^2 c. \quad (2)$$

Similarly, the scalar potential itself is expanded as

$$V = V^{(0)} + V^{(1)} = e^K (m_X^2 x^2 - 4c\mu^2 x + \mu^4 - 3c^2 + O(\mu^{10})),$$

and the condition for vanishing vacuum energy $V = 0$ as well as Eq. (2) leads to

$$c \simeq \frac{1}{\sqrt{3}} \mu^2 \left(1 - \frac{2}{3} \left(\frac{\mu}{m_X} \right)^2 \mu^2 \right). \quad (3)$$

These results are consistent with the assumption $x, c \sim \mu^2 \ll 1$ by recalling that $m_X \approx \mu$.

The vacuum value of the superpotential evaluated by Eqs. (2) and (3) is found to be

$$W \Big|_0 = c + \mu^2 x = \frac{1}{\sqrt{3}} \mu^2 + O(\mu^4),$$

and SUSY is broken at this Minkowski minimum due to

$$D_X W \Big|_0 = (W_X + K_X W) \Big|_0 = \mu^2 + O(\mu^4).$$

3 Moduli stabilization and F-term uplifting

Here, we study stabilization of the modulus T and uplifting. We assume the other moduli are stabilized at M_{Pl} , e.g. by flux background, but the single modulus T remains light. First, we consider the model, where the Kähler potential and superpotential are obtained as

$$K = -n_T \ln(T + \bar{T}), \quad W = c - A e^{-aT}.$$

The second term can be generated by gaugino condensation in the hidden sector, and in this case we have $A \sim 1$ and $a = O(10)$. Furthermore, we assume that $c \ll 1$. Its scalar potential is written as

$$V = e^K (K^{T\bar{T}} |D_T W|^2 - 3|W|^2).$$

The modulus T is stabilized at $D_T W = 0$ as $a\text{Re}(T) \simeq \ln A/c$, and the modulus mass is obtained as $m_T = am_{3/2}$. However, we obtain the negative vacuum energy $V = -3e^K |W|^2 < 0$. This corresponds to the second step in the KKLT scenario. In the KKLT scenario, the explicit SUSY breaking effect due to anti-D3 brane is added to uplift the potential and to realize de Sitter (Minkowski) vacuum. Here, we do not add such explicit SUSY breaking effect, but combine the moduli stabilization and spontaneous SUSY breaking models reviewed in the previous section to realize de Sitter (Minkowski) vacuum. Then, we study potential minima.

3.1 Polonyi-KKLT model

We study a combination of the Polonyi model and the KKLT-type model:¹

$$K = |\Phi|^2 - n_T \ln(T + \bar{T}), \quad W = c + \mu^2 \Phi - Ae^{-aT}.$$

As in the KKLT model, we assume

$$A \sim 1, \quad a \gg 1, \quad c, \mu^2 \ll 1,$$

in the unit $M_{Pl} = 1$.

The scalar potential is now a function of two chiral superfields Φ and T , and complicated. Therefore, we first find a ‘reference point’ which seems to be close to the genuine stationary point, and then estimate the deviation from the stationary point.

We define the reference point $(\Phi_0, T_0) = (\phi, t)$ such that the following conditions are satisfied there:

$$\begin{aligned} V_\Phi \Big|_0 &= 0, & (D_\Phi W \Big|_0 &\neq 0) \\ D_T W \Big|_0 &= 0, \end{aligned}$$

where $D_I W = W_I + K_I W$ for $I = (\Phi, T)$, and $f|_0 = f|_{\Phi=\phi, T=t}$ for a function $f = f(\Phi, T)$. The first condition $V_\Phi|_0 = 0$ is easily solved just by interpreting as $\tilde{c} = \mu^{-2}(c - Ae^{-at})$ in the previous analysis for the Polonyi model, and we find

$$\phi = \sqrt{3} - 1, \quad \mu^{-2}(c - Ae^{-at}) = 2 - \sqrt{3}.$$

¹ A similar model has been studied for another reason in Ref. [14].

The true minimum is represented by

$$\langle \Phi \rangle = \Phi_0 + \delta\Phi, \quad \langle T \rangle = T_0 + \delta T,$$

where $\delta\Phi/\Phi_0 \ll 1$ and $\delta T/T_0 \ll 1$ are assumed. The superpotential and its derivatives are expanded as

$$\begin{aligned} W &= W\Big|_0 + \partial_T W\Big|_0 \delta T + \dots \simeq \mu^2 + aAe^{-at} \delta T, \\ D_T W &= D_T W\Big|_0 + \partial_T D_T W\Big|_0 \delta T + \dots \simeq -a^2 Ae^{-at} \delta T, \\ D_\Phi W &= W_\Phi + K_\Phi W = \mu^2 + (\sqrt{3} - 1)(\mu^2 + aAe^{-at} \delta T + \dots) \\ &\simeq \sqrt{3}\mu^2 + (\sqrt{3} - 1)aAe^{-at} \delta T, \end{aligned}$$

where the ellipsis stands for the sub-dominant terms for which the coefficients are not enhanced by $a \gg 1$ or higher-order terms in powers of δT . From this, we find $\bar{F}^{\bar{T}} = -e^{K/2} K^{\bar{T}T} D_T W \simeq e^{K/2} K^{\bar{T}T} a^2 Ae^{-at} \delta T$, and then the scalar potential is expanded as

$$\begin{aligned} V &= K_{I\bar{J}} F^I \bar{F}^{\bar{J}} - 3e^K |W|^2 \\ &= e^K (|D_\Phi W|^2 - 3|W|^2) + K_{T\bar{T}} \bar{F}^{\bar{T}} F^T \\ &= e^K \Big|_0 \left\{ (D_\Phi \bar{W}) \Big|_0 (\sqrt{3}\mu^2 + (\sqrt{3} - 1)aAe^{-at} \delta T) - 3\bar{W} \Big|_0 (\mu^2 + aAe^{-at} \delta T) \right\} \\ &\quad + K_{T\bar{T}} e^{K/2} K^{\bar{T}T} \Big|_0 a^2 Ae^{-at} \delta T F^T + \dots, \end{aligned}$$

where $F^I = -e^{K/2} K^{I\bar{J}} D_{\bar{J}} \bar{W}$ for $I, J = (\Phi, T)$, and again the ellipsis represents sub-dominant terms.

Noticing $D_\Phi W\Big|_0 = \sqrt{3}\mu^2$ and $W\Big|_0 = \mu^2$, we obtain

$$V \simeq e^K (ae^{-K/2} F^T - \sqrt{3}\mu^2) aAe^{-at} \delta T.$$

Requiring $\partial V/\partial(\delta T) = 0$, the F-term F^T is determined as

$$F^T \simeq \sqrt{3}a^{-1} e^{K/2} W\Big|_0,$$

which is suppressed by a large factor $a \gg 1$. The order parameter F^T is vanishing at the reference point $F^T\Big|_0 = 0$, and generated by the small deviation $\delta\bar{T} (= \delta T)$ as $F^T \approx -e^{K/2} K^{\bar{T}T} a^2 Ae^{-at} \delta\bar{T}$. Then, from the above value of F^T , we can estimate δT as

$$\delta T/T_0 \approx \frac{1}{(at)^2} \ll 1,$$

where we have adopted $W\Big|_0 = -K_T^{-1} W_T\Big|_0 = -(K_T)^{-1}\Big|_0 aAe^{-at}$.

Then, as expected, the true minimum is very close to the reference point. The small deviation δT yields the small SUSY breaking order parameter F^T compared to F^Φ and $m_{3/2} = e^{K/2}W$. Particularly, the ratio between the anomaly mediation and modulus mediation is given by

$$\alpha = \frac{m_{3/2}}{\ln(M_{Pl}/m_{3/2})} \frac{T + \bar{T}}{F^T} \simeq \frac{at}{2\sqrt{3}\pi^2} \approx \frac{2}{\sqrt{3}},$$

where $\ln(M_{Pl}/m_{3/2}) \simeq 4\pi^2$ for $m_{3/2} = O(10)$ TeV, and $at \simeq 4\pi^2$ is determined by $D_T W|_0 = 0$.

This is compared to the original KKLT model, $\alpha_{KKLT} \simeq 1$. We remark that, in the KKLT model, SUSY is broken by the anti-D3 brane which generates an explicit SUSY breaking in the 4D effective $N = 1$ supergravity, while SUSY is broken within the $N = 1$ supergravity model studied in this section.

3.2 ISS-KKLT model

Here, we study a combination of the ISS model and the KKLT-type model:

$$\begin{aligned} V &= V^{(0)} + V^{(1)}, \\ V^{(0)} &= e^K (K^{I\bar{J}} (D_I W)(D_{\bar{J}} \bar{W}) - 3|W|^2), \\ V^{(1)} &= m_X^2 |X|^2, \end{aligned}$$

where $I, J = (X, T)$ and

$$K = |X|^2 - n_T \ln(T + \bar{T}), \quad W = c + \mu^2 X - Ae^{-aT}.$$

As in the analysis of Polonyi-KKLT model, we shall find a minimum of this model by the perturbation from the reference point $(X, T) = (X_0, T_0) = (x, t)$ where $V_X|_0 = 0$, $D_X W|_0 \neq 0$ and $D_T W|_0 = 0$ are satisfied. The first condition $V_X|_0 = 0$ is solved just by replacing c by $\tilde{c} = c - Ae^{-at}$ in the previous analysis of the pure ISS model. Then we find

$$x = 2 \left(\frac{\mu}{m_X} \right)^2 \tilde{c}.$$

The true minimum is assumed to be located close to the reference point,

$$\langle X \rangle = X_0 + \delta X, \quad \langle T \rangle = T_0 + \delta T,$$

where $\delta X/X_0, \delta T/T_0 \ll 1$.

Similarly to the Polonyi-KKLT model, the superpotential and its derivatives are expanded as

$$\begin{aligned} W &\simeq \frac{1}{\sqrt{3}} \mu^2 + aAe^{-at} \delta T, \\ D_T W &\simeq -a^2 Ae^{-at} \delta T, \\ D_X W &\simeq \mu^2. \end{aligned}$$

Then, the scalar potential is given by

$$\begin{aligned} V &= V^{(0)} + V^{(1)} \\ &\simeq e^K \Big|_0 (m_X^2 x^2 - 4\tilde{c}\mu^2 x + \mu^4 - 3\tilde{c}^2) + aAe^{-at}(ae^{K/2}F^T - \sqrt{3}\mu^2)\delta T. \end{aligned}$$

The stationary condition $\partial V/\partial(\delta T)$ determines F^T as

$$F^T \simeq 3a^{-1}e^{K/2}W \Big|_0,$$

and the condition for vanishing vacuum energy is the same as the pure ISS model besides the replacement $c \rightarrow \tilde{c}$, i.e.,

$$\tilde{c} \simeq \frac{1}{\sqrt{3}}\mu^2 \left(1 - \frac{2}{3} \left(\frac{\mu}{m_X}\right)^2 \mu^2\right).$$

The anomaly-to-modulus ratio for the SUSY breaking mediation in this case is given by

$$\alpha \simeq \frac{at}{6\pi^2} \approx \frac{2}{3}.$$

Note that $\alpha_{ISS-KKLT} \simeq \alpha_{P-KKLT}/\sqrt{3}$.

3.3 Stringy origin

In the ISS-KKLT model, the gravitino mass $m_{3/2}$ is determined by the constant c and supersymmetric mass μ^2 . To realize low-energy SUSY, we need suppressed values of c and μ^2 compared with M_{Pl} . Here we comment on what can be a source of such suppressed terms. Recall our first assumption, that is, we have assumed that all of moduli except the modulus T are stabilized at M_{Pl} . We denote these heavy moduli representatively by S . When these heavy moduli S appear in low-energy effective theory, they can be replaced by their VEVs. It is plausible that non-perturbative effects like gaugino condensation and string/D-brane instanton effects generate

$$W = Ce^{-\gamma S} + C'e^{-\gamma' S} q\bar{q},$$

where C, C' are $O(1)$ of constants.² The coefficients γ and γ' are constants determined by discrete numbers, e.g. beta-function coefficients for gaugino condensates. These terms become sources for c and μ^2 when we replace S by its VEV. Thus, when $\gamma\langle S \rangle$ and $\gamma'\langle S \rangle$ are of $O(10)$, we would have suppressed values of c and μ^2 . Furthermore, if $\gamma = \gamma'$, we expect $c = O(\mu^2)$, although we need fine-tuning between C and C' to realize almost vanishing vacuum energy.

² The possibility of the first term has been considered in Ref. [15, 16], and the possibility of the second term has been considered for the Higgs μ -term in Ref. [5, 7]. See also Ref. [17].

Furthermore, we have considered the simple case of T -dependent superpotential, i.e. $W_{np} = Ae^{-at}$ with $A = O(1)$ in the unit $M_{Pl} = 1$. However, when the gauge kinetic function of the hidden sector is written as a linear combination of S and T and this gaugino condensates, we would have the following superpotential term,

$$W_{np} = Ae^{-aT - a'S}.$$

instead of $W_{np} = Ae^{-aT}$. Using this superpotential term, we can analyze potential minima in a way similar to the previous section. Then, we have various values of α depending on a value of $a'S$ like Ref. [15], but its order would be obtained as $\alpha = O(1)$.

4 Soft SUSY breaking terms

Here we study soft SUSY breaking terms of the visible sector. The F-term of modulus F^T is smaller than the gravitino mass $m_{3/2}$ by $O(1/8\pi^2)$. That is, the modulus mediation and anomaly mediation are comparable, and its ratio α is of $O(1)$.

First, we evaluate gaugino masses, whose gauge kinetic functions are obtained as

$$f_a = k_a T + \Delta f_a,$$

where k_a and Δf_a are constants and Δf_a may depend on heavy moduli. The F^T contribution to gaugino mass is obtained as

$$M_a^{(T)} = \frac{k_a}{f_a + \Delta f_a} F^T.$$

In addition, there is the contribution from anomaly mediation,

$$M_a^{(AM)} = -\frac{\beta_{g_a^2}}{2g_a^2} m_{3/2},$$

where g_a is the gauge coupling and $\beta_{g_a^2}$ is the beta-function of g_a^2 . When $\alpha = O(1)$, these two contributions are comparable. That is the mirage mediation. If the Polonyi field and the field X in the ISS model appear in the gauge kinetic function, the situation would change.

Similarly, we evaluate scalar masses m_i of chiral multiplets Q^i in the visible sector. We may have several types of possibilities for assuming Kähler metric of chiral multiplets Q^i , in particular, how Q^i couple with the Polonyi field Φ and the field X in the ISS model. Here we consider three models. In the model I, the visible matter is separated from Φ and X in the form of $Y_i = e^{-K/3} Z_{i\bar{i}} |Q^i|^2$, where $Z_{i\bar{i}}$ is the Kähler metric of visible fields Q^i .³ Such assumption could be realized by the setup that the SUSY breaking source is

³ The authors would like to thank K. Choi for pointing this possibility.

localized far away from the visible matter fields in the compact space. In the model II, the Kähler metric $Z_{i\bar{i}}$ depends on only T and \bar{T} , but not Φ or X . Thus, the visible modes Q_i are not sequestered from Φ and X in Y_i . In the model III, the Kähler metric $Z_{i\bar{i}}$ includes a contact term like $c_i|X|^2|Q^i|^2/M_p^2$. The visible matter fields Q_i are not sequestered from Φ or X in Y_i or $Z_{i\bar{i}}$. We assume the almost vanishing vacuum energy, $V_0 \simeq 0$, in evaluation of scalar masses for all of three models.

In the model I, soft scalar masses squared are obtained at the tree-level as

$$m_i^2 = -|F^T|^2 \partial_T \partial_{\bar{T}} \ln Y_{i\bar{i}}, \quad (4)$$

where $Y_{i\bar{i}} = e^{-K/3} Z_{i\bar{i}}$. The F-term F^T is suppressed compared with $m_{3/2}$. Thus, the anomaly mediation is comparable with this tree-level effect. Then, we have the mirage mediation in soft scalar masses, too.

In the model II, soft scalar masses squared are obtained

$$m_i^2 = m_{3/2}^2 - |F^T|^2 \partial_T \partial_{\bar{T}} \ln Z_{i\bar{i}}.$$

Since the F-term F^T is suppressed compared with $m_{3/2}$, the first term is dominant in this case. Although there is the contribution from anomaly mediation, it is sub-dominant. Thus, in the model II, soft scalar masses are universal,

$$m_i^2 = m_{3/2}^2.$$

In the model III, there is a contact term like $c_i|X|^2|Q^i|^2/M_p^2$, soft scalar masses m_i also depend on $c_i|F^X|^2$. At any rate, the order of m_i is of $O(m_{3/2})$. In both models II and III, scalar masses are quite heavy compared with gaugino masses, and these would have radiative corrections.

Moreover, trilinear couplings of scalar fields, i.e. the so-called A-terms, can be calculated as

$$h_{ijk} = \sum_I F^I e^{K/2} [(\partial_I + K_I) Y_{ijk} - Y_{ijk} \partial_I \ln(Z_{i\bar{i}} Z_{j\bar{j}} Z_{k\bar{k}})],$$

where Y_{ijk} is the corresponding Yukawa coupling, and the index I includes the modulus T and the Polonyi field Φ and the field X of the ISS model. In the ISS-KKLT model, the VEV of X is suppressed and K_X is suppressed. Thus, the natural order of h_{ijk}/Y_{ijk} would be of $O(F^T/T)$. In this case, the anomaly mediation is also important and the size of h_{ijk}/Y_{ijk} is comparable with the gaugino masses. On the other hand, if the Yukawa coupling Y_{ijk} includes T -modulus like

$$Y_{ijk} \sim e^{-a'_{ijk} T}.$$

the contribution from F^T can be enhanced by the term $\partial_T Y_{ijk}$, and its order would be of $O(m_{3/2})$. In the model that all of Yukawa couplings are given in the above form, large

values of $|a'_{ijk}|$ correspond to suppressed Yukawa couplings. Thus, in such model, small Yukawa couplings like the first and second families would correspond to large A-terms of $O(m_{3/2})$, while large Yukawa couplings like the third family would correspond to smaller A-terms.

In the Polonyi-KKLT model, the VEV of Polonyi field is of $O(M_{Pl})$, and the dominant contribution to h_{ijk}/Y_{ijk} would be obtained as

$$\frac{h_{ijk}}{Y_{ijk}} = F^\Phi e^{K/2} K_\Phi + F^T e^{K/2} \partial_T \ln Y_{ijk}. \quad (5)$$

Then, it is naturally of $O(m_{3/2})$. However, when the Polonyi field Φ is sequestered from Q_i in Y_i , i.e. the model I, there is no contribution from F^Φ . Thus, when $\partial_T \ln Y_{ijk} = O(1)$, the size of h_{ijk}/Y_{ijk} would be comparable with the gaugino masses even in the Polonyi-KKLT model.

The magnitudes of the Higgs μ -term and the so-called B -term depend on how to generate the μ -term. Suppose that the μ -term is generated as

$$W_\mu = C e^{-hT} H_u H_d.$$

In this case, the would-be dominant part of B -term is obtained, e.g. in the ISS-KKLT model as

$$B = m_{3/2} - hF^T.$$

Thus, the natural scale of B is of $O(m_{3/2})$. If two terms $m_{3/2}$ and hF^T with $h = O(10)$ are canceled each other, then B can be of $O(m_{3/2}/8\pi^2)$. Indeed such cancellation happen in a certain case [7].

As a result, the full s-particle spectrum of our model is as follows. We choose the over-all mass scale such that the gaugino masses are of $O(100) - O(1000)$ GeV. When visible matter fields are sequestered from the spontaneous SUSY breaking sector, sfermion masses are of the same order as gaugino masses, and the gravitino are of $O(10)$ TeV and the mass of modulus T is of $O(100)$ TeV. When visible matter fields are not sequestered from the spontaneous SUSY breaking sector, the gravitino and sfermion masses are of $O(10)$ TeV. The size of A-terms can be of the same order as gaugino masses or gravitino masses, depending on T -dependence of Yukawa couplings and the spontaneous SUSY breaking mechanism. The natural scale of B -term is of $O(m_{3/2})$, but in a certain case we could obtain smaller value of B .

5 Conclusion and discussion

We have studied modulus stabilization with F-term uplifting. As explicit models, we have used the Polonyi model and the ISS model. Combining these spontaneous SUSY breaking

models and the KKLT type of superpotential, we have analyzed potential minima. At the potential minima, the size of modulus F-term F^T is similar to the KKLT model, and suppressed by a factor of $O(10)$ compared with the gravitino mass. We have also studied soft SUSY breaking terms. In the gaugino masses, modulus mediation and anomaly mediation are comparable, i.e. the mirage mediation. On the other hand, sfermion masses can be of the same order as the gaugino masses or of $O(m_{3/2})$, depending couplings with the spontaneous SUSY breaking sector. Thus, in the low-energy SUSY scenario, the gaugino masses are of $O(100) - O(1000)$ GeV, while sfermion masses are of the same order or $O(10)$ TeV. The gravitino mass is of $O(10)$ TeV and the modulus T is much heavier. The A-terms can be of the same order as the gaugino masses or gravitino masses. We have studied two explicit models with F-term uplifting, but these spectra would be expected of generic feature of F-term uplifting scenario. The natural size of B -term would be of $O(m_{3/2})$, but in a certain case it could be much smaller.

Recently, phenomenological aspects of the mirage mediation have been studied. However, spectra of our models, in particular sfermion masses as well as A-terms, can differ from the mirage mediation. It would be interesting to study phenomenological aspects of models, where the mirage mediation is dominant for the gaugino masses and sfermion masses are much heavier.⁴

Note to be added

While this paper was being finished, Ref. [20] appeared, where they also studied the same model as one of ours.

Acknowledgement

The authors would like to thank F. Brummer, K. Choi, T. Goto, A. Hebecker, K. Izawa, T. Onogi and M. Trapletti for useful discussions. H. A., T. H. and T. K. are supported in part by the Grand-in-Aid for Scientific Research #182496, #171643 and #17540251, respectively. T. K. is also supported in part by the Grant-in-Aid for the 21st Century COE “The Center for Diversity and Universality in Physics” from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- [1] S. Kachru, R. Kallosh, A. Linde and S. P. Trivedi, Phys. Rev. D **68**, 046005 (2003) [arXiv:hep-th/0301240].
- [2] S. B. Giddings, S. Kachru and J. Polchinski, Phys. Rev. D **66**, 106006 (2002) [arXiv:hep-th/0105097].

⁴ See e.g. Ref. [18, 19], where phenomenological aspects of similar s-particle spectra have been studied.

- [3] K. Choi, A. Falkowski, H. P. Nilles and M. Olechowski, Nucl. Phys. B **718**, 113 (2005) [arXiv:hep-th/0503216].
- [4] L. Randall and R. Sundrum, Nucl. Phys. B **557**, 79 (1999) [arXiv:hep-th/9810155]; G. F. Giudice, M. A. Luty, H. Murayama and R. Rattazzi, JHEP **9812**, 027 (1998) [arXiv:hep-ph/9810442].
- [5] K. Choi, K. S. Jeong and K. i. Okumura, JHEP **0509**, 039 (2005) [arXiv:hep-ph/0504037];
- [6] M. Endo, M. Yamaguchi and K. Yoshioka, Phys. Rev. D **72**, 015004 (2005) [arXiv:hep-ph/0504036]; A. Falkowski, O. Lebedev and Y. Mambrini, JHEP **0511**, 034 (2005) [arXiv:hep-ph/0507110]; K. Choi, K. Y. Lee, Y. Shimizu, Y. G. Kim and K. i. Okumura, arXiv:hep-ph/0609132.
- [7] K. Choi, K. S. Jeong, T. Kobayashi and K. i. Okumura, Phys. Lett. B **633**, 355 (2006) [arXiv:hep-ph/0508029];
- [8] R. Kitano and Y. Nomura, Phys. Lett. B **631**, 58 (2005) [arXiv:hep-ph/0509039]; O. Lebedev, H. P. Nilles and M. Ratz, arXiv:hep-ph/0511320; A. Pierce and J. Thaler, arXiv:hep-ph/0604192.
- [9] K. Choi and K. S. Jeong, arXiv:hep-th/0605108; J. P. Conlon and F. Quevedo, arXiv:hep-th/0605141; F. Brummer, A. Hebecker and M. Trapletti, Nucl. Phys. B **755**, 186 (2006) [arXiv:hep-th/0605232].
- [10] A. Saltman and E. Silverstein, JHEP **0411**, 066 (2004) [arXiv:hep-th/0402135]; M. Gomez-Reino and C. A. Scrucca, JHEP **0605**, 015 (2006) [arXiv:hep-th/0602246]; O. Lebedev, H. P. Nilles and M. Ratz, Phys. Lett. B **636**, 126 (2006) [arXiv:hep-th/0603047].
- [11] J. Polonyi, unpublished preprint KFKI-77-93, 1977.
- [12] K. Intriligator, N. Seiberg and D. Shih, JHEP **0604**, 021 (2006) [arXiv:hep-th/0602239].
- [13] S. Franco and A. M. Uranga, JHEP **0606**, 031 (2006) [arXiv:hep-th/0604136]; H. Ooguri and Y. Ookouchi, arXiv:hep-th/0606061; Phys. Lett. B **641**, 323 (2006) [arXiv:hep-th/0607183]; V. Braun, E. I. Buchbinder and B. A. Ovrut, Phys. Lett. B **639**, 566 (2006) [arXiv:hep-th/0606166]; JHEP **0610**, 041 (2006) [arXiv:hep-th/0606241]; S. Ray, Phys. Lett. B **642**, 137 (2006) [arXiv:hep-th/0607172]; S. Franco, I. Garcia-Etxebarria and A. M. Uranga, arXiv:hep-th/0607218; S. Forste, arXiv:hep-th/0608036; A. Amariti, L. Girardello and A. Mariotti, arXiv:hep-th/0608063; I. Bena, E. Gorbatov, S. Hellerman,

- N. Seiberg and D. Shih, arXiv:hep-th/0608157; M. Dine, J. L. Feng and E. Silverstein, arXiv:hep-th/0608159; C. Ahn, arXiv:hep-th/0608160; arXiv:hep-th/0610025; M. Eto, K. Hashimoto and S. Terashima, arXiv:hep-th/0610042; R. Argurio, M. Bertolini, S. Franco and S. Kachru, arXiv:hep-th/0610212; M. Aganagic, C. Beem, J. Seo and C. Vafa, arXiv:hep-th/0610249; S. A. Abel, C. S. Chu, J. Jaeckel and V. V. Khoze, arXiv:hep-th/0610334; N. J. Craig, P. J. Fox and J. G. Wacker, arXiv:hep-th/0611006.
- [14] M. Dine, R. Kitano, A. Morisse and Y. Shirman, Phys. Rev. D **73**, 123518 (2006) [arXiv:hep-ph/0604140].
- [15] H. Abe, T. Higaki and T. Kobayashi, Phys. Rev. D **73**, 046005 (2006) [arXiv:hep-th/0511160].
- [16] H. Abe, T. Higaki and T. Kobayashi, Phys. Rev. D **74**, 045012 (2006) [arXiv:hep-th/0606095].
- [17] R. Blumenhagen, M. Cvetič and T. Weigand, arXiv:hep-th/0609191; L. E. Ibanez and A. M. Uranga, arXiv:hep-th/0609213.
- [18] J. D. Wells, Phys. Rev. D **71**, 015013 (2005) [arXiv:hep-ph/0411041].
- [19] M. Ibe, T. Moroi and T. T. Yanagida, arXiv:hep-ph/0610277.
- [20] E. Dudas, C. Papineau and S. Pokorski, arXiv:hep-th/0610297.