

(Q uantum) tw isted Y angians: sym m etry, B axterisation and centralizers

N icolas C ram pe^{1a,b} and A nastasia D oikou^{2c}

^a International School for Advanced Studies,
V ia Beirut 2-4, 34014 Trieste, Italy

^b Istituto Nazionale di Fisica Nucleare
Sezione di Trieste

^c University of Bologna, Physics Department, INFN Section,
V ia Imerio 46, 40126 Bologna, Italy

A bstract

Based on the (quantum) tw isted Yangians, integrable system s with special boundary conditions, called soliton non-preserving (SNP), may be constructed. In the present article we focus on the study of subalgebras of the (quantum) tw isted Yangians, and we show that such a subalgebra provides an exact sym m etry of the rational transfer m atrix. W e discuss how the spectrum of a generic transfer m atrix m ay be obtained by focusing only on two types of special boundaries. It is also show n that the subalgebras, em erging from the asym ptotics of tensor product representations of the (quantum) tw isted Yangian, tum out to be dual to the (quantum) Brauer algebra. To deal with general boundaries in the trigonom etric case we propose a new algebra, which also provides the appropriate fram ew ork for the Baxterisation procedure in the SNP case.

¹ cram pe@ sissa .it

² doikou@ bo .infn .it

1 Introduction

It is well established by now that, for any \mathfrak{gl}_n algebra or the corresponding quantum deformation ($U_q(\mathfrak{gl}_n)$), one may consider integrable lattice models or quantum field theories with two distinct types of boundary conditions known as soliton preserving (SP) with underlying algebra the reflection algebra [1, 2] or soliton non-preserving (SNP) with underlying algebra the twisted Yangian [3] or quantum twisted Yangian [4]. Historically, soliton non-preserving boundary conditions were first introduced in the context of a new Toda field theories on the half line [5], whereas solutions of the (quantum) twisted Yangian first found in [6]. Nonetheless up to date, both SP and SNP boundary conditions have been extensively studied in the context of integrable quantum spin chains (see e.g. [7, 8, 9]).

In this paper, we focus on the SNP case and we provide a generic description of the underlying symmetry algebra. The rational and trigonometric cases are considered separately. For the rational case, we investigate the symmetry and we show that the algebra, emerging from the twisted Yangian as $j|j| - 1$, is isomorphic to $so(p; q)$ or $sp(n)$ and is an exact symmetry of the algebraic open transfer matrix for a particular relation between the left and right boundaries. In addition, for this particular relation between the boundaries, we show that it is sufficient to study the spectrum of the transfer matrix for only two special cases. Finally, we recall the duality between the Brauer algebra and this algebra [10].

Similar considerations are made for the trigonometric case. In this case, the corresponding finite twisted quantum algebra, obtained from the quantum twisted Yangian for $j|j| - 1$, does not seem to provide an exact symmetry of the corresponding open model. We recall however the link between the finite twisted quantum algebra and the quantum Brauer algebra established in [11] for a trivial boundary. In the case of a generic boundary, we need to introduce a new algebra. We prove that a subalgebra of this new algebra is the centralizer of the finite quantum twisted Yangian. The new framework allows us to find the spectral depending solution via a Baxterisation procedure.

The outline of this paper is as follows: In the next section, we give the algebraic setting by introducing the FRT presentation [12] for quantum groups and some subalgebras. We show for the rational case that the set of non local charges emerging from the asymptotic expansion of the tensor solution of the twisted Yangian, turn out to provide an exact symmetry of the transfer matrix. We then prove that the spectrum of any transfer matrix with generic boundary conditions can be deduced from the transfer matrix with special diagonal boundaries. We show that the entailed symmetry algebra commutes with appropriate representations of the Brauer algebra, depending on the choice of boundary conditions. In the last section, the trigonometric case is considered. Surprisingly the emerging boundary quantum algebra is not an exact symmetry of the trigonometric transfer matrix contrary to the rational case. Nevertheless, for trivial boundary conditions, we recall the duality between the quantum Brauer algebra and the twisted boundary quantum algebra. We finally provide a general framework, which allows us to deal with generic boundary conditions, and also discuss about the Baxterisation procedure.

2 (Quantum) twisted Yangian: general setting

Let $R(\lambda)$ be a solution of the Yang-Baxter equation [13, 14, 15]

$$R_{12}(\lambda_1, \lambda_2) R_{13}(\lambda_1) R_{23}(\lambda_2) = R_{23}(\lambda_2) R_{13}(\lambda_1) R_{12}(\lambda_1, \lambda_2); \quad (2.1)$$

acting on $\text{End}(\mathbb{C}^n)^{\otimes 3}$, and as usual $R_{12}(\lambda) = R(\lambda) \otimes I$; $R_{23}(\lambda) = I \otimes R(\lambda)$ and so on. In addition, the R matrices we shall deal with satisfy

(i) the unitarity condition

$$R_{12}(\lambda) R_{21}(\lambda) = I \quad (2.2)$$

(ii) the crossing relation

$$R_{12}^{t_i}(\lambda) M_{-1} R_{12}^{t_j}(\lambda - 2i) M_{-1}^{-1} = I; \quad (2.3)$$

where t_i denotes the transposition on the i^{th} space

(iii) the symmetry

$$M_{-1} M_2; R_{12}(\lambda) = 0; \quad (2.4)$$

Let us also define the matrix

$$R_{12}(\lambda) / R_{12}^{t_i}(\lambda - i); \quad (2.5)$$

which will be useful for our purposes here. The proportional factors have no influence on the definition of the algebra and will be chosen conveniently.

For an associative infinite algebra Y generated by $\{L_{ij}^{(m)} \mid 1 \leq i, j \leq n; m = 0, 1, \dots, g\}$, one may define the object

$$L(\lambda) = \sum_{i,j=1}^n E_{ij} L_{ij}(\lambda) \in \text{End}(\mathbb{C}^n) \otimes Y[\lambda^{-1}]; \quad (2.6)$$

where the second 'space' is occupied by formal series $L_{ij}(\lambda) = \sum_{m=0}^{\infty} \frac{L_{ij}^{(m)}}{\lambda^m}$ and E_{ij} is a n by n matrix such that $(E_{ij})_{kl} = \delta_{ik} \delta_{jl}$. The exchange relations may be written using the so-called FRT relation [12],

$$R_{12}(\lambda_1, \lambda_2) L_{13}(\lambda_1) L_{23}(\lambda_2) = L_{23}(\lambda_2) L_{13}(\lambda_1) R_{12}(\lambda_1, \lambda_2); \quad (2.7)$$

where the 1 and 2 stand for the two copies of $\text{End}(\mathbb{C}^n)$ whereas 3, for the space $Y[\lambda^{-1}]$.

In this article, we will be interested in two particular choices of the algebra called the $gl(n)$ (quantum) Yangian [16, 17] depending on the choice of a rational or trigonometric R matrix.

In this type of algebra defined by FRT relation, we can exhibit different homomorphisms. In particular, we will use $L(\lambda) \nabla L(\lambda) = f(\lambda) L^t(\lambda)$ i.e. in terms of generators (entries of L)

$$L_{ij}(\lambda) \nabla L_{ij}(\lambda) = f(\lambda) L_{ji}(\lambda) \quad (2.8)$$

where $f(\lambda)$ is any function chosen later conveniently.

The (quantum) Yangian defined by relation (2.7) are also Hopf algebras. In particular, they are equipped with a coproduct $\Delta: Y \rightarrow Y \otimes Y$ given by, for $i, j \in \{1, \dots, n\}$,

$$(L_{ij}(\lambda)) = \sum_{a=1}^{X^n} L_{aj}(\lambda) \otimes L_{ia}(\lambda) \quad (2.9)$$

which can be written also as

$$(\text{id} \otimes \Delta) L(\lambda) = L_{02}(\lambda) \otimes L_{01}(\lambda) \quad (2.10)$$

The index 0 stands for the space $\text{End}(\mathbb{C}^n)$, whereas 1 and 2 stand for the two copies of the algebra. This coproduct induces also

$$(\text{id} \otimes \Delta) L(\lambda) = L_{01}(\lambda) \otimes L_{02}(\lambda) \quad (2.11)$$

The Δ -coproduct $\Delta^{(2)}: Y \rightarrow Y \otimes Y$ is defined by the following iteration $\Delta^{(2)} = \text{id} \otimes \Delta^{(1)}$ with $\Delta^{(1)} = \Delta$.

In order to deal with integrable systems with non trivial boundary conditions, one has to consider appropriate subalgebras of Y . For this purpose, we define

$$K_0(\lambda) = \sum_{i,j=1}^{X^n} E_{ij} \quad K_{ij}(\lambda) = L_{01}(\lambda) K_0(\lambda) L_{01}(\lambda); \quad (2.12)$$

with $K(\lambda)$ a c -number matrix solution of the following equation

$$R_{12}(\lambda_1 - \lambda_2) K_1(\lambda_1) R_{12}(\lambda_1 + \lambda_2) K_2(\lambda_2) = K_2(\lambda_2) R_{12}(\lambda_1 + \lambda_2) K_1(\lambda_1) R_{12}(\lambda_1 - \lambda_2) \quad (2.13)$$

The generators encompassed in $K(\lambda)$ generate a subalgebra T of Y , whose exchange relations are given by (2.13), where now $K(\lambda)$ is replaced by $K(\lambda)$. Depending on the choice of the R matrix, T is the (quantum) twisted Yangian³ [3, 4]. It is clear that $K(\lambda)$ allows the expansion in powers of λ^{-1} as we shall see in subsequent section, providing explicit forms of the generators of the twisted Yangian.

The subalgebra T is not a Hopf algebra but has a structure of co-ideal inherited essentially from the (quantum) Yangian, $\Delta: T \rightarrow T \otimes Y$, such that (see also [18, 19]), for $i, j \in \{1, \dots, n\}$,

$$(K_{ij}(\lambda)) = \sum_{a,b=1}^{X^n} K_{ab}(\lambda) \otimes L_{ia}(\lambda) L_{bj}(\lambda) \quad (2.14)$$

³In the case of the twisted Yangian, a supplementary relation is required for K (see (3.7))

or, equivalently, $(\text{id} \otimes K) = L_{02} L_{01} K_0 L_{01} L_{02}$. One can exploit the existence of tensor product realizations of the (quantum) twisted Yangian in order to build the corresponding quantum system that is the open quantum spin chain with SNP boundary conditions. We define

$$T_0(\lambda) = (\text{id} \otimes^{(N)}) L(\lambda) = L_{0N}(\lambda) \cdots L_{01}(\lambda); \quad (2.15)$$

$$\bar{T}_0(\lambda) = (\text{id} \otimes^{(N)}) L(\lambda) = L_{01}(\lambda) \cdots L_{0N}(\lambda); \quad (2.16)$$

Then, the general tensor type solution of the (2.13) takes the form

$$B_0(\lambda) = (\text{id} \otimes^{(N)}) K(\lambda) = T_0(\lambda) K_0^{(R)}(\lambda) \bar{T}_0(\lambda); \quad (2.17)$$

where $K_0^{(R)}(\lambda)$ is a c-number matrix solution of (2.13) interpreted as the reflection on the right boundary. The entries $B_{ij}(\lambda)$ of the matrix $B(\lambda)$ can be computed explicitly by $B_{ij}(\lambda) = \text{tr}^{(N)}(K_{ij}(\lambda))$.

Finally, we introduce the transfer matrix of the open spin chain [2], which may be written as

$$t(\lambda) = \text{tr}_0 \left(K_0^{(L)}(\lambda) B_0(\lambda) \right); \quad (2.18)$$

where $K_0^{(L)}(\lambda) = K(\lambda, i)$ encodes the interaction with the left boundary and $K(\lambda)$ is a solution of (2.13). It can be shown [2, 7], using the fact that $B(\lambda)$ is a solution of (2.13), that transfer matrix (2.18) provides family of commuting operators i.e.,

$$[t(\lambda); t(\lambda')] = 0; \quad (2.19)$$

The latter commutation relation (2.19) ensures the integrability of the relevant models.

3 The rational case

One of the main objectives of the present study is the derivation of the exact symmetry for integrable models associated to (quantum) twisted Yangian. We shall first examine the rational case and, in the subsequent section, we shall proceed with the trigonometric case.

The gl_n R matrix is written in the following simple form,

$$R(\lambda) = I + \frac{i}{\lambda} P \quad (3.1)$$

where $P = \sum_{a \neq b=1}^n E_{ab} E_{ba}$ is the permutation operator, acting on $(\mathbb{C}^n)^{\otimes 2}$. Note that for the

rational case $\lambda = \frac{n}{2}$ and $M = I$ (see (2.3)). Let us define $P = I - Q$ where $Q = \sum_{a \neq b=1}^n E_{ab} E_{ab}$

is a one-dimensional projector satisfying

$$Q P = P Q = Q; \quad Q^2 = nQ; \quad (3.2)$$

and consequently $P^2 = -2I$. Then, the R matrix can be written as

$$R(\lambda) = I + \frac{i}{\lambda} P \quad (3.3)$$

For the rational case, a solution of the fundamental equation (2.7) is provided by evaluation map i.e.

$$L_{ij}(\lambda) = e_{ij} + \frac{ie_{ji}}{\lambda}; \quad L_{ij}(\lambda) = \frac{\lambda + i}{\lambda} L_{ji}(\lambda) \quad (i) = e_{ij} + \frac{i - e_{ij}}{\lambda} \quad ie_{ij} \quad (3.4)$$

where e_{ij} are the generators of the Lie algebra $gl(n)$ satisfying

$$[e_{ij}, e_{kl}] = \delta_{jk} e_{il} - \delta_{il} e_{kj} \quad (3.5)$$

Let us introduce $P = \sum_{ij} E_{ij} - E_{ji}$ and $\tilde{P} = \sum_{ij} E_{ij} - (\sum_{ij} E_{ij})$. Then, we can write

$$L(\lambda) = 1 + \frac{iP}{\lambda} \quad \text{and} \quad \tilde{L}(\lambda) = 1 + \frac{i\tilde{P}}{\lambda} \quad (3.6)$$

It is clear that P and \tilde{P} are the fundamental representations of the corresponding P and \tilde{P} .

3.1 Symmetry of the transfer matrix

We shall also need in what follows the c-number solution of (2.13) for the rational case, which is any constant matrix (independent) such that $K = K^t$ [8, 20]. We shall show that different choices of K provide different symmetry algebras. We suppose also that K is invertible and with real entries. We recall that there does not exist invertible antisymmetric $n \times n$ matrix for n odd (Jacobi's theorem easily checked by $\det(K) = (-1)^n \det(K)$). For this choice of R matrix, the elements encompassed in $B(\lambda)$ generated the twisted Yangian [3]. The elements satisfied the following supplementary symmetry relation⁴

$$B^t(\lambda) = B(\lambda)g(\lambda) + \frac{i}{2 + i} B(\lambda)g(\lambda) - B(\lambda); \quad (3.7)$$

where the upper (resp. lower) sign corresponds to K symmetric (resp. antisymmetric) and $g(\lambda) = \frac{\lambda + i}{\lambda} \lambda^{2N}$.

We shall consider here the expansion of $B(\lambda)$ as λ^{-1} in order to obtain explicit expressions of generators of the twisted Yangian (see also [21]). We shall keep up to λ^{-1} terms which is sufficient for our purposes. Bearing in mind the expansions of $L(\lambda)$ and $\tilde{L}(\lambda)$, we may easily deduce the asymptotic behavior of $B(\lambda)$ as λ^{-1} , i.e.

$$B_0(\lambda^{-1}) = K_0 + \frac{i}{\lambda} B_0^{(1)} + \dots = K_0 + \frac{i}{\lambda} \sum_{i=1}^N (K_0 P_{0i} + P_{0i} K_0) + \dots \quad (3.8)$$

⁴The differences between the relation presented here and this one in [3] are due to the shift in the spectral parameter of the matrix $R(\lambda)$ in the commutation relation (2.13) and to the factor in $L(\lambda)$.

Let us define the following combination emerging essentially from the asymptotic expansion of the generalized solution of (2.13)

$$K^{(1)} = K P + P K \quad \text{i.e.} \quad K_{ab}^{(1)} = K_{ab} \sum_{c=1}^{X^n} (K_{ac} e_{cb} - e_{ca} K_{cb}) : \quad (3.9)$$

The nonlocal charges i.e. the entries of (3.8), may be written simply as coproducts of the twisted Yangian elements namely $B^{(1)} = \sum_{c=1}^{X^n} (K^{(1)})$. In particular, the asymptotic expansion of (2.14) provides

$$(K^{(1)}) = K^{(1)} I + I K^{(1)} = K_0 P_{01} + P_{01} K_0 + K_0 P_{02} + P_{02} K_0 : \quad (3.10)$$

The elements $K_{ab}^{(1)}$ are called primitive and, in particular, their coproduct is cocommutative.

In the following, we will study more precisely the algebra $T^{(1)}$ spanned by $B_{ab}^{(1)}$ but, before that, we shall show that they commute with the transfer matrix of the system provided that special boundary conditions are considered. In the proof we shall simply exploit the underlying algebraic relations provided by (2.13) as $\sum_{i=1}^n 1$ i.e.

$$B_1^{(1)} ; B_2(\cdot) = P_{12} B_1(\cdot) K_2 - K_1 B_2(\cdot) + K_1 Q_{12} B_2(\cdot) - B_2(\cdot) Q_{12} K_1 : \quad (3.11)$$

Then, we extract the element in the position (a;b) in the space 1 and (c;d) in the space 2

$$B_{ab}^{(1)} ; B_{cd}(\cdot) = K_{ad} B_{cb}(\cdot) - K_{cb} B_{ad}(\cdot) + K_{ac} B_{bd}(\cdot) - K_{db} B_{ca}(\cdot) : \quad (3.12)$$

From the latter exchange relations it is entailed that for a generic $K^{(R)}$ matrix the elements $B_{ab}^{(L)}$ do not commute with the transfer matrix. If however we consider the case where the left boundary is $K^{(L)} = K^{(R)-1}$, the transfer matrix becomes

$$t(\cdot) = \sum_{c=1}^{X^n} K_{lc}^{(L)} B_{cl}(\cdot) : \quad (3.13)$$

From the latter equation and bearing in mind the exchange relations (3.12), it is easy to show that

$$t(\cdot) ; B_{ab}^{(1)} = 0 : \quad (3.14)$$

Note that an alternative proof may be formulated along the lines of [7, 22]. The particular choice of the left boundary indicates that the two boundaries of the chain have to be appropriately tuned so that all the elements $B_{ab}^{(L)}$ commute with the open transfer matrix. In fact, the special cases: (i) $K_{ab}^{(L)} = K_{ab}^{(R)}$, (ii) $K_{ab}^{(L)} = K_{a+1 b}^{(R)}$, (iii) $K_{ab}^{(L)} = (-1)^a K_{a+1 b}^{(R)}$ fall to the category above. The symmetry of cases (ii) and (iii) has been studied also in [9]. One may extract valuable information concerning the exact symmetry of a chain with generic boundary conditions exploiting the algebraic relations (3.12).

Now, we describe more precisely the algebra $T^{(1)}$ spanned by $\{B_{ab}^{(1)}\}_{a,b=1}^n$. We are going to show that in the case K symmetric it is isomorphic to the Lie algebra $so(p; q)$ whereas for K antisymmetric it is isomorphic to $sp(n)$. For convenience, we introduce the sign $\epsilon = \pm 1$ such that $K^t = \epsilon K$. By expanding expressions (3.7) and (3.12) up to $\mathcal{O}(\hbar^1)$, we get

$$B_{ba}^{(1)} = B_{ab}^{(1)} + 2N K_{ab} \quad (3.15)$$

$$B_{ab}^{(1)}; B_{cd}^{(1)} = K_{ad} B_{cb}^{(1)} - K_{cb} B_{ad}^{(1)} + K_{ac} B_{bd}^{(1)} - K_{db} B_{ca}^{(1)} \quad (3.16)$$

Let us now introduce the following $n \times n$ matrices

$$G^+ = \text{diag}(\underbrace{1, \dots, 1}_p; \underbrace{1, \dots, 1}_q) \quad \text{and} \quad G = \text{diag}(\underbrace{1, \dots, 1}_{n=2}) \quad \begin{matrix} 0 & 1 \\ 1 & 0 \end{matrix}; \quad (3.17)$$

where $p+q = n$ and the second case is valid only for n even. A well-known linear algebra result is that the matrices G are the normal forms of the matrix K over reals under congruence i.e. there exists an invertible real matrix U such that

$$UKU^t = G \quad (3.18)$$

The mapping

$$B^{(1)} \mapsto U(B^{(1)} - NK)U^t G = UB^{(1)}K^{-1}U^{-1} - N = M \quad (3.19)$$

$$\text{i.e. } B_{ab}^{(1)} \mapsto \sum_{i=1}^n U_a B^{(1)}_{i} - NK \quad U G_b = M_{ab} \quad (3.20)$$

defines an algebra isomorphism from $T^{(1)}$ to $so(p; q)$ (resp. to $sp(n)$) for $\epsilon = +$ (resp. $\epsilon = -$). The bijection is proven by the fact that U is invertible. By direct computation of the relations satisfying by M_{ab} starting from relations (3.15) and (3.16), we recognize the following defining relations of $so(p; q)$ (resp. $sp(n)$)

$$M^t G = -G M \quad (3.21)$$

$$[M_1; M_2] = [M_2; P_{12} - G_1 Q_{12} (G_1)^{-1}] \quad (3.22)$$

which show the algebra homomorphism.

3.2 Treatment of general boundary condition

Henceforth we focus on the fundamental representation i.e. $L_{0i}(\lambda) \in \mathbb{R}$ and $R_{0i}(\lambda) \in \mathbb{R}$ and we restrict ourselves to the case where $K^{(R)} = K^{(L)}^{-1} = K$. Then, we get the following representation for $B(\lambda)$,

$$B_0(\lambda) \in \mathbb{R} \quad R_{ON}(\lambda) \in \mathbb{R} \quad R_{01}(\lambda) \in \mathbb{R} \quad K_0 R_{01}(\lambda) \in \mathbb{R} \quad R_{ON}(\lambda) = B_0(\lambda) \quad (3.23)$$

In this case, we have the following important proposition which allows us to find the spectrum of the transfer matrix for any boundary condition by studying only the case where the boundary conditions are given by G .

Proposition 1 Let K be any solution of (2.13) and $t_K(\lambda) = \text{tr}_0(K_0^{-1}B_0(\lambda))$. There exists invertible matrix U such that $UKU^t = G$ where G are defined in (3.17). Let

$$t_G(\lambda) = \text{tr}_0(G_0)^{-1}R_{0N}(\lambda) \cdots R_{01}(\lambda)G_0R_{01}(\lambda) \cdots R_{0N}(\lambda) \quad (3.23)$$

Then $t_K(\lambda)$ and $t_G(\lambda)$ have the same eigenvalues, their eigenvectors (say V_K and V_G respectively) being related through

$$V_G = U_1 \cdots U_N V_K \quad (3.24)$$

Proof. The proof of this proposition is based on the relation

$$U_1 \cdots U_N t_K(\lambda) = t_G(\lambda) U_1 \cdots U_N \quad (3.25)$$

which is obtained using the property $[R_{0i}(\lambda); (U_0^t)^{-1}U_i] = 0$ and the symmetry relation (2.4). ■

This result is similar to the one obtained in the soliton preserving case in [8, 23, 24, 25]. This proposition allows us to restrict the study of the spectrum of the transfer matrix only to the cases where the boundaries are given by G . Let us emphasize that although the eigenvalues are identical the models obtained from t_K and t_G may be different.

3.3 Link with the Brauer algebra

We shall now focus on the relation between the symmetry algebra and the Brauer algebra [10]. A well-known presentation of the classical Brauer algebra $B_N(n)$ is realized by $2N - 2$ generators e_i and f_i ($1 \leq i \leq N - 1$) obeying exchange relations:

$$\begin{aligned} e_i^2 &= 1; & f_i^2 &= -n; & e_i e_i &= e_i e_i = -n; & i &= 1; \dots; N - 1 \\ e_i e_j &= e_j e_i; & f_i f_j &= f_j f_i; & e_i f_j &= f_j e_i; & |i - j| &> 1 \\ e_i e_{i+1} e_i &= e_{i+1} e_i e_{i+1}; & f_i f_{i-1} f_i &= -f_i; \\ e_i e_{i+1} e_i &= e_{i+1} e_i; & f_{i+1} f_i f_{i+1} &= f_{i+1} f_i; & i &= 1; \dots; N - 2 \end{aligned} \quad (3.26)$$

By direct computation, it is clear that the following map

$$e_i \mapsto P_{i+1} \quad \text{and} \quad f_i \mapsto K_i Q_{i+1} K_i^{-1}; \quad (3.27)$$

is a representation of the Brauer algebra $B_N(n)$ for any matrix K satisfying $K^t = -K$ with $n = \text{tr}(K^2)$.

By representing relation (3.8), the conserved quantities in the fundamental representation are given by

$$B_0^{(1)} = \sum_{j=1}^N P_{0j} K_0 + K_0 P_{0j} \quad (3.28)$$

and generate an algebra isomorphic to $so(p; q)$ and $sp(n)$ depending on the choice of the K matrix. Using relation (3.27), it is now straightforward to show that the conserved quantities are the centralizers of the Brauer algebra i.e.

$$[P_{i+i+1}; B_0^{(1)}] = 0 \quad \text{and} \quad [K_i Q_{i+i+1} K_i^{-1}; B_0^{(1)}] = 0 : \quad (3.29)$$

A few comments are in order at this point. In the SP case, the presence of a non trivial right boundary modifies naturally the form of the non-local charges, nevertheless they still commute with the transfer matrix as long as the left boundary is trivial. Furthermore, these charges continue to be the centralizers of an extended Hecke algebra called the B-type Hecke algebra (see [26]). In the case we examine here, the boundary non-local charges consist a symmetry algebra for the transfer matrix as long as the left and right boundaries are closely interrelated and are also centralizers of the Brauer algebra. In this spirit, it seems pointless to consider a 'boundary' extension of the Brauer algebra analogously to the Hecke case. However it is possible to conceive a generalization of the classical Brauer algebra regarding the K matrix as a representation of the extra element of the 'extended' algebra. Indeed, let us consider a possible extension of the Brauer algebra by introducing two supplementary generators b and b^{-1} , satisfying

$$b b^{-1} = b^{-1} b = 1; \quad b_j b = b_j; \quad b_j^{-1} b = b_j^{-1} \quad \text{for } j \geq 2 \quad (3.30)$$

$$b_1 b^{-1} b = b^{-1} b b_1; \quad (3.31)$$

$$b_1 b^{-1} b = b^{-1} b b_1; \quad b_1^{-1} b^{-1} b_1^{-1} b^{-1} = b^{-1} b_1^{-1} b^{-1} b_1^{-1} : \quad (3.32)$$

It is easy to prove that the following map

$$i \mapsto P_{i+i+1}, \quad i \mapsto Q_{i+i+1}, \quad b \mapsto K_1 \quad \text{and} \quad b^{-1} \mapsto K_1^{-1} \quad (3.33)$$

is a representation of this extended Brauer algebra. In the SP case, the representation of the extended Hecke algebra is important since it provides by Baxterisation new solutions of the reflection algebra. In our case, unfortunately, the Baxterisation does not apply given that no numerical solution depending on the spectral parameter of (2.13) exists.

The picture presented above offers a rather tempting interpretation of the K matrix in terms of the 'boundary' generator. However, another interpretation exists where the K matrix allows us to define new interaction in the bulk. Let us define, for any $K^t = K$,

$$R_{12}^0(\lambda) = K_1 R_{12}(\lambda) K_1^{-1} = K_2 R_{12}(\lambda) K_2^{-1} : \quad (3.34)$$

The transfer matrix, in the case where $(K^{(L)})^{-1} = K^{(R)} = K$, may be written as

$$t_K(\lambda) = \text{tr}_0 f R_{ON}(\lambda) :: R_{01}(\lambda) R_{01}^0(\lambda) :: R_{0N}^0(\lambda) g : \quad (3.35)$$

The commutativity of the transfer matrix (which provides, as usual, the integrability of the system) is ensured by

$$R_{12}(\lambda_1, \lambda_2) B_1^0(\lambda_1) R_{12}^0(\lambda_1 + \lambda_2) B_2^0(\lambda_2) = B_2^0(\lambda_2) R_{12}^0(\lambda_1 + \lambda_2) B_1^0(\lambda_1) R_{12}(\lambda_1, \lambda_2) \quad (3.36)$$

where $B_0^0(\lambda) = B_0(\lambda)K_0^{-1}$. The crucial point here is that the first non trivial terms in the expansion of $B_0^0(\lambda)$ for $|\lambda| \rightarrow 1$ i.e.

$$B_0^{0(1)} = \sum_{j=1}^N P_{0j} + K_0 \sum_{j=1}^N P_{0j} K_0^{-1} = N + \sum_{j=1}^N P_{0j} K_0 Q_{0j} K_0^{-1} \quad (3.37)$$

still provides the symmetry of the transfer matrix. In addition, it is still the centralizer of the representation of the Brauer algebra, i.e.

$$[P_{i i+1}; B_0^{0(1)}] = 0 \quad \text{and} \quad [K_i Q_{i i+1} K_i^{-1}; B_0^{0(1)}] = 0 \quad i=2, \dots, N-1 \quad (3.38)$$

This construction is more symmetrical in the sense that the same operator appears in the symmetry algebra (3.37) and in the representation of the Brauer algebra (3.27).

Within this frame of mind, it seems rather unnecessary to discuss about a 'boundary extension' of the Brauer algebra. To conclude we showed the duality between the symmetry algebra of the transfer matrix and the Brauer algebra for suitable representations, depending on the choice of boundaries.

4 The trigonometric case

The $U_q(\mathfrak{sl}_n)$ R matrix derived in [27] may be written in a compact form as

$$R(\lambda) = 2 \sinh(\lambda + i) P + 2 \sinh(\lambda) P U \quad \text{and} \quad U = \sum_{\substack{ij=1 \\ i \neq j}}^N (E_{ij} E_{ji} - q^{\text{sgn}(i-j)} E_{ii} E_{jj}) \quad (4.1)$$

where $q = e^i$. In this case, $n=2$ and the matrix M (2.3) is defined as

$$M_{ij} = q^{n-2j+1} E_{ij} \quad (4.2)$$

Defining $R_{12}[q] = R_{12} = P(q + U)$, we can write the R matrix more symmetrically as follows

$$R(\lambda) = e^{-i R_{12}} e^{R_{21}^{-1}} = e^{-i R_{12}[q]} e^{R_{12}[q^{-1}]^{t_1 t_2}} \quad (4.3)$$

The R matrix is given by

$$R(\lambda) = e^{-i R_{12}^{t_1}} e^{+i (R_{21}^{-1})^{t_1}} \quad (4.4)$$

In particular, we get $R(\lambda) = (q - q^{-1})Q$. It is well known that rational R matrix is a limit of trigonometric R matrix. Indeed, rescaling the spectral parameter λ in definition (4.1) of the trigonometric R matrix and taking the limit $\lambda \rightarrow 0$, we get rational R matrix (3.1). This limit, called the scaling limit, will be useful to compare the results of sections 3 and 4.

A solution of equation (2.7), where now R is a trigonometric matrix given above, may be written in the following simple form (see also [28])

$$L(\lambda) = e^{L^+} e^{-L}; \quad L^+ = \sum_{i,j=1}^n E_{ij} \psi_{ij}^+; \quad L = \sum_{i,j=1}^n E_{ij} \psi_{ij}; \quad (4.5)$$

with the matrices L^+ (L) apparently upper (lower) triangular and ψ_{ij}^+, ψ_{ij} the generators of the finite quantum group $U_q(\mathfrak{gl}_n)$. Its exchange relations can be also written using the FRT presentation as follows

$$R_{12} L_1 L_2 = L_2 L_1 R_{12} \quad (4.6)$$

$$R_{12} L_1^+ L_2 = L_2 L_1^+ R_{12}; \quad (4.7)$$

We have the following additional constraints on diagonal elements, for $1 \leq i \leq n$,

$$\psi_{ii}^+ \psi_{ii} = \psi_{ii} \psi_{ii}^+ = 1; \quad (4.8)$$

Similarly to the rational case, we introduce $L(\lambda) = L(\lambda - i)^t$ which can be written as

$$L(\lambda) = e^{L^+} e^{-L}; \quad L^+ = \sum_{i,j=1}^n E_{ij} \psi_{ij}^+; \quad L = \sum_{i,j=1}^n E_{ij} \psi_{ij}; \quad (4.9)$$

where $\psi_{ij} = e^{-i} \psi_{ji}$.

In [29], a classification of numerical solutions of equation (2.13) for trigonometric R matrix has been presented. More precisely, the general invertible solutions $K(\lambda)$, up to a global factor, can be written as

$$K(\lambda) = D G(\lambda) D \quad (4.10)$$

where D is any invertible constant matrix and $G(\lambda)$ is one of the following matrices

(i) $G(\lambda) = I$

(ii)
$$\frac{G(\lambda)}{q^{\frac{1}{2} + q^{-\frac{1}{2}}}} = e^{q^{\frac{n}{4}} \frac{B}{q}} \frac{q^{-1}}{q^{-1} + 1} \sum_{i=1}^n E_{ii} + \sum_{i,j=1}^n E_{ij} \frac{C}{A} e^{-q^{-\frac{n}{4}} \frac{B}{q}} \frac{q}{q+1} \sum_{i=1}^n E_{ii} + \sum_{i,j=1}^n E_{ij} \frac{C}{A}$$

where $\frac{1}{i} = \frac{p}{(1)^{in}}$ and by convention $\frac{1}{i < j} = 1$; $\frac{1}{i > j} = i$. The solution given here is linked to the solution given in [29] by transformation (4.10) where $D = \sum_{i=1}^n p \frac{X^n}{(1)^{in}} E_{ii}$ and by a global factor.

(iii)
$$G(\lambda) = e^2 q^{\frac{n-1}{2}} E_{1n} e^{-2} q^{-\frac{n-1}{2}} E_{n1} + \sum_{i=1}^{\frac{n-2}{2}} (q^{\frac{1}{2}} E_{2i 2i+1} - q^{\frac{1}{2}} E_{2i+1 2i}); \quad n \text{ even.}$$

$$(iv) G(\lambda) = \sum_{i=1}^n \frac{X^i}{q^{2i}} (q^{\frac{1}{2}} E_{2i-1, 2i} - q^{\frac{1}{2}} E_{2i, 2i-1}) = G_q; \quad n \text{ even.}$$

Notice that the solutions (i), (ii) coincide with previously known results found in [6, 18]. By the scaling limit, solutions (i) and (iv) become respectively G^+ (for $q=0$) and G^- (see relation (3.17)). Solution (iii) becomes $E_{1n} + E_{n1} + \sum_{i=1}^{\frac{n-2}{2}} E_{2i, 2i+1} - E_{2i+1, 2i}$ which is antisymmetric. Solution (ii), with the upper sign, becomes $\sum_{i,j=1}^n E_{ij}$ which is never invertible and, with the lower sign, $\sum_{i < j} (E_{ij} - E_{ji})$ which is antisymmetric. We recover different invertible solutions of the reflection equation with the rational R matrix and some non invertible solutions.

Finally, let us remark that each solution can be written as

$$G(\lambda) = G^+(\lambda; q) + G^-(\lambda; q^{-1})^t \quad (4.11)$$

where $G^+(\lambda; q)$ is equal to (i) $I=2$ ($\lambda = +$), (ii) $e^{-\frac{\lambda}{q^4}} \sum_{i=1}^n \frac{q}{q+1} E_{ii} + \sum_{i < j} E_{ij}$ ($\lambda = -$), (iii) $e^{\frac{\lambda}{q^2}} \sum_{i=1}^{\frac{n-1}{2}} E_{1n} + \sum_{i=1}^{\frac{n-2}{2}} q^{\frac{1}{2}} E_{2i, 2i+1}$ ($\lambda = +$) and (iv) $\sum_{i=1}^{\frac{n}{2}} q^{\frac{1}{2}} E_{2i-1, 2i}$ ($\lambda = -$). Note the similarity between relations (4.3) and (4.11) and the fact that $G^+(\lambda; q)$ is upper diagonal as R_{12} .

4.1 Finite subalgebras

The asymptotic expansion of $B(\lambda)$ as $\lambda \rightarrow 1$ will provide explicit expressions of the finite quantum twisted Yangian generators. We shall keep independent expressions, bearing in mind the form of L (4.5) and L (4.9) and also the generic solutions given above. It is clear that the first three solutions preserve the triangular decomposition of the $B(\lambda \rightarrow 1) = B = \sum_{i,j=1}^n E_{ij} - B_{ij}$ matrix, while for the last one B is not triangular anymore. Explicit expressions of B^+ for the various solutions (i) { (iv) are given below

$$(i) \text{ For } i < j, B_{ij}^+ = \sum_{k=i}^{X^j} v_{ik}^+ v_{kj}^+, B_{ii}^+ = e^i \text{ and for } i > j, B_{ij}^+ = 0.$$

$$(ii) \text{ For } i < j, B_{ij}^+ / \frac{q^{-1} X^j}{q^{-1} + 1} \sum_{k=i}^2 v_{ik}^+ v_{kj}^+ + \sum_{k=i}^{X^{-1} X^j} v_{kp}^+ v_{pj}^+, B_{ii}^+ / \frac{2e^i q^{-1}}{q^{-1} + 1} \text{ and for } i > j, B_{ij}^+ = 0$$

$$(iii) B_{1n}^+ / v_{11}^+ v_{nn}^+ \text{ and } 0 \text{ otherwise.}$$

$$(iv) B_{ij}^+ / q^{\frac{1}{2}} \sum_{k=\frac{i+1}{2}}^{X} v_{i, 2k-1}^+ v_{2k, j}^+ q^{\frac{1}{2}} \sum_{k=\frac{j}{2}}^{X} v_{i, 2k}^+ v_{2k-1, j}^+ \text{ for } i \leq j+1 \text{ and } 0 \text{ otherwise.}$$

We have used property (4.8) and the relations between v and v . Similarly, it is possible to compute explicitly the charges B_{ij} in terms of v_{ij} and v_{ij} .

The charges B_{ij} form an algebra, called T_f (f for finite), with the following defining exchange relations emerging from (2.13) as $q \rightarrow 1$, i.e. we get

$$R_{12} B_1^+ R_{12} B_2^+ = B_2^+ R_{12} B_1^+ R_{12} \quad (4.12)$$

$$R_{12} B_1 R_{12}^t B_2 = B_2 R_{12}^t B_1 R_{12} \quad (4.13)$$

$$R_{12} B_1^+ R_{12} B_2 = B_2 R_{12} B_1^+ R_{12} \quad (4.14)$$

$$R_{12} B_1^+ R_{12}^t B_2 = B_2 R_{12}^t B_1^+ R_{12} \quad (4.15)$$

where $R_{12} = (R_{21}^{-1})^t$. In [26, 30], analogous results have been obtained in the SP case.

Remarks (1) The subalgebra, denoted T_f^+ , generated by B_{ij}^+ is in fact isomorphic to the one, denoted T_f , generated by B_{ij} : it is proved using the invertible transformation $B_1^+ = M_1 (B_1^+)^{-1}$ and showing that B^+ satisfies relation (4.13).

(2) When $K(\cdot)$ is the solution (i) (resp. solution (iv)), the subalgebra T_f is the twisted quantum algebra $U_q^0(\mathfrak{so}_n)$ (resp. $U_q^0(\mathfrak{sp}_n)$): these algebras were introduced in [31, 32].

(3) For the solution (iii), there are only two non-vanishing generators B_{1n}^+ and B_{n1} . In this case, relations (4.12)–(4.15) reduce only to $[B_{1n}^+, B_{n1}] = 0$ which is an abelian algebra and is non-deformed.

(4) The algebra associated to the solution (ii) was discussed in [18], but here is treated in detail and gives interesting non-trivial results as we shall see in the subsequent sections.

In the following, we will not study the case (iii) since it provides a trivial finite algebra.

4.2 Symmetry of the transfer matrix

We shall present in what follows the exchange relations among the finite algebras studied previously and the entries of the transfer matrix (i.e. $\text{tr}_0 B_0(\cdot)$). Such relations will essentially manifest the existing (if any) symmetries of the model. More precisely, let us consider (2.13) for $q \rightarrow 1$. Bearing in mind that $R_{12}(q \rightarrow 1) / R_{12}$ and $R(q \rightarrow 1) / R_{12} = (R_{21}^{-1})^t$, one obtains the following set of exchange relations:

$$R_{12} B_1^+ R_{12} B_2(\cdot) = B_2(\cdot) R_{12} B_1^+ R_{12} : \quad (4.16)$$

Similar expressions may be derived for the case that $q \rightarrow 1$, but they are omitted here for brevity. Explicit expressions for the $U_q(\mathfrak{gl}_3)$ case are provided in Appendix A.

Let us focus on the charges next to diagonal, their explicit expressions for solution (ii) coincide with the ones found in [18] i.e.

$$B_{i,i+1}^+ = v_{ii}^+ v_{i,i+1}^+ + v_{i,i+1}^+ v_{i+1,i+1}^+ + (q+1) v_{ii}^+ v_{i+1,i+1}^+ : \quad (4.17)$$

For $K = I$, the expression is the same with the last term vanishing. Let $c = q - q^{-1}$. Then,

from (4.16), we obtain:

$$\begin{aligned}
 & \sum_{i=1}^X c_{i+1,j} B_{i,j}(\lambda) - q \sum_{i=1}^X c_{i,j} B_{i+1,j}(\lambda) + \sum_{j=1}^X f_j E_{j,l} B_{i,j+1}^+(\lambda) B_{j,l}(\lambda) \\
 = & \sum_{l=1}^X c_{i,l} B_{l,i+1}(\lambda) - q \sum_{l=1}^X c_{l,i+1} B_{l,i}(\lambda) + \sum_{l=j}^X f_j E_{l,j} B_{l,j}(\lambda) B_{i,j+1}^+(\lambda) : \quad (4.18)
 \end{aligned}$$

From the latter equations, one concludes that $B_{i,j+1}^+$ do not commute with the SNP transfer matrix for any choice of left boundary (see also Appendix A for the full set of exchange relations for the $U_q(\mathfrak{gl}_3)$ case) contrary to the SP case and to the rational SNP case where the exchange relations of the (4.16) allow the study of symmetry of the transfer matrix with boundary conditions. The higher order expansion of the transfer matrix provides naturally non-local quantities, which due to integrability commute with the transfer matrix, however such quantities being in involution form an abelian algebra. The main challenge within this context is the search of a non abelian algebra, which at the same time would form a symmetry of the transfer matrix as in the case of SP boundary conditions.

4.3 Quantum Brauer algebra

In this subsection, we basically recall the results obtained in [11]. The quantum deformation of the Brauer algebra $B_N(z; q)$ [11] is generated by $\{T_i, P_i, M_i\}_{i=1}^N$ subject to the following defining relations

$$\begin{aligned}
 T_i^2 &= (q - q^{-1}) T_{i+1}; & T_i T_j &= T_j T_i; & T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1}; \\
 T_i^2 &= \frac{z - z^{-1}}{q - q^{-1}} T_i; & P_i P_i &= P_i; & P_i &= q^{-1} \\
 T_i T_j T_i &= z T_j; & T_i P_i &= P_i T_i; & T_i &= 2; \dots; N-1 \\
 T_i (z q^{-1} + z^{-1} q) T_i &= (q^{-1} + q) T_i & & & & \\
 T_i (z q^{-1} + z^{-1} q) T_i &= (q^{-1} + q) T_i & & & & (4.19)
 \end{aligned}$$

where $T_i = T_{i+1} T_i T_{i+1}$. Notice that the generators f_i form the usual Hecke algebra $H_N(q)$. It has been shown in [11] that the map

$$T_i \mapsto U_{i,i+1} + q = P_{i,i+1} R_{i,i+1} = \hat{R}_{i,i+1} \quad \text{and} \quad P_i \mapsto P_{12}^i M_i^{-1} \quad (4.20)$$

is a representation of $B_N(q^{\pm 1}; q)$. The difference between the algebra defined here and the one introduced in [11] is such that, although we chose a different definition for the coproduct $T(\lambda)$, the quantum Brauer algebra is still the centralizer of the quantum twisted Yangian.

As in the rational case, we focus on the fundamental representation of $U_q(\mathfrak{gl}_n)$ acting on $(\mathbb{C}^n)^{\otimes N}$ i.e.

$$T(\lambda) \mapsto R_{0N}(\lambda) \dots R_{01}(\lambda); \quad (4.21)$$

which provides a representation of $B(\cdot) \nabla R_{ON}(\cdot) :: R_{01}(\cdot) K_0(\cdot) R_{01}(\cdot) :: R_{ON}(\cdot)$. In particular, we get

$$B_0^+ \nabla R_{ON} :: R_{01} K_0^+ R_{01} :: R_{ON} = B_0^+ \quad (4.22)$$

$$B_0 \nabla R_{N0}^{-1} :: R_{10}^{-1} K_0 R_{01}^{t_0} :: R_{ON}^{t_0} = B_0 \quad (4.23)$$

where $K = K(\cdot | \cdot | \cdot)$ and we recall that $R_{0i} = (R_{i0}^{-1})^{t_0}$.

A.I.M.olev, in [11], focused on the solution (i) ($K = I$) i.e. $U_q^0(\mathfrak{so}_n)$ and proved, in this case, that the actions of the quantum Brauer algebra $B_N(q^n; q)$ (see (4.20)) on the space $(\mathbb{C}^n)^{\otimes N}$ and the one of the algebra T_f , generated by B_{ij} , (see (4.23)) commute with each other i.e.

$$[U_{i;i+1} + q; B_0] = 0 \quad \text{and} \quad [P_{12}^{t_1} M_{1^{-1}}; B_0] = 0 : \quad (4.24)$$

We can prove similarly the same result for the whole algebra T_f (generated by B_{ij}).

Unfortunately, it seems impossible to prove similar results for all the solutions $K(\cdot)$. In addition, the matrices R and K cannot be written just using matrices similar to the ones representing \hat{R}_i and \hat{R}_i^{-1} whereas it has been possible in the rational case (see end of section 3.3). These remarks indicate that we need a more general framework we will introduce in the following section.

4.4 Generalized quantum Brauer algebra and Baxterisation

It is well known that the trigonometric R matrix can be obtained from generators of the Hecke algebra: such an operation is called Baxterisation. Indeed, starting from $\hat{R}_{i;i+1}$ satisfying Hecke algebra, the following linear combination

$$\hat{R}_{i;i+1}(\cdot) = e \hat{R}_{i;i+1} - e^{-1} \hat{R}_{i;i+1}^{-1} \quad (4.25)$$

satisfies the braided Yang-Baxter equation

$$\hat{R}_{12}(\cdot | \cdot | \cdot) \hat{R}_{23}(\cdot | \cdot | \cdot) \hat{R}_{12}(\cdot | \cdot | \cdot) = \hat{R}_{23}(\cdot | \cdot | \cdot) \hat{R}_{12}(\cdot | \cdot | \cdot) \hat{R}_{23}(\cdot | \cdot | \cdot) \quad (4.26)$$

and unitarity condition $\hat{R}(\cdot) \hat{R}(\cdot) / I$. As explained before, it does not seem possible to construct $R(\cdot)$ and $K(\cdot)$ in a similar fashion. However, there exist equivalent forms where the Baxterisation for these two matrices becomes possible.

Let us introduce the conjugate index $k = n + 1 - k$, the matrix $V = q^k \frac{N+1}{2} E_{kk}$ for n odd and $V = i(-1)^k q^k \frac{N+1}{2} E_{kk}$ for n even (such that $V^2 = I$ and $M = V^t V$) and $\mathfrak{B}_0(\cdot) = B_0(\cdot) V_0^t$. Then, we get from equation (2.13), the following braided reflection equation

$$\hat{R}_{12}(\cdot | \cdot | \cdot) \mathfrak{B}_1(\cdot | \cdot | \cdot) \mathfrak{R}_{12}(\cdot | \cdot | \cdot) \mathfrak{B}_1(\cdot | \cdot | \cdot) = \mathfrak{B}_1(\cdot | \cdot | \cdot) \mathfrak{R}_{12}(\cdot | \cdot | \cdot) \mathfrak{B}_1(\cdot | \cdot | \cdot) \hat{R}_{12}(\cdot | \cdot | \cdot) \quad (4.27)$$

where

$$\mathfrak{R}_{12}(\cdot) = P_{12} V_2^t R_{12}(\cdot) V_2^t = P_{12} V_2^t R_{12}^{t_1}(\cdot | \cdot | \cdot) V_2^t \quad (4.28)$$

Obviously, the algebras generated by $\mathbb{B}_0(\cdot)$ and $B_0(\cdot)$ are isomorphic. Whereas for $R(\cdot)$ there is no symmetric form, the matrix $\mathbb{R}(\cdot)$ can be written as

$$\mathbb{R}_{12}(\cdot) = e^{-i} \mathbb{R}_{12} e^{+i} \mathbb{R}_{12}^{-1} \quad (4.29)$$

where $\mathbb{R}_{12} = P_{12} V_2^t R_{12} V_2^t$.

All numerical solutions $\mathbb{R}(\cdot)$ of equation (4.27) can be obtained from the classification $K(\cdot)$ using the relation $\mathbb{R}(\cdot) = K(\cdot) V^t$. The only solution which gives a non-trivial finite algebra (see section 4.1) and depends on the spectral parameter is the solution (ii). The solution $\mathbb{R}(\cdot)$ corresponding to the cases (ii) can be also written symmetrically as

$$\mathbb{R}(\cdot) = e^{-q^{\frac{n}{4}}} \mathbb{R} e^{q^{\frac{n}{4}}} \mathbb{R}^{-1} \quad (4.30)$$

where $\mathbb{R} = \frac{1}{q^2 + q^{-2}} \left(\frac{q^{-1}}{q^{-1} + 1} \sum_i X_i^2 E_{ii} + \sum_{i < j} X_i X_j E_{ij} \right) V^t$.

We are now interested in the reciprocity: we want to find an algebra whose representation gives \hat{R} , \mathbb{R} and \mathbb{R} and which is sufficient to prove the different relations (Yang-Baxter equation, unitarity, reflection equation) that the matrices $\hat{R}(\cdot)$, $\mathbb{R}(\cdot)$ and $\mathbb{R}(\cdot)$ of the forms (4.25), (4.29) and (4.30) must satisfy. Let us define a new algebra, called $N_N(w; \cdot; q)$, generated by f_i, g satisfying the relation of the Hecke algebra and by the invertible generators f_i, g and f_i, g subject to the following defining relations,

$$f_i (f_i - 1) = w^2 (f_i - 1) \quad (4.31)$$

$$f_i g = g f_i, \quad f_i g = g f_i \quad \text{for } i > j \text{ or } i < j - 1 \quad (4.32)$$

$$f_i f_j = f_j f_i, \quad f_i f_j = f_j f_i \quad \text{for } |i - j| > 1 \quad (4.33)$$

$$f_{i+1} f_i f_{i+1} = f_i f_{i+1} f_i, \quad f_{i+1} f_i f_{i+1} = f_i f_{i+1} f_i \quad (4.34)$$

$$f_{i+1} f_i^{-1} f_{i+1} = f_i^{-1} f_{i+1} f_i, \quad f_{i+1} f_i^{-1} f_{i+1} = f_i^{-1} f_{i+1} f_i \quad (4.35)$$

$$f_{i+1} f_i^{-1} f_{i+1}^{-1} f_i = f_i^{-1} f_{i+1} f_i f_{i+1}^{-1} \quad (4.36)$$

$$f_{i+1} f_i f_{i+1}^{-1} f_i^{-1} f_{i+1} = f_i^{-1} f_{i+1} f_i f_{i+1}^{-1} \quad (4.37)$$

$$f_i b_i^{-1} b_i = b_i^{-1} b_i f_i, \quad f_i b_i b_i = b_i b_i f_i \quad (4.38)$$

$$f_i b_i b_i^{-1} b_i^{-1} b_i = b_i b_i f_i, \quad b_i^{-1} b_i^{-1} b_i^{-1} f_i \quad (4.39)$$

$$f_i b_i^{-1} b_i^{-1} b_i^{-1} b_i^{-1} b_i = b_i b_i^{-1} f_i, \quad b_i^{-1} b_i^{-1} f_i \quad (4.40)$$

$$f_i b_i^{-1} b_i^{-1} b_i^{-1} b_i^{-1} b_i^{-1} b_i = b_i b_i^{-1} b_i^{-1} f_i, \quad b_i^{-1} b_i^{-1} b_i^{-1} f_i \quad (4.41)$$

We define also two quotients of $N_N(w; \cdot; q)$, called $N_N^0(w; \cdot; q)$ (resp. $N_N^1(w; \cdot; q)$), defined by the supplementary relations

$$b_i^2 = 1 \quad (\text{resp. } (b_i + \overline{1} w^{-1})(b_i + b_i^{-1}) = 0); \quad (4.42)$$

where we recall that we used the convention $\overline{1} = 1$ and $\overline{1} = i$. Finally, let us introduce

$$\hat{f}_{i;i+1}(\cdot) = e^{-i} e^{-i^{-1}}, \quad \hat{e}_{i;i+1}(\cdot) = w^{-1} e^{-i} w e^{-i^{-1}} \quad (4.43)$$

and

$$k_i(\lambda) = \begin{cases} b_i & \text{for } N_N^0(w; \lambda; q) \\ e^{-w \frac{1}{2} b_i} + e^{-w \frac{1}{2} b_i^{-1}} & \text{for } N_N^1(w; \lambda; q); \end{cases} \quad (4.44)$$

where $\lambda = \lambda_i$ is a supplementary freedom. These algebras allow us to obtain the following proposition solving essentially the Baxterisation problem in the case of the quantum twisted Yangian:

Proposition 2 The generators $\hat{r}_{i,i+1}(\lambda)$ satisfy the braided Yang-Baxter equation. We get also

$$\hat{r}_{12}(\lambda_1, \lambda_2) \mathbf{e}_{23}(\lambda_1) \mathbf{e}_{12}(\lambda_2) = \mathbf{e}_{23}(\lambda_2) \mathbf{e}_{12}(\lambda_1) \hat{r}_{23}(\lambda_1, \lambda_2): \quad (4.45)$$

The generators $\hat{r}_{i,i+1}(\lambda)$, $\mathbf{e}_{i,i+1}(\lambda)$ and $k_i(\lambda)$ satisfy the braided reflection equation

$$\hat{r}_{12}(\lambda_1, \lambda_2) k_1(\lambda_1) \mathbf{e}_{12}(\lambda_1 + \lambda_2) k_1(\lambda_2) = k_1(\lambda_2) \mathbf{e}_{12}(\lambda_1 + \lambda_2) k_1(\lambda_1) \hat{r}_{12}(\lambda_1, \lambda_2): \quad (4.46)$$

We get also the following unitarity conditions

$$\frac{\hat{r}_{12}(\lambda) \hat{r}_{12}(\lambda^{-1})}{(q e^{-\lambda} - q^{-1} e^{\lambda})(q e^{-\lambda} - q^{-1} e^{\lambda})} = 1; \quad \frac{\mathbf{e}_{12}(\lambda) \mathbf{e}_{12}(\lambda^{-1})}{(w e^{-\lambda} - w^{-1} e^{\lambda})(w e^{-\lambda} - w^{-1} e^{\lambda})} = 1: \quad (4.47)$$

The unitarity condition for $k(\lambda)$ depends on the choice of the quotient: for $N_N^0(w; \lambda; q)$, we get $k_1(\lambda) k_1(\lambda^{-1}) = 1$, for $N_N^1(w; \lambda; q)$,

$$k_1(\lambda) k_1(\lambda^{-1}) = e^{2\lambda} + e^{-2\lambda} - w - w^{-1}; \quad (4.48)$$

and, finally, for $N_N^1(w; \lambda; q)$,

$$k_1(\lambda) k_1(\lambda + i\frac{\pi}{2}) = i e^{2\lambda} - e^{-2\lambda} - w + w^{-1}: \quad (4.49)$$

Proof. This proposition is proved by direct computation expressing the generators $\hat{r}_{i,i+1}(\lambda)$, $\mathbf{e}_{i,i+1}(\lambda)$ and $k_i(\lambda)$ thanks to (4.43)-(4.44) and using the defining exchange relations (4.31)-(4.42) of the algebras $N_N(w; \lambda; q)$, $N_N^0(w; \lambda; q)$ or $N_N^1(w; \lambda; q)$. For example, we get

$$\begin{aligned} \mathbf{e}_{12}(\lambda) \mathbf{e}_{12}(\lambda^{-1}) &= w^{-1} e^{-\lambda} - w e^{-\lambda^{-1}} - w^{-1} e^{-\lambda} - w e^{-\lambda^{-1}} \\ &= w^{-2} \frac{e^{2\lambda} - e^{-2\lambda}}{e^{\lambda} - e^{-\lambda}} + w^2 \frac{e^{2\lambda^{-1}} - e^{-2\lambda^{-1}}}{e^{\lambda^{-1}} - e^{-\lambda^{-1}}} \\ &= w^{-2} \frac{1 - e^{-4\lambda}}{1 - e^{-4\lambda}} + w^2 \frac{1 - e^{-4\lambda^{-1}}}{1 - e^{-4\lambda^{-1}}} + w^2 e^{2\lambda} - e^{2\lambda^{-1}} \\ &= w^{-2} + w^2 e^{2\lambda} - e^{2\lambda^{-1}}; \quad \text{using relation (4.31)}; \end{aligned}$$

which proves the second relation of (4.47). ■

The link between the generalized quantum Brauer algebra and the numerical previous solutions is given by the representation of $N_N^0(w; \lambda; q)$ or $N_N^1(w; \lambda; q)$. The map

$$i \mathbb{T} \hat{R}_{i,i+1} \quad , \quad i \mathbb{T} \hat{\mathbb{R}}_{i,i+1} \quad \text{and} \quad b_i \mathbb{T} V_i^t \quad \text{or} \quad i(G_q)_i V_i^t \quad (4.50)$$

is a representation of $N_N^0(q^{\frac{n}{2}}; (-1)^{n+1}; q)$ (we recall that $q^{\frac{n}{2}} = e^i$). It is proved by direct calculation. We then obtain the numerical matrices associated to the solutions (i) or (iv). Similarly, we can prove that

$${}_i \mathbb{T} \hat{R}_{i;i+1}, \quad {}_i \mathbb{T} \mathbb{K}_{i;i+1} \quad \text{and} \quad b_i \mathbb{T} \mathbb{K}_i \quad (\text{given after (4.30)}) \quad (4.51)$$

is a representation of $N_N^1(q^{\frac{n}{2}}; (-1)^{n+1}; q)$. In this case, we obtain the numerical solutions associated to solution (ii). Recall that we do not study the case (iii) since it provides a trivial finite algebra.

4.5 Quantum Brauer duality

We establish a result similar to the one of subsection 4.3 but now for any generic solution $K(\cdot)$ using the framework of the previous subsection.

We can define a finite algebra \mathbb{F}_f generated by $\mathbb{B}_0 = B_0 V_0^t$, which is obviously isomorphic to T_f . These generators can be represented by

$$\mathbb{B}_0^+ \mathbb{T} \mathbb{B}_0^+ = B_0^+ V_0^t = \hat{R}_{N_0} \hat{R}_{N-1;N} \cdots \hat{R}_{23} \hat{R}_{12} \mathbb{K}_1 \mathbb{K}_{12}^{-1} \mathbb{K}_{23}^{-1} \cdots \mathbb{K}_{N-1;N}^{-1} \mathbb{K}_N^{-1} \quad (4.52)$$

$$\mathbb{B}_0 \mathbb{T} \mathbb{B}_0 = B_0 V_0^t = \hat{R}_{N_0}^{-1} \hat{R}_{N-1;N}^{-1} \cdots \hat{R}_{23}^{-1} \hat{R}_{12}^{-1} \mathbb{K}_1^{-1} \mathbb{K}_{12} \mathbb{K}_{23} \cdots \mathbb{K}_{N-1;N} \mathbb{K}_N \quad (4.53)$$

where $\mathbb{K} = V^t$ for (i), $\mathbb{K} = iG_q V^t$ for (iv) and \mathbb{K} is given just after (4.30) for (ii). Recall that we do not investigate the case (iii) which gives a trivial finite algebra. Let us remark that with this form of the algebra, we succeed in writing the generators \mathbb{B} in terms of matrices appearing also in the representation (see (4.50) or (4.51)) of the algebra $N_N(w; \cdot; q)$.

The following proposition gives the analogue of the Brauer duality between the algebra generated by the elements of \mathbb{B}_0 and subalgebra of $N_N(q^{\frac{n}{2}}; \cdot; q)$:

Proposition 3 Let us introduce $g_1 = b_1^{-1} \frac{1}{q} \frac{1}{q^{-1}} b_1$. The actions of the algebra generated by $f_{i;j} \mid i, j \in N-1, g$ and g_1 (see (4.50)–(4.51)) and of the algebra \mathbb{F}_f on the space $(C^n)^{\otimes N}$ commute with each other.

Proof. The proof of this proposition is based in showing the following two relations, for $1 \leq i \leq N-1$,

$${}_i \mathbb{T} \hat{R}_{i;i+1} \mathbb{T} \mathbb{B}_0 = 0 \quad \text{and} \quad \mathbb{K}_1^{-1} \mathbb{K}_{12} \mathbb{K}_{12}^{-1} \mathbb{K}_1 \mathbb{T} \mathbb{B}_0 = 0 \quad (4.54)$$

We shall use the fact that all the matrices involved in this computation satisfy the exchange relations of $N_N(w; \cdot; q)$. Let us give explicitly the proof for $[\mathbb{K}_1^{-1} \mathbb{K}_{12} \mathbb{K}_{12}^{-1} \mathbb{K}_1 \mathbb{T} \mathbb{B}_0] = 0$. Since \mathbb{K}_1 and \mathbb{K}_{12} commute with matrices in space $0; 3; 4; \dots; N$, we can focus on proving

$${}_i \mathbb{T} \mathbb{K}_1^{-1} \mathbb{K}_{12} \mathbb{K}_{12}^{-1} \mathbb{K}_1 \hat{R}_{23} \hat{R}_{12} \mathbb{K}_1 \mathbb{K}_{12}^{-1} \mathbb{K}_{23}^{-1} = 0 : \quad (4.55)$$

Let us first compute

$$\begin{aligned}
\mathbb{R}_1^{-1} \mathbb{R}_{12} (\mathbb{R}_1 \hat{\mathbb{R}}_{23}) \hat{\mathbb{R}}_{12} \mathbb{R}_1 \mathbb{R}_{12}^{-1} \mathbb{R}_{23}^{-1} &= \mathbb{R}_1^{-1} \mathbb{R}_{12} \hat{\mathbb{R}}_{23} (\mathbb{R}_1 \hat{\mathbb{R}}_{12} \mathbb{R}_1 \mathbb{R}_{12}^{-1}) \mathbb{R}_{23}^{-1} \\
&= \mathbb{R}_1^{-1} (\mathbb{R}_{12} \hat{\mathbb{R}}_{23} \mathbb{R}_{12}^{-1}) \mathbb{R}_1 \hat{\mathbb{R}}_{12} \mathbb{R}_{23}^{-1} \mathbb{R}_1 \\
&= \mathbb{R}_{23}^{-1} \mathbb{R}_1^{-1} \hat{\mathbb{R}}_{12} \mathbb{R}_1 (\mathbb{R}_{23} \hat{\mathbb{R}}_{12} \mathbb{R}_{23}^{-1}) \mathbb{R}_1 \\
&= \mathbb{R}_{23}^{-1} (\mathbb{R}_1^{-1} \hat{\mathbb{R}}_{12} \mathbb{R}_1 \mathbb{R}_{12}^{-1}) \hat{\mathbb{R}}_{23} \mathbb{R}_{12} \mathbb{R}_1 \\
&= \mathbb{R}_{23}^{-1} \mathbb{R}_{12}^{-1} \mathbb{R}_1 \hat{\mathbb{R}}_{12} \hat{\mathbb{R}}_{23} \mathbb{R}_1^{-1} \mathbb{R}_{12} \mathbb{R}_1 \quad (4.56)
\end{aligned}$$

In the latter relations, the parenthesis indicate which part of the product is modified using different relations (4.31)–(4.41). We obtained similar result for $\mathbb{R}_1^{-1} \mathbb{R}_{12}^{-1} \mathbb{R}_1$. Then, we have proved that

$$\mathbb{R}_1^{-1} \mathbb{R}_{12}^{-1} \mathbb{R}_1 \mathbb{B}_0^+ = \mathbb{C}_0^+ \mathbb{R}_1^{-1} \mathbb{R}_{12}^{-1} \mathbb{R}_1; \quad (4.57)$$

where $\mathbb{C}_0^+ = \hat{\mathbb{R}}_{N0} \hat{\mathbb{R}}_{N-1N} \dots \hat{\mathbb{R}}_{34} \mathbb{R}_{23}^{-1} \mathbb{R}_{12}^{-1} \mathbb{R}_1 \hat{\mathbb{R}}_{12} \hat{\mathbb{R}}_{23} \mathbb{R}_{34}^{-1} \dots \mathbb{R}_N^{-1} \mathbb{R}_{N0}$. Multiplying on the left by $\mathbb{R}_1^{-1} \mathbb{R}_{12} \mathbb{R}_1$ the previous relation and knowing that it satisfies relation (4.31), we finish the proof. ■

Within this framework, the proof of this proposition requires only the exchange relations of the algebra $\mathcal{N}(w; ;q)$ whereas, in the previous case (see subsection 4.3), the proof of the analogous proposition is based on direct computation.

Acknowledgments: One of the authors (N.C.) wishes to thank the Physics Department of the University of Bologna and the INFN section of Bologna, where this work was initiated. This work was supported by INFN and the European Network 'EUCLID'; Integrable models and applications: from strings to condensed matter', contract number HPRN-CT-2002-00325.

A Appendix

In this appendix, we give the explicit expression of the commutation relations between the finite algebra and the quantum twisted Yangian for $U_q(\mathfrak{gl}_3)$ and for $K(\cdot) = I$. Let us express B as a matrix

$$B(\cdot) = \begin{pmatrix} 0 & & & 1 \\ A_1 & D_1 & D & \\ \mathbb{C} & C_1 & A_2 & D_2 & A \\ & C & C_2 & A_3 & \end{pmatrix} \quad (A.1)$$

Set also $c = q^{-1} q^1$ and $\mathbb{K}; Y_{\mathfrak{q}} = q X Y - q^{-1} Y X$. We recall that the diagonal charges are trivial B_{ii}^+ / I so we simply deal with the rest of them. Then from (4.16) and bearing in mind (A.1) various sets of equations are inferred:

$$\begin{aligned}
\frac{B_{12}^+; A_1}{h} &= q c C_1 + c D_1; & \frac{B_{12}^+; A_2}{h} &= q c C_1 - c D_1; & B_{12}^+; A_3 &= 0; \\
\frac{q B_{12}^+; D_1}{h} &= \frac{B_{12}^+; C_1}{h} = q c (A_2 - A_1); & \frac{B_{12}^+; D_2}{h} &= q^{\frac{1}{2}} c D \\
\frac{B_{12}^+; C_2}{h} &= q^{\frac{1}{2}} c C_1; & \frac{B_{11}^+; D}{h} &= q^{\frac{1}{2}} c D_2; & \frac{B_{12}^+; C}{h} &= q^{\frac{1}{2}} c C_2 \quad (A.2)
\end{aligned}$$

$$\begin{aligned}
\langle B_{23}^+; A_1 \rangle &= 0; & \langle B_{23}^+; A_2 \rangle &= q c C_2 + c D_2; & \langle B_{23}^+; A_3 \rangle &= q c C_2 - c D_2 \\
\langle q B_{22}^+; D_2 \rangle &= \langle B_{23}^+; C_2 \rangle = q c (A_3 - A_2); & \langle B_{23}^+; D_1 \rangle &= q^{\frac{1}{2}} c D \\
\langle B_{23}^+; C_1 \rangle &= q^{\frac{1}{2}} c C; & \langle B_{23}^+; D_{q^{\frac{1}{2}}} \rangle &= q^{\frac{1}{2}} c D_1; & \langle B_{23}^+; C_{q^{\frac{1}{2}}} \rangle &= q^{\frac{1}{2}} c C_1 \quad (A.3)
\end{aligned}$$

$$\begin{aligned}
\langle B_{13}^+; A_1 \rangle &= q c C + c D; & \langle B_{13}^+; A_3 \rangle &= q c C - c D; \\
\langle q B_{13}^+; D \rangle &= \langle B_{13}^+; C \rangle = q c (A_3 - A_1); & \langle B_{13}^+; D_1 \rangle &= c (q^{\frac{1}{2}} C_2 + q^{\frac{1}{2}} D B_{12}^+ - q^{\frac{1}{2}} A_1 B_{23}^+) \\
\langle B_{13}^+; C_1 \rangle &= q^{\frac{1}{2}} c (D_2 + B_{12}^+ C - B_{23}^+ A_1); & \langle B_{13}^+; D_2 \rangle &= c (q^{\frac{1}{2}} C_1 + q^{\frac{1}{2}} B_{12}^+ A_3 - q^{\frac{1}{2}} B_{23}^+ D); \\
\langle B_{13}^+; C_2 \rangle &= q^{\frac{1}{2}} c (A_3 B_{12} - C B_{23} - D_1); & \langle B_{13}^+; A_2 \rangle &= c (B_{23}^+ D_1 - C_1 B_{23}^+ + D_2 B_{12}^+ + B_{12}^+ C_2);
\end{aligned}$$

References

- [1] I.V. Cherednik, Factorizing particles on a half line and root system s , Theor. Math. Phys. 61 (1984) 977.
- [2] E.K. Sklyanin, Boundary conditions for integrable quantum systems, J. Phys. A 21 (1988) 2375.
- [3] G.I. Olshanski, Twisted Yangians and infinite-dimensional classical Lie algebras in Quantum Groups' (P.P. Kulish, Ed.), Lecture notes in Math. 1510, Springer (1992) 103; A.M.olev, M.Nazarov and G.I.Olshanski, Yangians and classical Lie algebras, Russ. Math. Surveys 51 (1996) 206 and hep-th/9409025.
- [4] A.I.Molev, E.Ragoucy and P.Sorba, Coideal subalgebras of quantum algebra, Rev. Math. Phys. 15 (2003) 789 and math.QA/0208140.
- [5] P. Bowcock, E. Corrigan, P.E. Dorey and R.H. Rietdijk, Classically integrable boundary conditions for affine Toda field theories, Nucl. Phys. B 445 (1995) 469 and hep-th/9501098; P. Bowcock, E. Corrigan and R.H. Rietdijk, Background field boundary conditions for affine Toda field theories, Nucl. Phys. B 465 (1996) 350 and hep-th/9510071.
- [6] G.M. Gaudinberger, New non-diagonal solutions to the $A_n^{(1)}$ boundary Yang-Baxter equation, hep-th/9911178.
- [7] A. Dolkou, Quantum spin chain with 'soliton nonpreserving' boundary conditions, J. Phys. A 33 (2000) 8797 and hep-th/0006197.

- [8] D. A. Maudon, J. Avan, N. Crampé, A. D. Dikou, L. Frappat, E. Ragoucy, General boundary conditions for the $sl(N)$ and $sl(M|N)$ open spin chains, *JSTAT* 08 (2004) P005, math-ph/0406021.
- [9] D. A. Maudon, N. Crampé, A. D. Dikou, L. Frappat and E. Ragoucy, Analytical Bethe Ansatz for open spin chains with soliton non preserving boundary conditions, *Int. J. Math. Phys.* A 21 (2006) 1537 and math-ph/0503014.
- [10] R. Brauer, On algebras which are connected with the semisimple continuous groups, *Ann. Math.* 38 (1937) 857.
- [11] A. I. Molev, A new quantum analog of the Brauer algebra, *Czech. J. of Physics* 53 (2003) 1073 and math-QA/0211082.
- [12] L. D. Faddeev, N. Yu. Reshetikhin and L. A. Takhtajan, Quantization of Lie groups and Lie algebras, *Leningrad Math. J.* 1 (1990) 193.
- [13] R. J. Baxter, Partition function of the eight-vertex lattice model, *Ann. Phys.* 70 (1972) 193;
J. Stat. Phys. 8 (1973) 25;
Exactly solved models in statistical mechanics (Academic Press, 1982).
- [14] A. B. Zamolodchikov and A. L. B. Zamolodchikov, Factorized S-matrices in two dimensions as the exact solutions of certain relativistic quantum field theory models, *Annals Phys.* 120 (1979) 253.
L. D. Faddeev, *Sov. Sci. Rev. C* 1 (1980) 107.
- [15] V. E. Korepin, New effects in the massive Thirring model: repulsive case, *Comm. Math. Phys.* 76 (1980) 165;
V. E. Korepin, G. Izergin and N. M. Bogoliubov, *Quantum inverse scattering method, correlation functions and algebraic Bethe Ansatz* (Cambridge University Press, 1993).
- [16] V. G. Drinfeld, Hopf algebras and the quantum Yang-Baxter equation, *Soviet. Math. Dokl.* 32 (1985) 254;
A new realization of Yangians and quantized affine algebras, *Soviet. Math. Dokl.* 36 (1988) 212.
- [17] V. G. Drinfeld, *Proceedings of the 1986 International Congress of Mathematicians, Berkeley ed. A. M. Gleason 1986* (Providence, RI: American Physical Society) pp 798.
- [18] G. Delius and N. M. Mackay, Quantum group symmetry in sine-Gordon and affine Toda field theories on the half-line, *Commun. Math. Phys.* 233 (2003) 173 and hep-th/0112023.
- [19] G. W. Delius, N. J. Mackay and B. J. Short, Boundary remnant of Yangian symmetry and the structure of rational reflection matrices, *Phys. Lett. B* 522 (2001) 335 and

- hep-th/0109115; Erratum *ibid.* B 524 (2002) 401;
 N.J.Mackay, Rational K-matrices and representations of twisted Yangians, *J. Phys. A* 35 (2002) 7865 and math.QA/0205155.
- [20] M.Mintchev, E.Ragoucy and P.Sorba, Spontaneous symmetry breaking in the $gl(N)$ -NLS hierarchy on the half line, *J. Phys. A* 34 (2001) 8345 and hep-th/0104079.
- [21] A.Doikou, On reflection algebras and twisted Yangians, *J. Math. Phys.* 46, 053504 (2005) and hep-th/0403277.
- [22] A.Doikou and R.J.Nepomechie, Duality and quantum algebra symmetry of the $A_N^{(1)}$ open spin chain with diagonal boundary fields, *Nucl. Phys. B* 530 (1998) 641 and hep-th/9807065.
- [23] N.Crampe, Approches algébriques dans les systèmes intégrables, University of Savoie PhD Thesis (June 2004), in French, Preprint LAPTH-These-1056/04, <http://wwwlapp.in2p3.fr/preplapp/psth/LAPTH1056.ps.gz>.
- [24] W.Gallas and M.J.Martins, Solution of the $SU(N)$ Vertex Model with Non-Diagonal Open Boundaries, *Phys. Lett. A* 335 (2005) 167 and nlin.SI/0407027.
- [25] D.Amaudon, N.Crampe, A.Doikou, L.Frappat, E.Ragoucy, Analytical Bethe Ansatz for closed and open $gl(n)$ -spin chains in any representation, *JSTAT* 02 (2005) P02007 and math-ph/0411021.
- [26] A.Doikou, From affine Hecke algebras to boundary symmetries, *Nucl. Phys. B* 725 (2005) 493 and math-ph/0409060.
- [27] M.Jimbo, A q -difference analog of $U(G)$ and the Yang-Baxter equation, *Lett. Math. Phys.* 10 (1985) 63.
- [28] L.D.Faddeev, N.Yu.Reshetikhin and L.A.Takhtajan, Quantization of Lie groups and Lie algebras, *Leningrad Math. J.* 1 (1990) 193.
- [29] D.Amaudon, N.Crampe, A.Doikou, L.Frappat, E.Ragoucy, Spectrum and Bethe ansatz equations for the $U_q(gl(N))$ closed and open spin chains in any representation, *Ann. Henri Poincaré* 7 (2006) 1217 and math-ph/0512037.
- [30] A.Doikou, A symmetric twin representation: the transfer matrix symmetry, *SIGMA* 3 (2007) 009 and math-ph/0606040.
- [31] A.M.Gavrilik and A.J.Klimyk, q -deformed orthogonal and pseudo-orthogonal algebras and their representations, *Lett. Math. Phys.* 21 (1991) 215.
- [32] M.Noumi, Macdonald's symmetric polynomials as zonal spherical functions on quantum homogeneous spaces, *Adv. Math.* 123 (1996) 16 and math.QA/9503224.