

PARTIAL REDUCTIONS OF HAMILTONIAN FLOWS AND
HESS-APPEL'ROT SYSTEMS ON $SO(n)$

BOZIDAR JOVANOVIĆ

Mathematical Institute SANU
Kneza Mihaila 35, 11000 Belgrade, Serbia
e-mail: bozaj@mi.sanu.ac.yu

Abstract. We study reductions of the Hamiltonian flows restricted to their invariant submanifolds. As examples, we consider partial Lagrange-Routh reductions of the natural mechanical systems such as geodesic flows on compact Lie groups and n -dimensional variants of the classical Hess-Appel'rot case of a heavy rigid body motion about a fixed point.
MSC : 37J35, 37J15, 53D20

Contents

1. Introduction	1
1.1. Hess-Appel'rot System	1
1.2. Reduction of Symmetries	3
1.3. Outline and Results of the Paper	3
2. Partial Reductions and Integrability	4
3. Partial Lagrange-Routh Reductions	7
3.1. Geodesic Flows on Compact Lie Groups	8
3.2. Local Description	10
4. n -Dimensional Hess-Appel'rot System	11
4.1. Invariant Relations	12
4.2. Right Trivialization and Reduction to S^{n-1}	12
4.3. Reduction to $(\mathfrak{so}(n), \mathbb{R}^n)$	14
4.4. L-A Pair	16
5. Hess-Appel'rot system on $(\mathfrak{so}(4), \mathfrak{so}(4))$	17
Acknowledgments	19
References	19

1. Introduction

In this paper we study reductions of the Hamiltonian flows restricted to their invariant submanifolds. Apparently, the lowering of order in Hamiltonian systems having invariant relations was firstly studied by Levi-Civita (e.g., see [17], ch. X).

1.1. Hess-Appel'rot System. The classical example of the system having an invariant relation is a celebrated Hess-Appel'rot case of a heavy rigid body motion

[15, 1]. Recall that the motion of a heavy rigid body around a fixed point, in the moving frame e , is represented by the Euler-Poisson equations

$$(1) \quad \frac{d}{dt} \mathfrak{m} = \mathfrak{m} \times \omega + GM \sim \times \mathfrak{r}; \quad \frac{d}{dt} \sim = \sim \times \omega; \quad \dot{\omega} = A \mathfrak{m};$$

where ω is the angular velocity, \mathfrak{m} the angular momentum, $I = A^{-1}$ the inertial tensor, M mass and \mathfrak{r} the vector of the mass center of a rigid body; \sim is the direction of the homogeneous gravitational field and G is the gravitational constant.

The equations (1) always have three integrals, the energy, geometric integral and the projection of angular momentum:

$$(2) \quad F_1 = \frac{1}{2} (\mathfrak{m}; \dot{\omega}) + MG (\mathfrak{r}; \sim); \quad F_2 = (\sim; \sim) = 1; \quad F_3 = (\mathfrak{m}; \sim):$$

For the integrability we need a fourth integral. There are three famous integrable cases: Euler, Lagrange and Kowalevskaya [2, 14].

A part of these cases, there are various particular solutions (e.g., see [14]). The celebrated is partially integrable Hess-Appell'rot case [15, 1]. Under the conditions:

$$(3) \quad r_2 = 0; \quad r_1 \frac{p}{a_3} - \frac{p}{a_2} - r_3 \frac{p}{a_2} - \frac{p}{a_1} = 0; \quad A = \text{diag}(a_1; a_2; a_3);$$

$a_3 > a_2 > a_1 > 0$, system (1) has an invariant relation given by

$$(4) \quad F_4 = (\mathfrak{m}; \mathfrak{r}) = r_1 m_1 + r_3 m_3 = 0:$$

The system is integrable up to one quadrature: the compact connected components of the regular invariant sets $F_1 = c_1; F_2 = 1; F_3 = c_3; F_4 = 0$ are tori, but not with quasi-periodic dynamics. The classical and algebro-geometric integration can be found in [14] and [9], respectively.

There is a nice geometrical interpretation of the conditions (3): the intersection of the plane orthogonal to \mathfrak{r} with the ellipsoid $(A \mathfrak{m}; \mathfrak{m}) = \text{const}$ is a circle (e.g., see [8]). If instead of the moving base given by the main axes of the inertia, we take the moving base $f_1; f_2; f_3$ such that the mass center of a rigid body \mathfrak{r} is proportional to f_3 ,

$$(5) \quad \mathfrak{r} = f_3; \quad \mathfrak{r} = \frac{q}{r_1^2 + r_3^2} f_3;$$

then the inverse of the inertial operator reads

$$(6) \quad A = \begin{pmatrix} 0 & & & 1 \\ & a_2 & 0 & a_{13} \\ & 0 & a_2 & a_{23} \\ & a_{13} & a_{23} & a_{33} \end{pmatrix} A$$

and the invariant relation (4) is simply given by

$$(7) \quad m_3 = 0:$$

Historical overview, with an application of Levi-Civita ideas to the Hess-Appell'rot system and other classical rigid body problems can be found in Borisov and Mamaev [8].

1.2. Reduction of Symmetries. Let G be a Lie group with a free proper Hamiltonian action on a symplectic manifold $(M; \omega)$. Let

$$(8) \quad \mu : M \rightarrow \mathfrak{g}^*$$

be the corresponding equivariant momentum map. Assume that λ is a regular value of μ , so that

$$(9) \quad M_\lambda = \mu^{-1}(\lambda)$$

and $M_0 = \mu^{-1}(0)$ are smooth manifolds. Here $O = G \cdot G$ is the coadjoint orbit of λ . The manifolds M_λ and M_0 are G -invariant and G -invariant, respectively.

There is a unique symplectic structure ω_λ on $N = M_\lambda/G$ satisfying

$$(10) \quad \pi^* \omega_\lambda = \omega - \langle d\mu, \xi \rangle$$

where $\pi : M \rightarrow N$ is the natural projection [20].

According to Noether's theorem, if h is a G -invariant function then the momentum mapping μ is an integral of the Hamiltonian system

$$(11) \quad \dot{x} = X_h$$

In addition, its restriction to the invariant submanifold M_λ projects to the Hamiltonian system $\dot{y} = X_H$ on the reduced space N with H defined by [20]

$$(12) \quad h(y) = H(\lambda)$$

An alternative description of the reduced space is as follows. Let $f; g$ be the canonical Poisson bracket on $(M; \omega)$. Then the manifold M_λ/G carries the induced Poisson structure and M_0/G is the symplectic leaf in M_λ/G . The mapping

$$(13) \quad \pi : N \rightarrow M_0/G$$

which assigns to the G -orbit of $x \in M_\lambda$ the G -orbit through x in M_0 establishes the symplectomorphism between N and M_0/G (e.g., see [24]).

1.3. Outline and Results of the Paper. The aim of this paper is to study Hamiltonian systems that naturally generalize geometrical properties of the Hess-Appelrot system (Definition 3, Section 4). This is the reason we are interested in the following: suppose that h is not a G -invariant function, but (9) is still an invariant manifold of the Hamiltonian system (11). As a modification of the regular Marsden-Weinstein reduction, we study (partial) reduction of the Hamiltonian system (11) from M to N and relationship between the integrability of the reduced and nonreduced system (Theorems 1, 2, Section 2).

In Section 3 we define partial Lagrange-Routh reductions (Theorem 3). The construction of the geodesic flows on compact Lie groups, having invariant manifolds foliated by invariant tori is presented. However, the flows over invariant tori are not quasi-periodic.

In Section 4 we consider an n -dimensional heavy rigid body motion about a fixed point. Recall that there are two natural multidimensional variants of a heavy rigid body which are described by the Euler-Poisson equations on the dual spaces of semidirect products $\mathfrak{so}(n) \ltimes \mathfrak{so}(n)$ and $\mathfrak{so}(n) \ltimes \mathbb{R}^n$. The integrable generalizations of the Lagrange top with respect to the first and the second n -dimensional variant are given in [25, 10] and [3], respectively. The n -dimensional Kowalevskaya top in the presence of q gravitational fields ($q = n$) represented by Euler-Poisson equations on the dual space of the semidirect product $\mathfrak{so}(n) \ltimes (\mathbb{R}^n)^q$ is given in [26].

Recently, an interesting construction of the Hess-Appel'rot system on $\mathfrak{so}(n)$ is studied by Dragović and Gajić [1]. Besides, by generalizing analytical and algebraic properties of the classical system (1), (3), they introduced a class of systems with invariant relations, so called Hess-Appel'rot type systems with remarkable property: there exist a pair of compatible Poisson structures, such that the system is Hamiltonian with respect to the first structure and invariant relations are Casimir functions with respect to the second structure (for more details see [1]).

We consider the another generalization of a heavy rigid body and define the n -dimensional Hess-Appel'rot system within the framework of partial reductions (Lemma 6). It is known that the motion of a rigid body mass center in the classical Hess-Appel'rot system is described by the spherical pendulum equations [29, 8]. We shall prove the same statement for the n -dimensional variant of the system (Theorem 7). On the other side, the L-A pair of the system written in the closed form of the Euler-Poisson equations on the dual space of the semidirect product $\mathfrak{so}(n) \ltimes \mathbb{R}^n$ is given. In particular, we get the L-A pair representation for the Euler-Poisson equations of the classical Hess-Appel'rot system (1), (5), (6) different from those given in [9] (Theorem 9).

Finally, we have made a relationship between our construction and the Dragović-Gajić $\mathfrak{so}(4) \ltimes \mathfrak{so}(4)$ system (Theorem 10, Section 5).

2. Partial Reductions and Integrability

Let G be a compact connected Lie group with a free Hamiltonian action on a symplectic manifold (M, ω) with the momentum map (8). Assume that $\mu^{-1}(c)$ is a regular value of μ .

Theorem 1. (i) Suppose that the restriction of h to $M_0 = \mu^{-1}(c)$ is a G -invariant function. Then M_0 is an invariant manifold of the Hamiltonian system (11) and $X_{h|_{M_0}}$ projects to the Hamiltonian vector field X_H

$$(14) \quad d(X_{h|_{M_0}})_{\mu^{-1}(c)} = X_H|_{\mu^{-1}(c)}$$

where H is the induced function on N defined by (12).

(ii) The inverse statement also holds: if (9) is an invariant submanifold of the Hamiltonian system (11) then the restriction of h to M_0 is a G -invariant function and $X_{h|_{M_0}}$ projects to the Hamiltonian vector field X_H on N , where H is defined by (12).

In both cases, the Hamiltonian vector field $X_{h|_{M_0}}$ is not assumed to be G -invariant on M_0 . Moreover $X_{h|_{M_0}}$ may not be G -invariant as well. It is invariant modulo the kernel of $d\mu$, which is sufficient the tools of symplectic reduction are still applicable.

Definition 1. We shall refer to the passing from $\underline{x} = X_{h|_{M_0}}$ to

$$(15) \quad \underline{y} = X_H$$

as a partial reduction.

If h is G -invariant, the complete integrability of the reduced and original system are closely related (see [16, 28]). In our case, we have the following corollary.

Theorem 2. (i) Suppose that the partially reduced system (15) is completely integrable, with a complete set of commuting integrals F_1, \dots, F_m , $m = \frac{1}{2} \dim N$.

Let $f_1 = F_1, \dots, f_n = F_n$. Then M is almost everywhere foliated by $(m + \dim G)$ -dimensional invariant isotropic manifolds

$$(16) \quad M_c = \{f_1 = c_1, \dots, f_m = c_m\} \subset g$$

of the system (11).

(ii) In addition, if μ is a regular element of g and f_1, \dots, f_m can be extended to the commuting G -invariant functions in some neighborhood of M_c , then the compact connected components of invariant manifolds (16) are tori. However, in general, the flow over the tori is not quasi-periodic.

Remark 1. If the partially reduced system (15) is integrable then we need $(\dim G)$ -additional quadratures for the solving of $\dot{x} = X_{h|_M}$ (the reconstruction equations). Also, if $G = G$, i.e., $O = f|_g$, then $m + \dim G = \frac{1}{2} \dim M$. Hence, the invariant manifolds (16) are Lagrangian.

The partial reduction can be seen as a special case of the symplectic reductions studied in [4, 19] (see also [18], Ch. III). There, the symplectic reduction of a symplectic manifold $(M; \omega)$ is any surjective submersion $p: N \rightarrow P$ of a submanifold $N \subset M$ onto another symplectic manifold $(P; \omega_P)$, which satisfies $p^*\omega_P = \omega|_N$. Also, instead of a compact group action one can consider a proper group action. However we work in the framework of a regular Marsden-Weinstein reduction and a compact group action, which allows us to easily describe the partial reductions of natural mechanical systems considered in this paper.

Proof of Theorem 1. (i) Let μ_1, \dots, μ_n be the base of g , $(\mu_i; \cdot) = \mu_i$ and

$$(17) \quad \mu = (\mu_i; \cdot) : M \rightarrow \mathbb{R}^n; \quad \mu_i = 1, \dots, n:$$

Then the level set of the momentum mapping (9) is given by the equations

$$\mu_i = 1; \quad \mu_n = 1, \dots, n:$$

The action of G is generated by Hamiltonian vector fields X_{μ_i} . Since $h|_{M_0}$ is G -invariant, for $x \in M_0$ we have

$$(18) \quad (dh; X_{\mu_i}) = f\mu_i; \quad g = (d\mu_i; X_{\mu_i}) = 0; \quad \mu_i = 1, \dots, n:$$

Thus M_0 is an invariant submanifold.

Let h be an arbitrary G -invariant function that coincides with h on M_0 . Then X_h is a G -invariant vector field on M which projects to X_H [20]:

$$(19) \quad d(X_h)|_x = X_H|_{\mu(x)}:$$

Let $\mu = h|_g$. From the condition $\dot{\mu} = 0$ we can express μ , in an open neighborhood of M_0 , as

$$\mu(x) = \sum_{i=1}^n \mu_i(x) (\mu_i(x) = 1):$$

Now, let f be a G -invariant function on M . Since $f|_g = 0$ (Noether's theorem), we get

$$f; fg|_x = \sum_{i=1}^n f; fg(\mu_i) + \sum_{i=1}^n f; fg|_{\mu_i} = 0:$$

Thus the Poisson bracket $f;fg_j = (df;X_j)_j$ vanish for an arbitrary G -invariant function f . In other words

$$(20) \quad X_j \in T_x(G \backslash M) \setminus \mathbb{T}M :$$

Combining (19), $X_h = X_h + X$ and (20) with the wellknown identity (e.g, see [24])

$$T_x(G \backslash M) \setminus \mathbb{T}M = T_x(G \backslash M) = \ker d_j ;$$

we prove the relation (14).

(ii) If G is a connected compact group, the coadjoint isotropy group G is connected. Thus, the function h is G -invariant if and only if it is invariant with respect to the infinitesimal action of G .

Suppose that M is an invariant submanifold of (11). Then (18) holds for $x \in M$. Since the action of G is generated by Hamiltonian vector fields X , from (18) we get that h is invariant with respect to the infinitesimal action of G . Therefore we have welldefined reduced Hamiltonian function H on N .

By using the diffeomorphism (13) and the fact that M_0 is a closed submanifold of M , we can find a G -invariant function h on M which coincides with h on M_0 . Now, the relation (14) follows from the proof of item (i).

Proof of Theorem 2. (i) Consider a regular invariant Lagrangian submanifold

$$(21) \quad N_c = \{F_1 = c_1; \dots; F_m = c_m\} \subset N ;$$

of the partially reduced system (15). From the relations (10) and (14) we get that $M_c = \pi^{-1}(N_c)$ is an invariant isotropic manifold of the system (11).

(ii) Let g be a regular element (G/T^r is a maximal torus of G , $r = \text{rank } G$). If the connected component N_c° of (21) is compact then, by Liouville's theorem, it is diffeomorphic to a m -dimensional torus T^m with quasi-periodic flow of (15). Thus the compact connected component $M_c^\circ = \pi^{-1}(N_c^\circ)$ of (16) is a torus bundle over T^m :

$$(22) \quad \begin{array}{c} T^r \rightarrow M_c^\circ \\ \# \\ T^m \end{array}$$

Let $I_1; \dots; I_r$ be the basic A_{dg} -invariant polynomials on g and $P_1; \dots; P_{n-r}$ be linear functions on g such that P_k, I are independent at g . Also, let $i = I$ and $p_k = P_k$ be the pullbacks of I and P_k by the momentum mapping, $i = 1; \dots; r, k = 1; \dots; n-r$.

Suppose that $f_1; \dots; f_m$ can be extended to commuting G -invariant functions in some G -invariant neighborhood V of M_c° . Then, within V , M_c° is given by the equations

$$f_1 = c_1; \dots; f_m = c_m; i_1 = I_1(g); \dots; i_r = I_r(g); p_1 = P_1(g); \dots; p_{n-r} = P_{n-r}(g) :$$

From the Noether theorem the functions i, p_k commute with all G -invariant functions on M and, since i are G -invariant, the following commuting relations hold on V :

$$(23) \quad \begin{aligned} ff_a; f_b g = ff_a; i g = fi; i g = ff_a; p_k g = fi; p_k g = 0; \\ a; b = 1; \dots; m; i = 1; \dots; r; k = 1; \dots; n-r : \end{aligned}$$

Now, as in the case of non-commutative integrability of Hamiltonian systems [13, 23], M_c° is a torus with tangent space spanned by X_{f_a}, X_{i_1} . Namely, from the

vanishing of the Poisson brackets (23) we get that the vector fields X_{f_a}, X_{f_i} are tangential to M^c . Since they are independent, from the dimensional reasons, they span the tangent spaces $T_x M^c, x \in M^c$. Taking into account (23) and the relations $\{X_{f_i}, X_{f_j}\} = f_i f_j - f_j f_i, [X_{f_i}, X_{f_j}] = X_{f_i f_j - f_j f_i}, f_i, f_j \in C^1(M)$, we obtain that M^c is isotropic. Furthermore the vector fields X_{f_a}, X_{f_i} commute between them selves. Since M^c is a compact manifold admitting $m+r = \dim M^c$ independent commuting vector fields it is a $(m+r)$ -dimensional torus, i.e., the bundle (22) is trivial.

In general, the flow of $\underline{x} = X_h$ over the torus M^c is not quasi-periodic: the vector field X_h do not commute with vector fields $X_{f_1}, \dots, X_{f_m}; X_{f_{i_1}}, \dots, X_{f_{i_r}}$ (although Poisson brackets $f_h; f_a, f_h; f_i$ vanish on M^c).

Remark 2. Let $x \in M^o$ and $z = \pi(x)$, where $\pi: M \rightarrow M/G$ is the canonical projection. By the use of (13) and the existence of local canonical coordinates on the Poisson manifold M/G within some neighborhood U of z (e.g., see [18]), the functions f_i can be always extended to G -invariant commuting functions in the G -invariant neighborhood $V = \pi^{-1}(U)$. Thus, if M^o is "small enough", it is a torus.

Remark 3. It is clear that complete commutative integrability of the reduced system in Theorem 2, can be replaced by the condition of non-commutative integrability.

Remark 4. The Hamiltonian flow (11) preserves the canonical measure $\mu = \int_{\dim M=2}$ (the Liouville theorem). If M^o is an invariant manifold of the equations (11), then the functions (17) are particular integrals. Therefore the time derivative of μ is of the form

$$-\dot{\mu}(x) = \int_M \text{tr} \left(\frac{dX}{dt} \right) \mu(x);$$

where μ are smooth functions. Let tr be the trace of the matrix $\frac{dX}{dt}$. After straightforward calculations, one can prove that the flow (11) restricted to the invariant manifold M^o preserve the restriction of μ to M^o if and only if

$$\text{tr} \left(\frac{dX}{dt} \right) = 0;$$

for $x \in M^o$. In particular, if the Hamiltonian h is a G -invariant function, then $\int_{M^o} \mu$ is an invariant volume form.

3. Partial Lagrange-Routh Reductions

Let $(Q; l)$ be a natural mechanical system with Lagrangian $l = \frac{1}{2} \langle q; q \rangle + v(q)$, where the metric $\langle \cdot; \cdot \rangle$ is also regarded as a mapping $\langle \cdot; \cdot \rangle: TQ \rightarrow T^*Q$. The motion of the system is described by the Euler-Lagrange equations

$$(24) \quad \frac{\partial l}{\partial q} - \frac{d}{dt} \frac{\partial l}{\partial \dot{q}} = 0$$

or by the Hamiltonian equations on the cotangent bundle T^*Q with the Hamiltonian $h(q; p) = \frac{1}{2} \langle p; p \rangle + v(q)$ being the Legendre transformation of l .

Let G be a compact connected Lie group acting freely on Q and $\pi: Q \rightarrow B = Q/G$ be the canonical projection. The G -action can be naturally extended to the Hamiltonian action on $T^*Q: g \cdot (q; p) = (g \cdot q; (dg)^{-1} p)$ with the momentum mapping given by (e.g., see [18])

$$\langle (q; p); \xi \rangle = \langle p; \xi \rangle; \quad \xi \in \mathfrak{g}$$

Here ξ is the vector given by the action of one-parameter subgroup $\exp(t\xi)$ at q .

For Lagrangian system s , it is convenient to work with tangent bundle reductions. Let $V_q = \ker \tau_q$ be the tangent space to the fiber G at q (vertical space at q) and $V = \ker \tau$ be the vertical distribution. Then

$$(TQ)_0 = \tau^{-1}(0) = \ker V_q; \quad \ker V_q = \{p \in T_q Q \mid \tau(p) = q\} = 0; \quad \tau^{-1}(0) = 0;$$

Consider the horizontal distribution $H = \ker V_q$ orthogonal to V with respect to the metric g . Equivalently, H is the zero level-set of the tangent bundle momentum mapping μ :

$$(25) \quad H = \tau^{-1}(0); \quad \mu(q; \dot{q}) = \frac{\partial l}{\partial \dot{q}}; \quad \mu_q = \left(\frac{\partial l}{\partial \dot{q}}; \dot{q} \right); \quad \tau^{-1}(0) = 0;$$

Since $\mu_q(H_q) = \ker V_q$, we see that H is invariant with respect to the "twisted" G -action

$$(26) \quad g(q; X) = (g(q; g_q^{-1}(d\tau^{-1}(X)))); \quad X \in T_q Q;$$

that is the pullback of canonical symplectic G -action on TQ via metric g :

$$\begin{array}{ccc} TQ & \xrightarrow{g} & TQ \\ \downarrow \tau & \# & \downarrow \tau \\ TQ & \xrightarrow{g} & TQ \end{array}$$

From Theorem 1 we obtain

Theorem 3. (i) The horizontal distribution (25) is an invariant submanifold of the Euler-Lagrange equations (24) if and only if the potential v and the restriction μ_H of the metric g to H are G -invariant with respect to the action (26).

(ii) If H is an invariant submanifold of the system $(Q; l)$ then the trajectories $q(t)$ of the natural mechanical system with velocities $\dot{q}(t)$ that belong to H project to the trajectories $b(t) = \tau(q(t))$ of the natural mechanical system $(B; L)$ with the potential $V(b) = v(q)$ and the metric K obtained from μ_H via identification $H = G \times B$.

Note that when g is G -invariant, the twisted G -action (26) coincides with usual G -action: $g(q; X) = (g(q; d\tau(X)))$ and the induced metric K is the submersion metric. In this case Theorem 3 is exactly the classical method of E. J. Routh for eliminating cyclic coordinates [27, 2, 21].

Definition 2. By the analogy with the Lagrange-Routh reduction we shall call the procedure of passing from the Lagrangian system $(Q; l)$ to the system $(B; L)$ a partial Lagrange-Routh reduction.

Remark 5. The distribution (25), in general, is nonintegrable. One can interpret the restriction of the Lagrangian system $(Q; l)$ to H as a nonholonomic system $(Q; l; H)$ with a property that the reaction forces are equal to zero.

3.1. Geodesic Flows on Compact Lie Groups. Let G be a compact Lie group, \mathfrak{g} be the Lie algebra of G and $\langle \cdot, \cdot \rangle$ be a \mathfrak{A} -invariant scalar product on \mathfrak{g} . Let $a \in \mathfrak{g}$ be an arbitrary element and let b belongs to the center of \mathfrak{g}_a . Here $\mathfrak{g}_a = \{X \in \mathfrak{g} \mid [a, X] = 0\}$ is the isotropy algebra of a . Let $\mathfrak{g} = \mathfrak{g}_a \oplus \mathfrak{d}$ be the orthogonal

decomposition. Consider the linear operator (so called sectional operators [13]) $A_{a,b;C} : \mathfrak{g} \rightarrow \mathfrak{g}$, defined by

$$A_{a,b;C}(\xi) = \text{ad}_a^{-1} \text{ad}_b \text{pr}_{\mathfrak{g}}(\xi) + C(\text{pr}_{\mathfrak{g}_a}(\xi));$$

where $\text{pr}_{\mathfrak{d}}$ and $\text{pr}_{\mathfrak{g}_a}$ are the orthogonal (with respect to $\langle \cdot, \cdot \rangle$) projections to \mathfrak{d} and \mathfrak{g}_a , respectively and $C : \mathfrak{g}_a \rightarrow \mathfrak{g}_a$ is symmetric. We can always find b and C such that $A_{a,b;C}$ is positive definite.

Identify \mathfrak{g} with \mathfrak{g} by means of the scalar product $\langle \cdot, \cdot \rangle$ and consider the left-trivialization:

$$T G \rightarrow G \quad g = f(g; \xi)g;$$

Then the quadratic form

$$(27) \quad h_{a,b;C} = \frac{1}{2} h A_{a,b;C}(\xi); \xi$$

can be regarded as the Hamiltonian of a left-invariant Riemannian metric on G . Denote this metric by $h_{a,b;C}$.

Let G_a be the adjoint isotropy group of the element a and consider the right G_a -action on G . With the above notation, the momentum mapping and its zero level-set are given by

$$(28) \quad \mu(g; \xi) = \text{pr}_{\mathfrak{g}_a}(\xi);$$

$$(29) \quad (T G)_0 \rightarrow G \rightarrow \mathfrak{d};$$

Lemma 4. The metric $h_{a,b;C}$ is invariant with respect to the G_a -action, i.e., the momentum map (28) is preserved along the geodesic flow

$$(30) \quad \frac{d}{dt} \mu(\gamma; A_{a,b;C}(\dot{\gamma})) = 0; \quad \gamma = g \cdot A_{a,b;C}(\xi)$$

if and only if the quadratic form $h_{a,b;C}(\xi)$ is Ad_a -invariant:

$$(31) \quad \langle \xi; C(\xi) \rangle = 0; \quad \xi \in \mathfrak{g};$$

If a is regular element of the Lie algebra \mathfrak{g} , then \mathfrak{g}_a is Abelian and the condition (31) is satisfied for an arbitrary operator C .

Suppose (31) holds. Then we can project the metric $h_{a,b;C}$ to the homogeneous space G/G_a , that is to the adjoint orbit of $a: G/G_a \rightarrow O(a) = f \text{Ad}_g(a) \text{ for } g \in G$. Since we deal with the right action, the vertical distribution is left-invariant: $V_g = \mathfrak{g} \cdot g$, while from the definition of $h_{a,b;C}$, the horizontal distribution is $H_g = \mathfrak{g} \cdot \mathfrak{d}$ and the submersion metric does not depend on C .

Denote the submersion metric by $K_{a,b}$. The cotangent bundle $T^*O(a)$ can be realized as a submanifold of $\mathfrak{g} \times \mathfrak{g}$

$$(32) \quad T^*O(a) = \{ (x;p) \mid x = \text{Ad}_g(a); p \in \mathfrak{g}_x^2 \};$$

with the pairing between $p \in T_x^*O(a)$ and $\dot{x} \in T_xO(a)$ given by $\langle p, \dot{x} \rangle = \langle p; \dot{x} \rangle$. Let $b_x = \text{Ad}_g b$. Then the Hamiltonian and the geodesic flow for the metric $K_{a,b}$ in redundant variables $(x;p)$ are given by (see [6])

$$(33) \quad H_{a,b}(x;p) = \frac{1}{2} \langle \text{ad}_{b_x} p; \text{ad}_x p \rangle = \frac{1}{2} \langle \text{ad}_x \text{ad}_{b_x} p; p \rangle;$$

$$(34) \quad \dot{x} = \text{ad}_x \text{ad}_{b_x} p = [\mathfrak{b}_x; p]; x;$$

$$(35) \quad \dot{p} = \text{ad}_x^{-1} [\mathfrak{p}; [\mathfrak{b}_x; p]] + \text{pr}_{\mathfrak{g}_x} [\mathfrak{b}_x; p]; p;$$

Now, let us perturb the metric $h_{a,b;c}$ as follows. Take $h = h_B(\text{pr}_{g_a})$; $i + \frac{1}{2}h_C(\text{pr}_{g_a})$; $i, B : d! g_a, C : g_a! g_a$, such that

$$(36) \quad h(g;) = h_{a,b;c} + \frac{1}{2}h_A ; i$$

is positive definite. Then (36) will be the Hamiltonian function of the left-invariant metric that we shall denote by h . Since $h = h|_{(T G)_0}$, the geodesic flow of

$$(37) \quad \dot{g} = [; A ()]; \quad \dot{g} = g A ():$$

has the invariant relation (29). Hence we can perform the partial reduction. The metrics h and $h_{a,b;c}$ induce the same metric $K_{a,b}$ on the orbit $O(a)$, but their horizontal distributions H and H are different for $B \notin 0$:

$$H_g = \pi^{-1}(g - d) \in g - d = \mathbb{H}:$$

In the case when a is a singular element of g we can take $h = \frac{1}{2}h_C(\text{pr}_{g_a})$; i such that h is not A_{g_a} -invariant. Then the perturbed metric has the same horizontal distribution as the non-perturbed one: $H_g = g - d$.

The geodesic flow (34), (35) is completely integrable in the non-commutative sense. Moreover, the system is also integrable in the usual commutative sense by means of analytic functions, polynomial in momenta and which can be lifted to commuting G_a -invariant functions on $T G$ (see [5, 22, 6]). Thus, according Theorem 2 we obtain

Corollary 5. The equations (30) and (37) have the same invariant isotropic foliation of (29).

3.2. Local Description. To clear up the difference between geodesic flows (30) and (37), let us write down the problem in local coordinates.

Suppose a is a regular element of g . Then $G_a \cong T^n$ is a maximal torus. Locally, we have $T G \cong T O(a) \cong T T^n$. We take coordinates $(q;'; p;)$ in $T G$ such that

$$(q; p) = (q_1; \dots; q_m; p_1; \dots; p_m) \quad \text{and} \quad (q;';) = (q'_1; \dots; q'_n; p'_1; \dots; p'_n)$$

are canonical local coordinates on $T O(a)$ and $T T^n$. Locally, the right T^n -action is given by translations in q' coordinates and $(T G)_0 = f(q;'; p; 0)g$.

The Hamiltonians $h_{a,b;c}$ and h are of the form:

$$h_{a,b;c} = \frac{1}{2} \sum_{i,j} A^{ij} p_i p_j + \sum_i B^i p_i + \frac{1}{2} \sum_{k,j} C^{kj} p_k p_j ;$$

$$h = \frac{1}{2} \sum_{i,j} A^{ij} p_i p_j + \sum_i B^i p_i + \frac{1}{2} \sum_{k,j} C^{kj} p_k p_j ;$$

where A^{ij}, B^i, C^{kj} do not depend on the variables q' .

The partially reduced system

$$(38) \quad \dot{q}_i = \frac{\partial H_{a,b}}{\partial p_i} = \sum_{j,k} A^{ij} p_j ; \quad \dot{p}_i = - \frac{\partial H_{a,b}}{\partial q_i} = - \sum_{k,j} \frac{\partial A^{kj}}{\partial q_i} p_k p_j ; \quad i = 1; \dots; m$$

is completely integrable and we can treat $q(t)$ and $p(t)$ as known functions of time. Here $H_{a,b} = \frac{1}{2} \sum_{i,j} A^{ij} p_i p_j$ is the reduced Hamiltonian (33).

The equations of the geodesic flows (30) and (37) on (29) in variables $q_i; p_i$ are given by (38). The equations in variables q', p' , respectively, are given by:

$$\dot{q}'_i = \sum_{j,k} B^i(q) p_j \quad \text{and} \quad \dot{p}'_i = - \sum_{j,k} B^i(q;') p_j ; \quad i = 1; \dots; n:$$

While the first system is solvable, the second one, generically, is not. Globally, (29) is foliated on invariant tori with quasi-periodic and non-quasi-periodic flows of (30) and (37), respectively.

4. n-Dimensional Hess-Appel'rot System

Consider the motion of an n-dimensional rigid body around a fixed point $O = (0; 0; \dots; 0)$ in the n-dimensional Euclidean vector space $(\mathbb{R}^n; (\cdot, \cdot))$. The configuration space of the system is the Lie group $SO(n)$: the element $g \in SO(n)$ maps the moving coordinate system (attached to the body) to the fixed one (e.g., see [12]). Let $F_1; \dots; F_n, E_1; \dots; E_n$ and $f_1; \dots; f_n, e_1; \dots; e_n$ be the orthonormal bases attached to the body and fixed in the space regarded in the space frame and moving frame:

$$\begin{aligned} E_1 = f_1 = (1; 0; \dots; 0; 0)^T; \dots; E_n = f_n = (0; 0; \dots; 0; 1)^T; \\ E_i = g \cdot e_i; \dots; E_n = g \cdot e_n; F_1 = g \cdot f_1; \dots; F_n = g \cdot f_n \end{aligned}$$

We can consider the components of the vectors $e_1; \dots; e_n$ (or $F_1; \dots; F_n$) as redundant coordinates on $SO(n)$.

For a path $g(t) \in SO(n)$, the angular velocity in the body frame and angular velocity in the space frame are defined by $\dot{g}(t) = g^{-1} \dot{g}(t) \in \mathfrak{so}(n)$ and $\dot{g}(t) = g \cdot \dot{g}^{-1} = g \cdot \dot{g}^{-1} = g \cdot \dot{g}^{-1} = \text{Ad}_g \dot{g}^{-1}$, respectively. From the conditions $0 = E_i = g \cdot e_i$ and $F_i = g \cdot f_i$, the vectors $e_1; \dots; e_n$ and $F_1; \dots; F_n$ satisfy Poisson equations

$$\dot{e}_i = \dot{g} \cdot e_i \quad \dot{F}_i = \dot{g} \cdot F_i; \quad i = 1; \dots; n$$

The kinetic energy of a rigid body is a left-invariant quadratic form $\frac{1}{2} \langle I^{-1} \dot{g}, \dot{g} \rangle$, where $I : \mathfrak{so}(n) \rightarrow \mathfrak{so}(n)$ is a non-degenerate inertia operator and $\langle X, Y \rangle = \frac{1}{2} \text{tr}(XY)$ denotes the Killing metric on $\mathfrak{so}(n)$. For a "physical" rigid body, I^{-1} has the form $I^{-1} = I^{-1} + \langle \cdot, \cdot \rangle$, where I is a symmetric $n \times n$ matrix [12]. We will relax this condition, considering an arbitrary positive definite operator.

Further, suppose that the body is placed in the homogeneous gravitational force field in the direction e_n and the position of the center of mass of a rigid body is f_n . Then the potential is $v = -GM \langle f_n, e_n \rangle$, where G is the gravitational constant and M is the mass of the body. Let $m = I^{-1} \dot{g}$ be the angular momentum in the body frame and $A = I^{-1}$. By the use of Killing metric we can identify $\mathfrak{so}(n)$ and $\mathfrak{so}(n)$. Then the Hamiltonian in the left-trivialization

$$(39) \quad T_{SO(n)} \times_{SO(n)} \mathfrak{so}(n) = f(g; m)g$$

reads

$$h = \frac{1}{2} \langle m, m \rangle + GM \langle f_n, e_n \rangle$$

and the equations of the system, in redundant variables $(e_1; \dots; e_n; m)$, take the form of the Euler-Poisson equations

$$(40) \quad \dot{m} = [m, \dot{g}] + GM f_n \wedge e_n; \quad \dot{g} = Am$$

$$(41) \quad \dot{e}_i = \dot{g} \cdot e_i \quad i = 1; \dots; n$$

be identified with the positions of the vector F_n :

$$\begin{aligned} SO(n-1) &\cong SO(n) \\ &\cong \#_{S^{n-1}}; \quad (F_1; \dots; F_n) = F_n : \end{aligned}$$

The equations of motion (40), (41) in the space frame take the form

$$\begin{aligned} (48) \quad \dot{M} &= GM F_n \wedge E_n; \\ \dot{F}_i &= F_i; \quad i = 1; \dots; n; \end{aligned}$$

where the angular momentum M and velocity \dot{F}_i in the space are related to the angular momentum m and velocity \dot{f}_i in body coordinates via $M = Ad_g m$ and $\dot{F}_i = Ad_g \dot{f}_i$, respectively. Therefore

$$(49) \quad \dot{M} = Ad_g(\dot{m}) = Ad_g(A^{-1} Ad_1(M)):$$

Since $F_i \wedge F_j = Ad_g(f_i \wedge f_j)$, the invariant submanifold (42) in the right-trivialization reads

$$(\Gamma SO(n))_0 \cong f(g; D_g); D_g = Ad_g(d)g:$$

Let $K_g = Ad_g(k)$ and $X = X_{K_g} + X_{D_g}$ be the decomposition of $X \in \mathfrak{so}(n)$ with respect to $\mathfrak{so}(n) = K_g + D_g$. Note that the orthogonal projection to D_g given by

$$(50) \quad X_{D_g} = (X - F_n) \wedge F_n:$$

On $(\Gamma SO(n))_0$ we have $\dot{F}_i = A_d(\dot{m}) + B(\dot{m}) = a \dot{m} + B(\dot{m})$. Thus

$$\dot{F}_i = X_{K_g} + X_{D_g}; \quad X_{K_g} = Ad_g(B^{-1} Ad_1(\dot{M})); \quad X_{D_g} = a^{-1} \dot{M};$$

Furthermore, from (50) we have $F_n = D_g^{-1} F_n$. Therefore, on $(\Gamma SO(n))_0$, the Hess-Appell'rot system from the closed system in variables $(F_n; D_g)$:

$$\begin{aligned} (51) \quad -D_g &= a GM F_n \wedge E_n; \\ \dot{F}_n &= D_g^{-1} F_n; \end{aligned}$$

From the second equation we get

$$(52) \quad D_g = F_n \wedge F_n; \quad i.e.; \quad M = \frac{1}{a} F_n \wedge F_n$$

and the first equation can be rewritten in the form:

$$(53) \quad \dot{F}_n = a GM E_n + F_n:$$

Here the Lagrange multiplier is determined from the condition $(F_n; F_n) = 1$.

Therefore we obtain:

Theorem 7. The motion of the center of the mass of the n -dimensional Hess-Appell'rot system (40), (41), (42), (45) in the space frame is described by the spherical pendulum equations (53).

The same statement, for the classical Hess-Appell'rot system is proved by Zhukovski (see [29, 8]).

It is well known that the partially reduced system (53) is completely integrable in the non-commutative sense. The tangent bundle TS^{n-1} is foliated by two-dimensional invariant manifolds that are level-sets of the energy and integrals

$$\langle F_n \wedge F_n; E_i \wedge E_j \rangle = (F_n; E_i)(F_n; E_j) - (F_n; E_j)(F_n; E_i); \quad 1 \leq i < j \leq n-1:$$

Using (52), we see that the lifting of these integrals to $(T^*SO(n))_0$ are components of the momentum map of the left $SO(n-1)$ -action:

$$(54) \quad G_{ij} = a \, hM; \mathbb{E}^i \wedge E_j; \quad 1 \leq i < j \leq n-1$$

(note that (54) are integrals on the whole phase space $T^*SO(n)$ as well). Whence Corollary 8. The invariant set (42) of the Hess-Appellrot system is almost everywhere foliated by $(2 + (n-1)(n-2)=2)$ -dimensional isotropic invariant manifolds, level sets of the Hamiltonian function and integrals (54).

The above considerations motivate us for the following definition

Definition 3. We shall say that a Hamiltonian system (11) satisfies geometrical Hess-Appellrot conditions if it has an invariant relation (9) and the partially reduced system (15) is completely integrable.

The geodesic flows on compact Lie groups presented in Section 3 provide examples of systems that satisfy geometrical Hess-Appellrot conditions. Note that the condition that the partially reduced system is integrable differs from the notion of restricted integrability given in [11].

4.3. Reduction to $(so(n), \mathbb{R}^n)$. The system (40), (41) is always left $SO(n-1)$ -invariant (rotations of a rigid body "around" the vector e_n). Denote $\mathbb{E} = e_n$, $r = f_n$. The equations (40) together with the last Poisson equation,

$$(55) \quad \dot{m}_n = [m; \mathbb{E}] + GM \, r^{\wedge}; \quad \dot{r}_i = -\mathbb{E}^i; \quad \dot{r}_n = Am;$$

can be seen as a left $SO(n-1)$ -reduction of (40), (41). This is a Hamiltonian system on the dual space of Lie algebra $\mathfrak{e}(n) = so(n) \times \mathbb{R}^n$ with respect to the usual Lie-Poisson bracket. For $n=3$, after identification (47), the equations (55) get the familiar form (1).

Now suppose the Hess-Appellrot condition (45) is satisfied. Then the Euler-Poisson equations (55) on the invariant set (46) coordinately read:

$$(56) \quad \begin{aligned} \dot{m}_{in} &= \sum_{j=1}^{X-1} \mathbb{E}^j m_{jn} - GM \, \mathbb{E}^i; \quad i=1; \dots; n-1; \\ \dot{r}_i &= -a_n m_{in} - \sum_{j=1}^{X-1} \mathbb{E}^j r_j; \quad i=1; \dots; n-1; \\ \dot{r}_n &= a \sum_{j=1}^{X-1} m_{jn}; \end{aligned}$$

where $\mathbb{E}^i = hM; f_i \wedge f_j; \quad 1 \leq i < j \leq n-1$. The equations (56) always have three integrals

$$(57) \quad \begin{aligned} F_1 &= \frac{a}{2} \sum_{i=1}^{X-1} m_{in}^2 + GM \, r_n; \quad F_2 = \sum_{i=1}^{X-1} r_i^2 = 1; \\ F_3 &= \sum_{1 \leq i < j \leq n-1} (m_{in} r_j - m_{jn} r_i)^2; \end{aligned}$$

which correspond to integrals (2). As we saw above, the variable $\mathbb{E} = (e_n; f_n) = (E_n; F_n)$ satisfies the vertical component equation for the motion of the spherical pendulum and can be found by quadratures.

In general, equations (56) have no smooth invariant measure. The non-existence of a smooth invariant measure is the reflection of the non-solvability by quadratures of the classical system (1), (3), (4). Namely, by the Euler-Jacobi theorem (e.g., see [2], page 131), the invariant measure together with the integrals (2), (4) would imply solvability of the system by quadratures.

Example 1. As an illustration, consider the case $n = 4$ and Hamiltonian

$$h = \frac{1}{2}(a_1 m_{23}^2 + a_2 m_{13}^2 + a_3 m_{12}^2) + \frac{a}{2}(m_{14}^2 + m_{24}^2 + m_{34}^2) + b_1 m_{12} m_{14} + b_2 m_{12} m_{24} + b_3 m_{12} m_{34} + GM_n;$$

Then the equations in variables $m_{12}; m_{23}; m_{13}$ are

$$\begin{aligned} \underline{m}_{12} &= m_{13} m_{23} (a_2 - a_1) + m_{12} (b_1 m_{24} - b_2 m_{14}); \\ \underline{m}_{23} &= m_{12} m_{13} (a_3 - a_2) + m_{12} (b_2 m_{34} - b_3 m_{24}) + m_{13} (b_1 m_{14} + b_2 m_{24} + b_3 m_{34}); \\ \underline{m}_{13} &= m_{23} m_{12} (a_1 - a_3) + m_{12} (b_1 m_{34} - b_3 m_{14}) - m_{23} (b_1 m_{14} + b_2 m_{24} + b_3 m_{34}); \end{aligned}$$

while the equations in other variables, on the invariant set

$$(58) \quad m_{12} = m_{13} = m_{23} = 0;$$

take the form

$$\begin{aligned} \underline{m}_{14} &= m_{24} (b_1 m_{14} + b_2 m_{24} + b_3 m_{34}) - GM_{-1}; \\ \underline{m}_{24} &= m_{14} (b_1 m_{14} + b_2 m_{24} + b_3 m_{34}) - GM_{-2}; \\ \underline{m}_{34} &= GM_{-3}; \\ (59) \quad \underline{a}_1 &= a_4 m_{14} - 2 (b_1 m_{14} + b_2 m_{24} + b_3 m_{34}); \\ \underline{a}_2 &= a_4 m_{24} + \underline{a}_1 (b_1 m_{14} + b_2 m_{24} + b_3 m_{34}); \\ \underline{a}_3 &= a_4 m_{34}; \\ \underline{a}_4 &= a_4 (m_{14} + m_{24} + m_{34}); \end{aligned}$$

Together with (57) equations (59) have a supplementary integral

$$(60) \quad F_{12} = m_{14}^2 - m_{24}^2;$$

implying the foliation on 3-dimensional invariant manifolds of (58).

In redundant coordinates $(F_1; F_2; F_3; F_4; M)$ the integral (60) on $T^*SO(4)$ reads

$$\begin{aligned} (61) \quad F_{12} &= \int m_1 f_1 \wedge f_4 i(e_4; f_2) - \int m_2 f_2 \wedge f_4 i(e_4; f_1) \\ &= \int m_1 f_1 \wedge F_4 i(E_4; F_2) - \int m_2 f_2 \wedge F_4 i(E_4; F_1); \end{aligned}$$

It is clear that F_{12} is independent of integrals (54). Whence the 5-dimensional invariant isotropic manifolds outlined in Corollary 8 are foliated on 4-dimensional invariant level-sets of (61).

Note that the system (59) has an invariant measure if and only if $b_1 = b_2 = 0$. Also, recall that for $a_1 = a_2 = a_3 \notin a, b_1 = b_2 = b_3 = 0$, the Euler-Poisson equations represent the motion of a dynamically symmetric heavy rigid body, 4-dimensional version of the Lagrange top [3].

Therefore, if $b_1 = b_2 = b_3 = 0$ and $a_1 \notin a_2 \notin a_3$ then the equations (59) coincides to the 4-dimensional Lagrange top equations restricted to the invariant

level-set (58), although the systems are different on the whole phase space. They have additional integrals

$$(62) \quad \begin{aligned} F_{13} &= m_{14} m_{31} (= hM ; F_1 \wedge F_3 i(E_4; F_3) - hM ; F_3 \wedge F_4 i(E_4; F_1)); \\ F_{23} &= m_{24} m_{34} (= hM ; F_2 \wedge F_4 i(E_4; F_3) - hM ; F_3 \wedge F_4 i(E_4; F_2)); \end{aligned}$$

Since the equations (59) preserve the standard measure in variables $m_{14}; m_{24}; m_{34}, m_{11}; m_{22}; m_{33}; m_{44}$ and the invariant manifolds given by integrals (57), (60), (62) are two-dimensional, by the Euler-Jacobi theorem, they are solvable by quadratures. On the other side, the Hamiltonian and functions (54), (61), (62) provide 3-dimensional invariant foliation of 9-dimensional manifold $(T^*SO(4))_0$.

4.4. L-A Pair. The non-commutative integrability of the n-dimensional Lagrange top is proved by Beljajev [3] and the L-A pair is given by Reymann and Semenov-Tian-Shansky [26]. The L-A pair of the similar form can be used for the Hess-Appel'rot system on $so(n) \times \mathbb{R}^n$. Let $m, !, r$ be the $so(n+1)$ -matrices given by

$$\begin{aligned} m &= \begin{pmatrix} m & 0 \\ 0 & 0 \end{pmatrix}; \quad = \begin{pmatrix} 0 & \\ & t & 0 \end{pmatrix}; \\ ! &= \begin{pmatrix} ! & 0 \\ 0 & 0 \end{pmatrix}; \quad r = \begin{pmatrix} 0 & f_n \\ f_n^t & 0 \end{pmatrix}; \end{aligned}$$

where 0 is the zero $n \times n$ matrix.

Then, under the Hess-Appel'rot conditions (45) and (46), the Euler-Poisson equations (55) are equivalent to the matrix equation with a spectral parameter

$$(63) \quad L(\lambda) = [L(\lambda); A(\lambda)];$$

where $L(\lambda) = m + \lambda^{-2} \frac{1}{a} GM r$ and $A(\lambda) = ! + GM r$.

Indeed, the straightforward computations show that the terms with λ^{-4} and λ^{-1} in (63) are equivalent to the equations (55). The left hand side term with λ^{-2} is identically equal to zero. It can be proved that the right hand side term with λ^{-2} vanishes, on the invariant submanifold (46), if and only if (45) holds.

On the invariant set (46) the spectral curve is

$$(64) \quad \begin{aligned} p(\lambda; \mu) &= \det(L(\lambda) - \mu Id) = (\mu)^n - \lambda^{-3} \mu^4 + \lambda^{-2} P(\mu) + Q(\mu)^2 = 0; \\ P(\mu) &= F_2 + \frac{2}{a} \mu^2 F_1 + \mu^4 \frac{GM}{a}; \quad Q(\mu) = F_3; \end{aligned}$$

where $F_1; F_2; F_3$ are integrals (57). The curve (64) is reducible (for $n > 3$) and consists of the $n-3$ copies of the rational curve $\mu = 0$ and the singular algebraic curve

$$(65) \quad \mu^4 + \lambda^{-2} P(\mu) + Q(\mu)^2 = 0;$$

The spectral curves of the form (65) appear in the algebro-geometric analysis given in [10, 11].

In particular the above consideration leads to the L-A pair of the classical system different from those obtained in [9].

Theorem 9. Under the Hess-Appel'rot conditions (5), (6), (7), the Euler-Poisson equations (1) are equivalent to the matrix equation (63), where

$$L(\cdot) = \begin{pmatrix} 0 & 0 & m_3 & m_2 & 1 & 1 \\ m_3 & 0 & m_1 & 0 & 2 & 2 \\ m_2 & m_1 & 0 & 0 & 3 & 2 \\ 0 & 1 & 2 & 3 & 2 & 2 \\ 0 & 0 & 1 & 2 & 3 & 2 \\ 0 & 1 & 2 & 3 & 2 & 2 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \frac{2GM}{a_2} \frac{C}{A};$$

$$A(\cdot) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \frac{C}{A};$$

Remark 6. Recall that the spectral curve for the classical system given in [9] is reducible and consists of the rational and the elliptic curve. In our case, due to the involution $(\cdot; \cdot) \mapsto (\cdot; \cdot)$, the singular spectral curve (65) is a two-covering of the regular (for the generic values of the integrals (2)), genus-3 hyperelliptic curve

$$0 = u^2 + uP(\cdot) + Q(\cdot)^2 = 0$$

(e.g., see Lemma 1 in [10]). In the above part, the point $(0;0)$ is an ordinary double point as well as the ramification point of the covering.

5. Hess-Appel'rot system on $(so(4), so(4))$

Dragovic and Gajić defined a class of systems of the Hess-Appel'rot type by using the analytical and algebraic properties of the classical system [11]. Subtle difference between Dragovic-Gajić systems and those constructed here has to be studied carefully, but there is a big coherence between these two approaches. Remarkably, the $(so(4), so(4))$ system given in [11] considered on the whole phase space $T^*SO(4)$ satisfies geometrical Hess-Appel'rot conditions as well.

We start from the example of the geodesic flows given in Section 3. Suppose, in addition, the G_a -invariant potential force $v(g)$ is given. Then (29) remains to be an invariant manifold for natural mechanical systems with kinetic energies $K_{a,b,c}$ and \dots . After Lagrange-Routh and partial Lagrange-Routh reductions, the systems project to the system with kinetic energy $K_{a,b}$ and potential force $V(x)$ on the adjoint orbit $O(a)$, where $V(x) = v(g)$, $x = Ad_g a$.

The reduced system is a modification of (34), (35) by the potential force:

$$(66) \quad \dot{x} = ad_x ad_{b_x} p = [b_x; p; x];$$

$$(67) \quad \dot{p} = ad_x^{-1} [p; x; b_x; p] - \frac{\partial V(x)}{\partial x} + \sum_{i=1}^n \lambda_i \xi_i(x);$$

Here $\xi_i(x); \dots; \xi_n(x)$ is a base of \mathfrak{g}_x and Lagrange multipliers λ_i are chosen such that trajectory $(x(t); p(t))$ belongs to (32). For a proportional to b and a linear potential $V(x) = \langle h; x \rangle$, the system represents analog of spherical pendulum on adjoint orbit $O(a)$, which is integrable on an arbitrary orbit $O(a)$ (see Bolsinov and Jovanovic [7]).

Now consider the above construction for the case $G = SO(4)$, i.e., for the 4-dimensional rigid body motion about a fixed point (we use notation from the previous section). Let

$$(68) \quad b = (J_1 + J_3)a; \quad a = a_{12} f_1 \wedge f_2 + a_{34} f_3 \wedge f_4;$$

We have $\mathfrak{so}(4)_a = \text{span}\{f_1 \wedge f_2, f_3 \wedge f_4\} = \mathfrak{so}(2) \oplus \mathfrak{so}(2)$. Define

$$C : \mathfrak{so}(4)_a \rightarrow \mathfrak{so}(4)_a; \quad C(f_1 \wedge f_2) = 2J_1 f_1 \wedge f_2; \quad C(f_3 \wedge f_4) = 2J_3 f_3 \wedge f_4;$$

Then the left-invariant kinetic energy $h_{a,b;C}$ (see (27)) takes the form

$$h_{a,b;C}(g; m) = J_1 m_{12}^2 + J_3 m_{34}^2 + \frac{J_1 + J_3}{2} (m_{13}^2 + m_{14}^2 + m_{23}^2 + m_{24}^2);$$

Consider the perturbation of the metric

$$(69) \quad h = h_{a,b;C} + \frac{1}{2} h_A m; m; i; \\ (g; m) = m_{12} (J_{14} m_{14} - J_{13} m_{23}) + m_{34} (J_{13} m_{14} - J_{14} m_{23})$$

and the potential function

$$(70) \quad v = h \text{Ad}_{g^{-1}}(a); a; i = h a_{12} e_1 \wedge e_2 + a_{34} e_3 \wedge e_4; a; i \\ = h a; A \text{d}_g(a); i = h a; a_{12} F_1 \wedge F_2 + a_{34} F_3 \wedge F_4; i$$

(written in the coordinates $(e_1; e_2; e_3; e_4)$ and $(F_1; F_2; F_3; F_4)$, respectively).

The rigid body system with Hamiltonian $h = h + v$, in the left-trivialization (39), takes the form

$$(71) \quad \underline{m} = [m; A(m)] + [a_{12} e_1 \wedge e_2 + a_{34} e_3 \wedge e_4; a]; \quad \underline{g} = g \cdot A(m);$$

The system (71) has two invariant relations

$$(72) \quad m_{12} = 0; \quad m_{34} = 0$$

and it is reducible to the oriented Grassmannian variety $G r^+(4;2) \subset O(a)$ of oriented two-dimensional planes in the four-dimensional vector space. The diffeomorphism $\pi : O(a) \rightarrow G r^+(4;2)$ is simply given by

$$(\underline{x}) = F_1 \wedge F_2; \quad \text{where } \underline{x} = \text{Ad}_g(a) = a_{12} F_1 \wedge F_2 + a_{34} F_3 \wedge F_4;$$

Recall that $F_1; F_2; F_3; F_4$ is the moving base regarded in the space frame e , so $F_1 \wedge F_2$ is the oriented two-plane attached to the body.

From (33), (68), (70) we get the reduced Hamiltonian

$$H(\underline{x}; p) = H_{a,b}(\underline{x}; p) + V(\underline{x}) = \frac{J_1 + J_3}{2} h[\underline{x}; p]; [\underline{x}; p]; i + h \underline{x}; a; i$$

After calculating the Lagrange multipliers in (66), (67) the partially reduced system on $T G r^+(4;2)$ become

$$(73) \quad \underline{\dot{x}} = (J_1 + J_3) [\underline{x}; p]; \underline{x};$$

$$(74) \quad \underline{\dot{p}} = (J_1 + J_3) [\underline{x}; p]; p] \cdot a + p r_{g_x} a;$$

It follows from Theorem 4 [7] that the reduced system is completely integrable by means of integrals that can be extended to commuting $SO(4)_a$ -invariant functions on $T SO(4)$. Hence we get the following qualitative behavior of the system

Theorem 10. The rigid body system (71) satisfies geometrical Hess-Appellrot conditions: the partial reduction of the system from (72) to the oriented Grassmannian variety $G r^+(4;2)$ is completely integrable pendulum type system (73), (74). Further, 10-dimensional invariant manifold (72) is almost everywhere foliated by invariant 6-dimensional Lagrangian tori that project to the 4-dimensional Liouville tori of the reduced system (73), (74). We need two additional quadratures to solve the reconstruction problem.

By introducing $\mu = a_{12}e_1 \wedge e_2 + a_{34}e_3 \wedge e_4$, from (71) we can write the closed system in variables $(m; \mu)$:

$$(75) \quad \underline{m} = [m; \mu] + [; a]; \quad \underline{\mu} = [; \mu]; \quad \dot{\mu} = A(m; \mu);$$

Regarding μ as the free $\mathfrak{so}(4)$ -variable, the system (75) becomes the Hamiltonian system with respect to the Lie-Poisson bracket on the dual space of the semidirect product $\mathfrak{so}(4) \ltimes \mathfrak{so}(4)$ (see Ratiu [25]). Moreover, with the above choice of h we have

$$A(m; \mu) = Jm + mJ; \quad J = \begin{matrix} 0 & & & & 1 \\ & J_1 & & & 0 \\ & & 0 & & J_{13} \\ & & & 0 & & 0 \\ & & & & & & C \\ & & & & & & & A \end{matrix};$$

and (75) coincides with the Hess-Appel'rot system defined in [11].

Remark 7. A similar statement can be proved for the Hess-Appel'rot systems on $\mathfrak{so}(n) \ltimes \mathfrak{so}(n)$ for $n > 4$ (see [11]), where one should take

$$b = (J_1 + J_3)a; \quad a = a_{12}f_1 \wedge f_2;$$

Then $\mathfrak{so}(n)_a = \mathfrak{so}(2) \oplus \mathfrak{so}(n-2)$ and the systems are partially reducible to the orbits $O(a)$, which are now diffeomorphic to the oriented Grassmannian varieties $Gr^+(n; 2)$ of oriented two-dimensional planes in the n -dimensional vector space.

Relations (72) define 6-dimensional invariant manifolds within generic symplectic leaves of $(\mathfrak{so}(4) \ltimes \mathfrak{so}(4))$. Apart of the Hamiltonian function, the system (75) has a supplementary integral (see [11])

$$F = {}_{34}a_{12} + {}_{12}a_{34} + (J_1 + J_3)(m_{12}m_{34} + m_{23}m_{14} - m_{13}m_{24})$$

implying the foliation on 4-dimensional invariant varieties. The L-A pair representation as well as the algebro-geometric integration of (75) up to two quadratures are given in [11].

Acknowledgments. I am very grateful to Vladimir Dragovic and Borislav Gajić for useful discussions and to the referees for suggestions which help me to improve the exposition of the paper. The research was supported by the Serbian Ministry of Science, Project "Geometry and Topology of Manifolds and Integrable Dynamical Systems".

References

- [1] Appel'rot, G. G.: The problem of motion of a rigid body about a fixed point. Uchenye Zap. Mosk. Univ. Otdel. Fiz. Mat. Nauk 11 (1894) 1-112 (in Russian).
- [2] Arnold, V. I., Kozlov, V. V. and Neishtadt, A. I.: Mathematical aspects of classical and celestial mechanics. Itogi Nauki i Tekhniki. Sovr. Probl. Mat. Fundamental'nye Napravleniya, Vol. 3, VINITI, Moscow 1985. English transl.: Encyclopaedia of Math. Sciences, Vol. 3, Springer-Verlag, Berlin 1989.
- [3] Beljaev, A. V.: Motion of a multidimensional rigid body with a fixed point in a gravitational field, Mat. Sb. 114 (156) (1981) no. 3, 465-470 (Russian).
- [4] Benenti, S. and Tulczyjew, W. M.: Remarques sur les reductions symplectiques, C. R. Acad. Sc. Paris, 294 Serie I (1982) 561-564.
- [5] Bolsinov, A. V. and Jovanovic, B.: Complete involutive algebras of functions on cotangent bundles of homogeneous spaces. Mathematische Zeitschrift 246 no. 1-2, 213-236 (2004).
- [6] Bolsinov, A. V. and Jovanovic, B.: Magnetic Geodesic Flows on Coadjoint Orbits, J. Phys. A: Math. Gen. 39, L247-L252 (2006), arXiv: math-ph/0602016.

- [7] Bolsinov, A. V. and Jovanovic, B.: Magnetic Flows on Homogeneous Spaces, arXiv: math-ph/0609005.
- [8] Borisov A. V. and Mamaev I. S.: The Hess case in the dynamics of a rigid body, Pribl. Mat. Mekh. 67 (2003) no. 2, 256-265 (Russian); English transl. J. Appl. Math. Mech. 67 (2003) no. 2, 227-235.
- [9] Dragovic, V. and Gajić, B.: An L-A pair for the Hess-Apffelrot system and a new integrable case for the Euler-Poisson equations on $\mathfrak{so}(4)$ $\mathfrak{so}(4)$. Proc. Roy. Soc. Edinburgh Sect. A 131 (2001) no. 4, 845-855, arXiv math-ph/9911047.
- [10] Dragovic, V. and Gajić, B.: The Lagrange bitop on $\mathfrak{so}(4)$ $\mathfrak{so}(4)$ and geometry of the Pym varieties. Amer. J. Math. 126 (2004) no. 5, 981-1004, arXiv: math-ph/0201036.
- [11] Dragovic, V. Gajić, B.: Systems of Hess-Apffelrot type, Comm. Math. Phys. 256 (2006) no. 2, 397-435, arXiv: math-ph/0602022.
- [12] Fedorov, Yu. N. and Kozlov, V. V.: Various aspects of n-dimensional rigid body dynamics, Amer. Math. Soc. Transl. Series 2, 168 (1995), 141-171.
- [13] Fomenko, A. T. and Troshchinskii, V. V.: Geometric and algebraic mechanisms of the integrability of Hamiltonian systems on homogeneous spaces and Lie algebras. In: Dynamical Systems VII (Eds.: V. I. Arnold, S. P. Novikov), Springer, 1994.
- [14] Golubev, V. V.: Lectures on integration of the equations of motion of a rigid body about a fixed point. Moskva, Gostekhizdat, 1953 (in Russian); English translation: Transl. Philadelphia, PA: Coronet Books, 1953.
- [15] Hess, W.: Über die Euler'schen Bewegungsgleichungen und über eine neue particulare Lösung des Problems der Bewegung eines starren Körpers um einen festen Punkt. Math. Ann. 37 (1890) no. 2, 178-180.
- [16] Jovanovic, B.: On the Integrability of Geodesic Flows of Submanifolds, Lett. Math. Phys. 61 (2002) 29-39; arXiv: math-ph/0204048.
- [17] Levi-Civita, T. and Amaldi, V.: Lezioni di Meccanica Razionale. Volume Secondo, Parte Seconda, Bologna, 1927.
- [18] Libermann, P. and Marle, C. M.: Symplectic geometry and analytic mechanics. Math. and its Appl. 35, Reidel, Boston, 1987.
- [19] Lichnerowicz, A.: Variété symplectique et dynamique associée à une sous-variété, C. R. Acad. Sc. Paris 280 Serie A (1975) 523-527.
- [20] Marsden, J. E. and Weinstein A.: Reduction of symplectic manifolds with symmetry, Rep. Math. Phys. 5 (1974) 121-130.
- [21] Marsden, J. E., Ratiu, T. S. and Scheurle J.: Reduction theory and the Lagrange-Routh reduction. J. Math. Phys. 41 (2000), no. 6, 3379-3429.
- [22] Mykytyuk, I. V. and Panasyuk A.: Bi-Poisson structures and integrability of geodesic flows on homogeneous spaces. Transformation Groups 9, no. 3, 289-308 (2004)
- [23] Nekhoroshev, N. N.: Action-angle variables and their generalization. Tr. Mosk. Mat. O.-va. 26 (1972) 181-198, (Russian); English translation: Trans. Mosc. Math. Soc. 26 (1972) 180-198.
- [24] Ortega, J. P. and Ratiu, T. S.: Momentum maps and Hamiltonian reduction. Progress in Math. 222, Birkhäuser, 2004.
- [25] Ratiu, T.: Euler-Poisson equations on Lie algebras and the N-dimensional heavy rigid body. Amer. J. Math. 104 (1982) 409-448.
- [26] Reyman, A. G. and Semenov-Tian-Shansky, M. A.: Lax representation with a spectral parameter for the Kovalevskaya top and its generalizations. Funktsional. Anal. i Prilozhen. 22 (1988), no. 2, 87-88 (Russian); translation in Funct. Anal. Appl. 22 (1988), no. 2, 158-160.
- [27] Routh, E. J.: Treatise on the Dynamics of a System of Rigid Bodies. Macmillan, London 1860.
- [28] Zung, N. T.: Torus actions and integrable systems In: Topological methods in the theory of integrable systems (Bolsinov A. V., Fomenko A. T., Oshemkov A. A. eds) Cambridge Scientific Publ., (to appear), math DS/0407455
- [29] Zhukovskii, N. E.: Geometrische Interpretation des Hess'schen Falles der Bewegung eines schweren starren Körpers um einen festen Punkt. Jber. Deutschen Math. Verein. 3 (1894) 62-70.