

A symmetry test for quasilinear coupled systems

V.V. Sokolov

Centre for Nonlinear Studies &
Landau Institute for Theoretical Physics,
Kosygina 2, Moscow 117940, Russia
em ail: sokolov@landau.ac.ru

Thomas Wolf

Queen Mary & Westfield College, University of London,
Mile End Road, London E1 4NS, UK
em ail: T.Wolf@maths.qmw.ac.uk

May 3, 2021

Abstract

All quasilinear NLS-type systems with fourth order symmetry are listed. Vector generalizations of some of them are constructed. Local master symmetries for several systems from the list are found.

1 Classification result.

It is well known that the following class of systems of evolution equations

$$\begin{aligned} u_t &= u_{xx} + F(u; v; u_x; v_x); \\ v_t &= v_{xx} + G(u; v; u_x; v_x); \end{aligned} \quad (1)$$

is very rich in integrable cases. In the papers [1]–[5] by Mikhailov, Shabat and Yamilov, all systems (1), possessing higher conservation laws, were classified. Hence, these authors have found all systems (1) that can be integrated by the inverse scattering method (S-integrable equations in the terminology by F. Calogero [6]).

However, there are integrable cases that are not in their classification. As an example, consider the system

$$\begin{aligned} u_t &= u_{xx} - 2uu_x - 2vu_x - 2uv_x + 2u^2v + 2uv^2; \\ v_t &= v_{xx} + 2vu_x + 2uv_x + 2vv_x - 2u^2v - 2uv^2; \end{aligned} \quad (2)$$

rst discussed in [7] (see [8] for generalizations). It can be reduced to

$$U_t = U_{xx}; \quad V_t = V_{xx}$$

by the following substitution of the Cole-Hopf type

$$u = \frac{U_x}{(U + V)}; \quad v = \frac{V_x}{(U + V)}; \quad (3)$$

The system (2) has no higher order conservation laws, but it has higher order symmetries. This is a typical feature of linearizable systems like the Burgers equation (C-integrable equations). Therefore, it would be interesting to classify the systems (1), which have higher order symmetries. As a result, all S-integrable and C-integrable systems would be found.

The complete classification problem is very difficult. Here we consider only the most interesting (from our opinion) subclass of systems (1). Namely, we consider equations linear in all derivatives of the form

$$\begin{aligned} u_t &= u_{xx} + A_1(u;v)u_x + A_2(u;v)v_x + A_0(u;v) \\ v_t &= v_{xx} + B_1(u;v)v_x + B_2(u;v)u_x + B_0(u;v); \end{aligned} \quad (4)$$

without any restrictions on the functions $A_i(u;v); B_i(u;v)$. We apply to such systems the simplest version of the symmetry test (see [9]-[12]).

Lemma. If system (4) has a fourth order symmetry

$$\begin{aligned} u &= u_{xxxx} + f(u;v;u_x;v_x;u_{xx};v_{xx};u_{xxx};v_{xxx}); \\ v &= v_{xxxx} + g(u;v;u_x;v_x;u_{xx};v_{xx};u_{xxx};v_{xxx}); \end{aligned} \quad (5)$$

then the system is of the following form

$$\begin{aligned} u_t &= u_{xx} + (a_{12}uv + a_1u + a_2v + a_0)u_x + (p_2v + p_{11}u^2 + p_1u + p_0)v_x + A_0(u;v); \\ v_t &= v_{xx} + (b_{12}uv + b_1v + b_2u + b_0)v_x + (q_2u + q_{11}v^2 + q_1v + q_0)u_x + B_0(u;v); \end{aligned}$$

where A_0 and B_0 are polynomials of at most fifth degree.

The coefficients of the last system satisfy an overdetermined system of algebraic equations. The most essential equations are

$$\begin{aligned} p_2(b_{12} - q_{11}) &= 0; & p_2(a_{12} - p_{11}) &= 0; & p_2(a_{12} + 2b_{12}) &= 0; \\ q_2(b_{12} - q_{11}) &= 0; & q_2(a_{12} - p_{11}) &= 0; & q_2(b_{12} + 2a_{12}) &= 0; \end{aligned}$$

$$\begin{aligned} a_{12}(a_{12} - b_{12} + q_{11} - p_{11}) &= 0; & b_{12}(a_{12} - b_{12} + q_{11} - p_{11}) &= 0; \\ (a_{12} - p_{11})(p_{11} - q_{11}) &= 0; & (b_{12} - q_{11})(p_{11} - q_{11}) &= 0; \\ (a_{12} - p_{11})(a_{12} - b_{12}) &= 0; & (b_{12} - q_{11})(a_{12} - b_{12}) &= 0; \end{aligned}$$

As usual, such factorized equations lead to a tree of variants, which was investigated by the computer algebra program Crack¹ [19].

Solving the overdetermined system, we don't consider so called triangular systems like the following

$$u_t = u_{xx} + 2uv_x; \quad v_t = v_{xx} - 2vv_x; \quad (6)$$

Here the second equation is separated and the first is linear with the variable coefficients defined by a given solution of the second equation.

Theorem . Any nonlinear nontriangular system (4), having a symmetry (5), up to scalings of t, x, u, v , shifts of u and v , and the involution

$$u \leftrightarrow v; \quad t \leftrightarrow -t \quad (7)$$

belongs to the following list:

$$\begin{aligned} u_t &= u_{xx} + (u + v)u_x + uv_x; \\ v_t &= v_{xx} + (u + v)v_x + vu_x; \end{aligned} \quad (8)$$

$$\begin{aligned} (\quad & u_t = u_{xx} - 2(u + v)u_x - 2uv_x + 2u^2v + 2uv^2 + u + v + ; \\ & v_t = v_{xx} + 2(u + v)v_x + 2vu_x - 2u^2v - 2uv^2 - u - v - ; \end{aligned} \quad (9)$$

$$\begin{aligned} u_t &= u_{xx} + vu_x + uv_x; \\ v_t &= v_{xx} + vv_x + u_x; \end{aligned} \quad (10)$$

$$\begin{aligned} u_t &= u_{xx} + 2vu_x + 2uv_x + 2uv^2 + u^2 + u + v + ; \\ v_t &= v_{xx} - 2vv_x - u_x; \end{aligned} \quad (11)$$

$$\begin{aligned} (\quad & u_t = u_{xx} + v_x + (u + v)^2 + (u + v) + ; \\ & v_t = v_{xx} + u_x - (u + v)^2 - (u + v) - ; \end{aligned} \quad (12)$$

$$\begin{aligned} (\quad & u_t = u_{xx} + (u + v)u_x + 4v_x + (u + v)^2 + (u + v) + ; \\ & v_t = v_{xx} + (u + v)v_x + 4u_x - (u + v)^2 - (u + v) - ; \end{aligned} \quad (13)$$

$$\begin{aligned} (\quad & u_t = u_{xx} + 2u^2v_x + 2uvu_x + (- 2)u^3v^2 + u^2v + u; \\ & v_t = v_{xx} + 2v^2u_x + 2uvv_x - (- 2)u^2v^3 - uv^2 - v; \end{aligned} \quad (14)$$

¹Wolff, T. and Brand, A., The Computer Algebra Package CRACK for Investigating PDEs, manual + software in the REDUCE network library (1992)

$$\begin{cases} u_t = u_{xx} + 2uvu_x + (\alpha + u^2)v_x; \\ v_t = v_{xx} + 2uvv_x + (\alpha + v^2)u_x; \end{cases} \quad (15)$$

$$\begin{cases} u_t = u_{xx} + 2uvu_x + 2u^2v_x + u^3v^2 + \alpha u; \\ v_t = v_{xx} + 2v^2u_x + 2uvv_x + u^2v^3 + \alpha v; \end{cases} \quad (16)$$

$$\begin{cases} u_t = u_{xx} + 2uvu_x + 2(\alpha + u^2)v_x + u^3v^2 + \alpha u^3 + uv^2 + \alpha u; \\ v_t = v_{xx} + 2uvv_x + 2(\alpha + v^2)u_x + u^2v^3 + \alpha u^2v + \alpha v^3 + \alpha v; \end{cases} \quad (17)$$

$$\begin{cases} u_t = u_{xx} + 4uvu_x + 4u^2v_x + 3vv_x + 2u^3v^2 + uv^3 + \alpha u; \\ v_t = v_{xx} + 2v^2u_x + 2uvv_x + 2u^2v^3 + v^4 + \alpha v; \end{cases} \quad (18)$$

$$\begin{cases} u_t = u_{xx} + 4uu_x + 2vv_x; \\ v_t = v_{xx} + 2vu_x + 2uv_x + 3u^2v + v^3 + \alpha v; \end{cases} \quad (19)$$

$$\begin{cases} u_t = u_{xx} + vv_x; \\ v_t = v_{xx} + uu_x; \end{cases} \quad (20)$$

$$\begin{cases} u_t = u_{xx} + 6(u+v)v_x + 6(u+v)^3 + (u+v)^2 + (u+v) + \alpha; \\ v_t = v_{xx} + 6(u+v)u_x + 6(u+v)^3 + (u+v)^2 + (u+v) + \alpha; \end{cases} \quad (21)$$

$$\begin{cases} u_t = u_{xx} + vv_x; \\ v_t = v_{xx} + uu_x; \end{cases} \quad (22)$$

We omitted the term $(cu_x; cv_x)^T$ on the right hand sides of all systems. It is a Lie symmetry, corresponding to the invariance of our classification problem with respect to the shift of x .

2 Discussion.

1. Admissible transformations. Some equations in the list contain arbitrary constants $\alpha; \beta; \gamma$. Not all of them are essential.

Let us consider, for instance, the equations (15). It is easy to see that those constants α and β which are not equal to zero can be reduced to 1 via scalings of $t; x; u$ and v . In this way, (15) actually describes three different equations without parameters. These equations correspond to $\alpha = \beta = 1$, $\alpha = \beta = 0$ and $\alpha = 1; \beta = 0$.

The following parameters: α in (12) and (13), β in (14), both α and β in (17) are not essential in the same sense. For (14) and (16) the essential parameter is the ratio of α and β . Note, that if $\alpha = \beta = 0$ then (14) coincides with the nonlinear Schrödinger equation, which is not a separate equation in our list.

For some equations from the list, there exist admissible transformations of the form

$$u \rightarrow p(x;t)u + q(x;t); \quad v \rightarrow r(x;t)v + s(x;t); \quad (23)$$

"Admissible" means that the resulting equation does not depend explicitly on x and t and has the same form (4). Using such admissible transformations, one can remove some of constants in the equations of the list.

In particular, with the help of the transformation $u \rightarrow \exp(ct)u; v \rightarrow \exp(-ct)v$ one can remove the terms $(cu; -cv)^T$ in (14), (16), (17).

Using the transformations $u \rightarrow u + t + x; v \rightarrow v - t - x$ it is possible to remove u and v in (12), (13). The equation (21) can be reduced to the form

$$\begin{cases} u_t = u_{xx} + 6(u+v)v_x - 6(u+v)^3 + cv_x; \\ v_t = v_{xx} + 6(u+v)u_x + 6(u+v)^3 + cu_x \end{cases} \quad (24)$$

by such a transformation and by shifts of u and v . It seems to us that the essential constant c was missed in the classification result of [3].

More general transformations are described in [4],[5] which reduce some of the equations (8)–(22) to others in this list. For simplicity in applying our results, we will not rely on these non-trivial transformations, and will instead operate with the complete list (8)–(22).

2. Three groups of equations. All equations of the list can be divided into three groups. The first group contains the so called NLS-type equations (8), (10), (13), (14), (15). Besides a higher symmetry (5) every such equation possesses a symmetry of the form

$$\begin{aligned} u &= u_{xxx} + (u;v;u_x;v_x;u_{xx};v_{xx}); \\ v &= v_{xxx} + (u;v;u_x;v_x;u_{xx};v_{xx}); \end{aligned} \quad (25)$$

This is typical for equations having the Lax representations in $sl(2)$.

The equations of the Boussinesq type form the second group (12), (20), (21), (22). They have no symmetries of third order. This indicates the existence of a Lax representation in $sl(3)$. We have chosen the existence of the symmetry (5) as a criterion of integrability for (4) since the choice of the simplest ansatz (25) leads to the loss of all equations of the second group.

The last group (9), (11), (16), (17), (18), (19) consists of "linearizable" equations, which have no higher conservation laws. Some of them seem to be new.

Equations (16), (17) : In [18],[6] one can find a linearization procedure for (16). Namely, a non-local substitution

$$U = u \exp \int u v dx; \quad V = v \exp \int u v dx \quad (26)$$

reduces it to the linear equation $U_t = U_{xx} + U; V_t = V_{xx} - V$:

Consider equation (17). It is easy to see that it has the following symmetry

$$\begin{cases} u = u_{xxx} + 3uvu_{xx} + 6uu_xv_x + 3vu_x^2 + 3u^2v^2u_x; \\ v = v_{xxx} + 3uvv_{xx} + 6vu_xv_x + 3uv_x^2 + 3u^2v^2v_x; \end{cases} \quad (27)$$

for any u, v ; Under a reduction $u = v$, (27) coincides with the well known equation (see [13], [11], [6])

$$u_t = u_{xxx} + 3u^2 u_{xx} + 9uu_x^2 + 3u^4 u_x;$$

which can be linearized by the substitution $U = u \exp^R u^2 dx$. It is easy to verify that the following very similar substitution (cf. also with (26))

$$U = u \exp^Z uv dx; \quad V = v \exp^Z uv dx \quad (28)$$

reduces (27) to $U_t = U_{xxx}; V_t = V_{xxx}$. The same substitution reduces (17) to a linear system

$$U_t = U_{xx} + 2V_x + U; \quad V_t = V_{xx} - 2U_x - V;$$

Generalizing the formula (28) one can find the following vector generalizations of the systems (17) and (27):

$$\begin{aligned} \begin{cases} u_t = u_{xxx} + 2 \langle u; v \rangle u_x + 2 \langle u; v_x \rangle u + \langle u; v \rangle^2 u + \\ 2 \langle v; v \rangle u_x + \langle u; u \rangle u + 2 \langle u; v \rangle v + \langle v; v \rangle u + u; \end{cases} \\ \begin{cases} v_t = v_{xxx} - 2 \langle u; v \rangle v_x - 2 \langle v; u_x \rangle v - \langle u; v \rangle^2 v \\ 2 \langle u; u \rangle v_x + \langle v; v \rangle v + 2 \langle u; v \rangle u + \langle u; u \rangle v + v; \end{cases} \end{aligned}$$

$$\begin{aligned} u_t &= u_{xxx} + 3 \langle u; v \rangle u_{xx} + 3u \langle u_x; v_x \rangle + 3 \langle u; v \rangle_x u_x + 3 \langle u; v \rangle^2 u_x; \\ v_t &= v_{xxx} + 3 \langle u; v \rangle v_{xx} + 3v \langle u_x; v_x \rangle + 3 \langle u; v \rangle_x v_x + 3 \langle u; v \rangle^2 v_x; \end{aligned}$$

where u and v are N -dimensional vectors and $\langle \cdot \rangle$ is a scalar product. Both of them can be linearized just as in the scalar case:

$$U = u \exp^Z \langle u; v \rangle dx; \quad V = v \exp^Z \langle u; v \rangle dx \quad (29)$$

In contrast with (16), (17), equations (9) and (11) are related to linear systems

$$U_t = U_{xx} + c_1 U_x + c_2 V_x + c_3 U; \quad V_t = V_{xx} - k_1 V_x - k_2 U_x - k_3 V \quad (30)$$

via local differential substitutions.

Equation (9): For $c_1 = c_2 = c_3 = 0$ the substitution is given by (3). In the general case such a substitution is defined by

$$u = \frac{U_x}{(U+V)} + \frac{(c_1 - k_2)U + c_2 V}{2(U+V)}; \quad v = \frac{V_x}{(U+V)} + \frac{(c_1 - c_2)V + k_2 U}{2(U+V)};$$

where the constants in (30) and (9) satisfy the following conditions

$$k_1 = c_1; \quad k_3 = c_3 = \frac{c_1(c_1 - c_2 - k_2)}{4};$$

$$2 = k_2(c_1 + c_2); \quad 2 = c_2(c_1 + k_2); \quad 2 = c_1 k_2 c_2:$$

Equation (11): The substitution is of the form

$$u = \frac{c_2 U_x}{V} + \frac{c_2^2}{2}; \quad v = V_x = V + \frac{c_1}{2}:$$

The relations between constants are the following

$$k_1 = c_1; \quad k_2 = c_2; \quad k_3 = 0;$$

$$2 = c_1^2 - 2c_2^2 + 2c_3; \quad 2 = c_1 c_2^2; \quad 4 = c_2^2 (2c_1^2 + c_2^2 - 2c_3):$$

Equation (18), (19): These equations are related to triangular systems. Equations (19) have been obtained in [17]. The substitution

$$u = \frac{U_x}{2U}; \quad v = \frac{V}{U};$$

found first by Marichin [15], links equations (19) with

$$U_t = U_{xx} + 2V^2; \quad V_t = V_{xx} + V:$$

The last system is linear in the following sense. To find V we need to solve a linear equation. For a given function V , the function U satisfies a linear equation with variable coefficients.

Similarly the following substitution

$$u = \frac{1}{3} U^{-2} V^{-1} U_x; \quad v = U^{-1} V;$$

reduces (18) to

$$U_t = U_{xx} - 2V^{-1} V_x U_x + 3U + 3V^3; \quad V_t = V_{xx}:$$

3. Master-symmetries. For all equations (8)–(22) we have found all symmetries of order less or equal than four using the computer program LiePde [20]. It turns out that in many of the equations have symmetries depending on t and x explicitly. For example, computing for equation (16) the general symmetry of n^{th} order of the form

$$u = P(t)u_{nx} + \dots; \quad v = P(t)v_{nx} + \dots$$

the polynomial $P(t)$ is an arbitrary polynomial of degree n .

It is well known that symmetries which are linearly depending on t and x are closely related to local master-symmetries [14]. The equations (8)–(11), (15)–(17) have such symmetries. To obtain the master-symmetries one has to simply put t equal to zero in these time-dependent symmetries.

The resulting master-symmetry is of the form

$$\begin{aligned} u_t &= 2x (u_{xx} + F(u; v; u_x; v_x)) + f(u; v; u_x; v_x); \\ v_t &= 2x (v_{xx} + G(u; v; u_x; v_x)) + g(u; v; u_x; v_x); \end{aligned} \quad (31)$$

where F and G are the right hand sides of the corresponding equation (1) and f and g are given by the following list

$$\begin{aligned} (8) : f &= u^2 + 3uv + 4u_x; & g &= v^2 + 3uv - 4v_x; \\ (9) : f &= 2u^2 + 4uv + 3u_x; & g &= 2v^2 - 4uv - 3v_x; \\ (10) : f &= 4uv + 5u_x; & g &= v^2 + 4u - 5v_x; \\ (11) : f &= 4uv + 3u_x; & g &= 2v^2 - 2u - 3v_x; \\ (15) : f &= 2u^2v + 2v + 3u_x; & g &= 2uv^2 + 2u - 3v_x; \\ (16) : f &= 2u^2v - 2xu + 2u_x; & g &= 2v^2u + 2xv - 2v_x; \\ (17) : f &= 2u^2v + 2v + 2u_x; & g &= 2uv^2 - 2u - 2v_x; \end{aligned}$$

Acknowledgement

The first author (V.S.) was supported, in part, by RFBR grant 99-01-00294, EPSRC grant GR/L99036 and LMS (FSU) grant 5324. He is grateful to the Queen Mary & Westfield College (Univ. of London) for its hospitality. The second author (T.W.) thanks the Symbolic Computation Group at the University of Waterloo for its hospitality during the authors sabbatical in the fall of 1998.

References

- [1] Mikhailov A.V., Shabat A.B., Integrability conditions for systems of two equations $u_t = A(u)u_{xx} + B(u; u_x)$: I., *Theor. Math. Phys.*, 62 (2) 163-185 (1985).
- [2] Mikhailov A.V., Shabat A.B., Integrability conditions for systems of two equations $u_t = A(u)u_{xx} + B(u; u_x)$: II., *Theor. Math. Phys.*, 66 (1) 47-65 (1986).
- [3] Shabat A.B., Yamilov R.I., On a complete list of integrable systems of the form $iu_t = u_{xx} + f(u; v; u_x; v_x)$, $iv_t = v_{xx} + g(u; v; u_x; v_x)$. Preprint BFAN Ufa, 28 pages (1985).
- [4] Mikhailov A.V., Shabat A.B., Yamilov R.I., Extension of the module invertible transformations. Classification of integrable systems. *Commun. in Math. Phys.*, 115 1-19 (1988).
- [5] Mikhailov A.V., Shabat A.B., Yamilov R.I., Symmetry approach to classification of nonlinear equations. Complete lists of integrable systems. *Russian Math. Surveys*, 42 (4) 1-63 (1987).

- [6] Calogero F., Why Are Certain Nonlinear PDE's Both Widely Applicable and Integrable?, in book "What is integrability?" Springer-Verlag (Springer Series in Nonlinear Dynamics), 1-62 (1991).
- [7] Borovik A.E., Popkov V.Yu., Robuk V.N., Generation of nonlinear structures in exactly solvable dissipative systems, Dokl. AN SSSR, 305 (4) 841-843 (1989).
- [8] Sokolov V.V., On the symmetries of evolution equations, Russian Math. Surveys, 43 (5) 165-204 (1988).
- [9] N.Kh. Ibragimov and A.B. Shabat, Evolutionary equations with nontrivial Lie-Backlund algebra. Func. Anal. Appl., 14 (1) 19-28 (1980).
- [10] Fokas A.S., A symmetry approach to exactly solvable evolution equations, J. Math. Phys., 21 (6) 1318-1325 (1980).
- [11] Sokolov V.V. and Shabat A.B., Classification of integrable evolution equations Soviet Sci. Reviews, Section C, 4 221-280 (1984).
- [12] Mikhailov A.V., Shabat A.B., Sokolov V.V., Symmetry Approach to Classification of Integrable Equations, in the book "What is integrability?", Springer-Verlag (Springer Series in Nonlinear Dynamics), 115-184 (1991).
- [13] N.Kh. Ibragimov and A.B. Shabat, Infinite Lie-Backlund algebras. Func. Anal. Appl., 14 (4) 313-315 (1980).
- [14] Fuchssteiner B., Mastersymmetries, higher order time-dependent symmetries and conserved densities of nonlinear evolution equations. Prog. Theor. Phys., 70 (6) 1508-1522 (1983).
- [15] Marihin V.G., Private communication, (1998).
- [16] Sokolov V.V., Svinolupov S.I., Wolf T. On linearizable evolution equations of the second order, Phys. Lett. A, 163 415-418 (1992).
- [17] O'lvér P.J., Sokolov V.V., Integrable Evolution Equations on Associative Algebras, Commun. in Math. Phys., 193 245-268 (1998).
- [18] F. Calogero and W. Eckhaus, Nonlinear evolution equations, rescalings, model PDEs and their integrability. I, Inverse Problems, 3 229{262 (1987).
- [19] Wolf, T., The program CRACK for solving PDEs in General Relativity, in Relativity and Scientific Computing, Eds. F.W. Hehl, R.A. Puntigam, H. Ruder, Springer, 241{257 (1996).
- [20] Wolf, T., An efficiency improved program LIEPDE for determining Lie-symmetries of PDEs, Proceedings of Modern Group Analysis: advanced analytical and computational methods in mathematical physics, Catania, Italy October 1992, Kluwer Academic Publishers, 377{385 (1993)